

1973/164  
Copy 3


Restricted until after publication.  
Manuscript submitted for publication  
to: Bull. *Geol. Surv. Aust.* *Geol.*



DEPARTMENT OF  
MINERALS AND ENERGY

# BUREAU OF MINERAL RESOURCES, GEOLOGY AND GEOPHYSICS

RECORD 1973/164



THE SILESIAN IN AUSTRALIA - A REVIEW

by

E.C. DRUCE AND P.J. JONES

The information contained in this report has been obtained by the Department of Minerals and Energy as part of the policy of the Australian Government to assist in the exploration and development of mineral resources. It may not be published in any form or used in a company prospectus or statement without the permission in writing of the Director, Bureau of Mineral Resources, Geology and Geophysics.

BMR  
Record  
1973/164  
c.3

RECORD 1973/164

THE SILESIAN IN AUSTRALIA - A REVIEW

by

E.C. DRUCE AND P.J. JONES

THE SILESIAN IN AUSTRALIA - A REVIEW

by

E. C. Druce & P.J. Jones

- 
- I INTRODUCTION
- II VISÉAN-NAMURIAN BOUNDARY
- III GEOLOGICAL SUMMARY
- Eastern Australia
- New South Wales
- Queensland
- Central to Western Australia
- IV PALAEOONTOLOGICAL SEQUENCES
- Brachiopods
- Plants
- Spores and Pollens
- V CORRELATION
- VI CARBONIFEROUS - PERMIAN BOUNDARY
- VII TECTONICS
- VIII CONCLUSION

# ABSTRACT

The Silesian in Australia is well defined by faunas at the Viséan - Namurian boundary below, and the Sakmarian above. Between these limits the geology is complicated by tectonism and glaciation, and correlation is hampered by faunal and floral provincialism.



# THE SILESIAN IN AUSTRALIA - A REVIEW

by

E.C. Druce and P.J. Jones \*

## I. INTRODUCTION

The Upper Carboniferous (Silesian) is poorly represented in Australia. Most sequences are known from eastern Australia, where both marine and non-marine strata are represented. In central and western Australia widespread occurrences of Silesian rocks have been reported. In general the Australian Silesian sequence contains glaciogene sediments (Crowell & Frakes, 1971).

Carboniferous rocks were first described from Australia by Strzelecki (1845) from the Hunter Valley, New South Wales, from a sequence now referred to the Silesian. Most of his fossil collection was described by Morris (Lonsdale described the polyzoans and corals), who concluded that "the deposits containing them may probably belong to that division of the Palaeozoic series, usually termed carboniferous". Strzelecki's (1845, pp. 88-9) observations of the sequence exposed on the steep banks of the River Karuah, west of Stroud, are remarkably accurate, in terms of present-day interpretations. There is no doubt that the tripartite sequence ("siliceous breccia, slaty blue argillaceous rock, and sandstone and conglomerate") which he described as overlying greenstones (= Nerong Volcanics), refer to the Karuah, Booral, and Isaacs Formations of Campbell (1961). The rich fauna of gastropods, ostracods, *Spirifer*, and fish remains, which Strzelecki described from Booral, probably belong to the *Levipustula levis* Zone of the Booral Formation, (Campbell, 1961). Strzelecki suggested that the "sandstones and conglomerates" of his uppermost unit in the Booral district, (i.e. the Isaacs Formation) "at least in a mineralogical point of view" are similar to those of Raymond Terrace (i.e., Lower Permian Dalwood Group and Branxton Formation of the Maitland Group). It is to his credit that he was aware that such a lithological correlation lacked palaeontological control, "for the absence of natural sections precludes the discovery of the fossils".

Considerable information on the Silesian has been published by various authors; a recent synthesis of the Carboniferous System has been given by Brown et al., (1968). A detailed account of the sequence and affinities of eastern Australian Carboniferous invertebrates has been provided by Campbell & McKellar (1969).


The present review briefly outlines the geology and fauna of the Silesian of the Australian continent. Not every Silesian outcrop has been considered in detail; rather we have used generalized or typical sections to represent each of the major areas of sedimentation.

These sections are shown on Fig 2, and the localities appear on the generalized map of the Silesian (Fig 1).

We thank Dr K.S.W. Campbell (Australian National University) and our BMR colleagues Drs Elizabeth Kemp and J.M. Dickins for discussions on various aspects of this paper.

\* Published with the permission of the Director Bureau of Mineral Resources, Geology, and Geophysics, Canberra.

Figure 1.



## II. THE VISÉAN - NAMURIAN BOUNDARY.

Palaeontological data are not continuous in those sections which appear to cross the boundary. Campbell & McKellar (1969) have reviewed the evidence and considered that the Margininugus barringtonensis Assemblage Zone may span this interval. Their evidence was based on the coral fauna and in addition they pointed out that "Lissochonetes, which is found only in the upper part of the zone in one area, is not reliably reported from pre-Namurian rocks anywhere in the world" (op. cit. p. 88). Subsequent faunas are separated by either a stratigraphic break or a disconformity (op. cit. p. 103).

This fauna is extensive in eastern Australia, being known from central Queensland (Baywulla Formation, Yarrol Basin; McKellar, 1965, 1967) to the central coast of New South Wales (uppermost Copeland Road and lowermost Buckets Gap Formations; Campbell & McKelvey, 1972). Other Namurian faunas in New South Wales include the sequence with Cravenoceras at Kempsey (Campbell, 1962), and possibly the Taree Limestone (Pickett, 1966).

In eastern Australia sedimentation continued from the Viséan into the Namurian and Westphalian. In Western Australia there is generally a hiatus between the marine Viséan and the dominantly non-marine Upper Carboniferous. In the Bonaparte Gulf Basin Veevers & Roberts (1968, p. 108) considered that the uppermost beds of the Point Spring Sandstone are Namurian. The Point Spring Sandstone contains Anthracospirifer milliganensis, which is close to A. bisulcatus Sowerby sensu stricto, and its close allies from the Viséan and Namurian of Great Britain and Belgium (Thomas, 1971, p. 24). Roberts (1971, p. 22) recognized a subsequent brachiopod fauna, the Echinoconchus gradatus Fauna, from sample 458/12. Although this species occurs in the Viséan of eastern Australia it is considered to be Namurian in the Bonaparte Gulf Basin because it is later than A. milliganensis. The Echinoconchus gradatus Fauna has also been found in Bonaparte No. 1 Well (206-210 m) and Kulshill No. 1 Well (2042-2048 m). The Namurian age is confirmed by the presence of foraminifers indicating the Viséan-Namurian boundary in Bonaparte No. 1 Well in the interval 204 - 207 m (Mamet & Belford, 1968, p. 344).

In the Canning Basin the top of the Carboniferous Unit A of Bischoff has yielded conodonts of either latest Viséan or earliest Namurian age in Barlee No. 1 Well (Bischoff, fide Jones et al., in press). This unit is unconformably overlain by the Grant Formation, a non-marine unit which may be as old as Westphalian.

## III GEOLOGICAL SUMMARY

### Eastern Australia

#### New South Wales

The Upper Carboniferous in New South Wales extends as a U-shaped belt from the Queensland border as far south as Newcastle. Sections in the youngest Carboniferous are developed in the southern part of the Lower and Middle Hunter Valley and in the Stroud-Gloucester area. Three distinct facies are recognized (Engel, 1965): the Kuttung, Myall, and Burindi facies.

Figure. 2.

EUROPEAN STAGE		WESTERN AUSTRALIAN BRACHIOPOD ZONES	WESTERN AUSTRALIA					QUEENSLAND			NEW SOUTH WALES				EASTERN AUSTRALIAN MARINE INVERTEBRATE ZONES				
			BONAPARTE GULF BASIN			CANNING BASIN	CARNARVON BASIN	NORTH QUEENSLAND <small>(Generalised sequence Western part)</small>	YARROL TROUGH		WARWICK - GOONDIWINDI	TAMWORTH TROUGH							
			NORTH-WESTERN PLATFORM	BASINAL AREA	SOUTH-EASTERN PLATFORM				ROCKHAMPTON	CANIA		CANNINDAH CREEK-YARROL	BALICKERA-CLARENCE-TOWN	BOORAL		GLOUCESTER BARRINGTON			
SAKMARIAN (in part)			52	52	52	28'9	31	54,55	19,30	28,129	35	25,1151	43,24	7	12				
<div>STEPHANIAN</div> <div>WESTPHALIAN</div> <div>NAMURIAN</div>				KULSHILL FM					DINNER CREEK CONGLOM	YOULAMBIE CONGLOM	BURNETT FM								
				KEEP INLET BEDS															
<div>Polynoid key</div> <div>P = <i>Potoneisporites</i></div> <div>U = unnamed microflora</div> <div>S = <i>Grandsport</i></div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
<div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div> <div><i>Echinocanthus gradolus</i> fauna</div>																			
												</							

To accompany Record 1973/164

AUS 4/20

The Kuttung facies is essentially a non-marine volcanic and glacial sequence, mainly Late Carboniferous (Engel, 1966) and laterally equivalent to the Myall facies.

The Myall facies is a mixed sequence of marine and non-marine arenite and mudstone with beds of glacial outwash material and volcanics flows.

The Burindi facies is a marine mudstone-arenite sequence which is virtually confined to the Lower Carboniferous according to Engel (1966), and is distinguished from the Myall facies in that it lacks glacial outwash sediments and contains little, if any, volcanic rocks and terrigenous material. However, Campbell & McKelvey (1972) consider that the Burindi facies is extensively developed in the Upper Carboniferous and that it is temporally equivalent to both the Kuttung and Myall facies. Some confusion has arisen because these facies names were originally used in a stratigraphic sense and Roberts & Oversby (1973) suggest that the terms should be abandoned. Though we agree with Roberts and Oversby that they should not be used in a stratigraphic sense, they are used here strictly as a convenient way of delineating different lithological sequences of the same age.

Campbell & McKelvey (1972) illustrate all three facies as being present during the earliest Namurian (Margininugus barringtonensis time). The Kuttung facies consists of volcanic rocks separated from the Myall facies by a north-northwest shoreline in the Port Stephens - Dungog area. Farther offshore, to the east, in the area of Myall Lake and Forster, the Burindi facies was developed. This sedimentary pattern continued through the Namurian (Oriocrassatella time) and into the Westphalian (Levipustula levis time and later), although in the region where the Burindi facies was probably developed after Levipustula levis time, no information is available.

The Kuttung facies is well developed in the area north of Raymond Terrace (Rattigan, 1967a). Part of the sequence, the Wallaringa Formation and the overlying Gilmore Volcanics, is Lower Carboniferous. The Upper Carboniferous is represented by the Kings Hill Group, comprising the Balickera Conglomerate, the Italia Road Formation, and the Grahamstown Lake Formation (Rattigan, op. cit.). The Balickera Conglomerate consists of about 420 m of polymictic boulder conglomerate with two tuff and ignimbrite horizons. Plant fossils are indeterminate. The Italia Road Formation, corresponding to part of the "Glacial Stage" of Osborne (1922), contains 360 m of massive lithic arenite with interbeds of fine laminated sandstone, shale, and poor coal (Rattigan, 1967a). The formation yields abundant plant material (Rattigan, 1967a) and spores (Playford & Helby, 1968; Helby, 1969b). The Grahamstown Lake Formation, corresponding to the "Main Glacial Beds" (Osborne, 1922), consists of 363 m of laminated red, green, and mauve claystone, siltstone, fine sandstone with zeolitic sandstone, and pebble conglomerate (Rattigan, 1967a). The plant Rhacopteris is present. Both Rattigan (1967a) and Playford & Helby (1968) considered that the Grahamstown Lake Formation is conformable on the Italia Road Formation, however, Jones et al., (in press) have suggested a possible stratigraphic break between the two as an attempt to reconcile the differences between the correlations of Norris, and unpublished palynological data of Helby.



in press

Helby (vide Jones et al.) considers the microflora of the Isaacs Formation to be intermediate between that of the Italia Road Formation and that of the Seaham Formation. The latter has been lithologically correlated with the Grahamstown Lake Formation. However, no floras are known from the Grahamstown Lake Formation (Helby, pers. comm.) and this break should be considered as questionable.

The Myall facies has been described from the Barrington District by Campbell & McKelvey (1972), where the Silesian is represented by the Buckets Gap and the McInnes Formations. The Buckets Gap Formation consists of up to 610 m of mudstone, siltstone, and fine sandstone with interbedded lithic sandstones. The unit is fossiliferous, yielding brachiopods, corals, polyzoans, pelecypods, gastropods, and a trilobite (Campbell & McKelvey, 1972). The McInnes Formation is about 620 m thick, consisting of medium to coarse sandstone and fine conglomerate with interbedded dark siltstone containing plant fossils. The upper part of the unit consists of a volcanic breccia which Campbell & McKelvey (1972, p. 17) considered to be a separate unit, but the top of the sequence is faulted, and it is not named.

The Burindi facies is developed farther east in the vicinity of Bulahdelah and Port Stephens (Engel, 1962) and is conspicuously lacking in glacially derived material. This facies, referred to the Crawford Formation, is about 2440 m thick in the type section and consists of lithic arenite, friable mudstone, and small amounts of conglomerate and tuff beds (Engel, 1962). Fossils are rare and occur in bands; they include brachiopods, pelecypods, and gastropods with occasional plant fragments.

On the eastern limb of the Stroud - Gloucester Syncline near Booral, Campbell, (1961) has divided the Crawford Formation into the Karuah, Booral and Isaacs Formations; this locality is probably the one described by Strzelecki.

Recent studies in the southern Sydney Basin (Gostin & Herbert, 1973) have demonstrated the presence of terrestrial valley-fill sediments (Talaterang Group) which define a palaeodrainage system active during the Late Carboniferous. The basal unit - the Pigeon House Creek Siltstone - contains a microflora closely comparable with that recorded from the Seaham Formation in the Hunter Valley (Helby & Herbert, 1971).

### Queensland

The scattered outcrops of Upper Carboniferous rocks in Queensland represent marine, mixed and non-marine facies, although nowhere are they seen as an interfingering sequence as in New South Wales. The Carboniferous outcrop consists of two linear belts running north-south. The eastern belt consists of the Yarrol Trough, which is probably a northward extension of the Tamworth Trough. The western outcrop belt consists of the Drummond, Bundock, Clarke River, Burdekin, and Hodgkinson Basins (Brown et al., 1968, fig. 6, 2). Silesian rocks are rare in the western belt but are relatively more common in the Yarrol Trough.

Terrestrial sediments and volcanics are common in the western outcrop belt to the west and south of Townsville. In most areas the sequence consists of a basalt unit of andesite or rhyolite overlain by sediments, and an upper volcanic unit, with a total thickness of about 3000 m.

The sediments consist of conglomerate, sandstone, subgreywacke, siltstone, and carbonaceous shale. Fossils include palaeoniscid fish and plant remains including cf. Rhacopteris. Because of extensive faulting, uplifting, and folding many formation names have been proposed; however, the similarity of lithology and fossils suggests that all the formations are part of a single depositional cycle in the Bundock, Clarke River, and Burdekin Basins. The Drummond Basin is the southerly extension of this outcrop belt and deposition consisted of glacio-lacustrine sediments referred to the Joe Joe Formation, which consists of about 600 m of lithic conglomerate and sandstones with striated polymictic pebbles, grading up into fine sandstone and siltstone, interlaminated siltstone and claystone, carbonaceous beds, and thin coaly beds. Carboniferous plants, Cardiopteris, and spores are the only fossils. The sequence was interpreted as fluvioglacial, with tills in the basal part; the upper part is mainly lacustrine (Dickins & Malone, 1973, p. 112). Equivalent sediments are developed in the Galilee Basin to the west (Olgers, 1973.).

The Hodgkinson Basin forms the northern extremity of the outcrop belt; the Nychum Volcanics have yielded a mixed Glossopteris - Cardiopteris flora and a Rb/Sr age dating of  $294 \pm 9$  m.y. (Black et al., 1972) suggesting a late Carboniferous age. This has been challenged by Balme (1973) who stated that the real interest of this plant fossil assemblage "is not that it is 'mixed', but that it provides the first clear Australian record of cardiopterid fronds in association with diverse elements of the Glossopteris flora". Such an association is known from India, South Africa, and Brazil, in strata which most Australian palaeontologists and stratigraphers regard as Early Permian (see full discussion by Balme, op. cit.).

The easternmost deposits in Queensland are in the Yarrol Basin which probably extends south to join the Tamworth Trough in New South Wales. The earliest Silesian is represented by part of the Baywulla Formation; McKellar (1967) considers that his Fauna G (= Marginirugus barringtonensis Zone) straddles the Viséan - Namurian boundary.

Unconformably overlying the interbedded limestone and medium grained greywackes of the Baywulla Formation are up to 580 m of subgreywacke, conglomerate, and argillite referred to the Branch Creek Formation (McKellar, o. cit.). The fauna consists of polyzoans, brachiopods (including Levipustula levis), coelenterates, molluscs, and a trilobite, Australosutura gardneri (Mitchell).

The upper part of the sequence consists of the Poperima Formation, up to 400 m of mudstone and shale, and the Rands Formation comprising up to 1500 m of subgreywacke with interbeds of conglomerate, shale and mudstone. The fauna of both these formations consists of polyzoans, brachiopods, and molluscs. The Rands Formation is conformably overlain by Lower Permian sediments and McKellar (1967, p. 16) considered that the Boiling Creek Group, which comprises these formations, to include the Westphalian and Stephanian because it overlies, albeit with slight unconformity, the late Viséan-early Namurian Baywulla Formation and is overlain by the Lower Permian (Sakmarian) Burnett Formation.

In the northern part of the Yarrol Basin the Silesian is represented by the 2150 m thick Neerkol Formation (Kirkegaard et al., 1970), which disconformably overlies the Rockhampton Group. It is dominated by mudstone and fine grained sandy siltstone with minor amounts of sandstone, conglomerate, and limestone (Kirkegaard et al., op. cit.). The fauna consists of polyzoans, brachiopods, and molluscs (Fleming, 1969) and is similar to that in the central part of the trough; the formation is equivalent to the Poperima and Rands Formations (McKellar, 1967) and possibly to the basal Permian (Sakmarian) Burnett Formation (Fleming, 1969).

In southern Queensland in an area between the Yarrol and Tamworth Troughs two exposures in areas of structural complexity have yielded Silesian faunas. In the Murgon area, 160 km northwest of Brisbane, limestones of the 'Wondai Series' (thought to be part of the Neranleigh Fernvale Group) are exposed in a faulted block. Palmieri (1969) has described a Silesian conodont fauna from them which ranges in age from late Namurian to basal Westphalian (Jones et al., in press). At Mount Barney 100 km south southwest of Brisbane about 2000 m of sediments occurs; a mudstone 600 m above the base has yielded the Levipustula levis fauna (McKellar, 1967).

#### Central and Western Australia

No undoubted Silesian marine sediments are known from Central and Western Australia. However, floral evidence suggests possible widespread non-marine sedimentation in the area, usually including glaciogene sediments.

Evans (1969) reported his Stage 1 microflora, which he regarded as Late Carboniferous, from the Lake Phillipson Bore in the northern part of South Australia (Balme, 1957; Ludbrook, 1961), the southern part of the Northern Territory, and parts of the Murray Basin. This microflora has since been reported from Tasmania, from Hellyer Gorge in the basal unit of the Wynard Tillite of the Quamby Group (Evans vide Banks & Clarke, 1973). Gulline's (1967, p. 332-3) discovery of 'Rhacopteris' ovata in these sediments provided the first proof of their Late Carboniferous age.

In Western Australia Silesian sediments are known from the Bonaparte Gulf, Canning and possibly the Carnarvon Basins.

In the Bonaparte Gulf Basin the late Viséan - early Namurian Point Spring Sandstone is unconformably overlain by the Border Creek Formation. In outcrop the Border Creek Formation has yielded Phyllothea like plants (Veevers & Roberts, 1968, p. 110) suggesting a Late Carboniferous or Permian age. In Pelican Island No. 1 well the Potonieisporites microflora is present in the unit but in Bonaparte No. 2 Well the Border Creek yielded a Tournaisian - Viséan microflora (Jones et al., in press). This anomalous result suggests that either the microflora has been recycled, or the sediments have been incorrectly identified.



In the Canning Basin the subsurface Anderson Formation was considered on conchostracan evidence to be Upper Carboniferous (Öpik in McWhae et al., 1958; Balme, 1960). However conodont evidence from Yulleroo No. 1 Well and Grant Range No. 1 well suggest that it is confined to the Lower Carboniferous (Bischoff, fide Jones et al., 1973).  
*(in press)*

The Grant Formation unconformably overlies the Anderson Formation; it consists of up to 3000 m of tillite, sandstone, and shale and is, in large part, Permian in age. However, the lower part contains Late Carboniferous microfloras (Balme, fide Jones et al., 1973).  
*(in press)*

The Harris Sandstone in the Carnarvon Basin is of indefinite age, but plants suggest that it is Upper Carboniferous (Krausel, 1961).

#### IV PALAEONTOLOGICAL SEQUENCES

The Australian Silesian faunas are markedly provincial (Brown et al., 1968), and faunal links between Eastern Australia and the northern hemisphere were broken, probably during the Namurian (Campbell & McKellar, 1969). The endemic fauna and flora, including brachiopods, plants, and spores, have been used extensively to subdivide the Late Carboniferous.

##### Brachiopods

In New South Wales (Tamworth Trough) and Queensland (Yarrol Trough) sequences of marine brachiopod faunas have been described from the Carboniferous (McKellar, 1967; Campbell & McKellar, 1969; Campbell & McKelvey, 1972).

McKellar recognized three successive Silesian faunas in the Yarrol Trough, Faunas G-I (1967, p. 28). Fauna G occurs in the Baywulla Formation; the index species are Marginirugus barringtonensis (Dun) and Tylothyris (?) multicostatus Maxwell. Fauna H includes Werriea regina McKellar, Lissochonetes montinis McKellar, Neospirifer campbelli Maxwell, Alispirifer laminosus Campbell, and Kitakamithyris booralensis (Campbell), a fauna with no similarity to Fauna G. It occurs in Unit B of the Branch Creek Formation and is the first appearance of Levipustula levis Maxwell and Beecheria aff. patagonica Amos.

The overlying fauna, I contains two diagnostic species, Neospirifer pristinus Maxwell and Spinuliplica spinulosa Campbell, as well as Levipustula levis. This fauna is characteristic of the Poperima Formation.

The overlying Rands Formation contains some elements of Fauna I, including Levipustula levis, Spiriferellina neerkolensis Maxwell and Beecheria aff. patagonica Amos; Punctospirifer ambiguus Maxwell occurs in the upper part. McKellar made no attempt to designate these later faunas.

The faunal sequence in New South Wales is closely similar, although Campbell & McKellar (1969) have merged Fauna's H and I into a Levipustula levis Zone. Campbell & McKelvey (1972) have added the Oricrassatella fauna, which is intermediate between Fauna G and H, to the L. levis Zone; they recognized an interval which they call the Oriocrassatella Levipustula levis time (p.27). McKellar's Fauna G is equivalent to the Marginirugus barringtonensis Zone (Campbell & McKellar, 1969).

A post-Levipustula levis Zone fauna is recognized in the Tamworth Trough, but is dissimilar to faunas of the same age in the Yarrol Trough. No zones or faunal groupings have been erected for these faunas; they are generally considered to be Carboniferous because they show strong similarity to underlying faunas and little, if any, to the overlying Permian faunas.

### Plants

The Silesian in Australia has also been defined florally; it is characterized by the Rhacopteris flora and is underlain by the Lepidodendron flora of generally lower Carboniferous age (Walkom, 1920). The upper limit of the Silesian has been taken at the boundary between the Rhacopteris and Glossopteris floras.

Recently Rigby (1973) has reviewed the floras and extended the range of Lepidodendron upward to a horizon considered to be equivalent to the Westphalian-Stephanian boundary. The range of Rhacopteris is unchanged and this interval is called the Pseudorhacopteris flora (Rigby, 1973, p.5). Overlying this flora is the Gondwanidium flora, which is succeeded by the Gangamopteris/Glossopteris flora of Permian and questionably latest Carboniferous age. There is still considerable discussion on the taxonomy of Australian Silesian plants, and Norris (*fide* Jones et al., in press 1973) considers that Lepidodendron does not extend into the Silesian.

### Spores and Pollen

Many petroleum exploration wells have intersected Silesian strata in Australia. The sequence has been divided on palynological data and several schemes have been proposed. The first detailed discussion of Australian pre-Tertiary floras was by Balme (1964), who reported little information from the Carboniferous except that the early Carboniferous was dominated by Lycospora. In the Permian he recognized two assemblages in his Striatites Microflora, the Nuskoisporites and Vittatina Assemblages. The lower part of the Nuskoisporites Assemblage is now considered to be Silesian because it first appears within the range of the Rhacopteris Flora (Evans, 1969). This interval is Stage 1 of Evans (op. cit., p. 43) in which Rhacopteris is associated with monosaccate pollen; and with the spores Retusotriletes diversiformis, Punctatisporites gretensis and species of Verrucosisporites in the Seaham Formation of New South Wales. This is the Potonieisporites Microflora of Helby (1969).

Helby (1969b, fig. 30) recognized an earlier Grandispora Microflora, first described by Playford & Helby (1968), which is long ranging and extends down into the Wallaringa Formation of late Viséan age (Jones et al., in press). Helby (*fide* Jones et al., in press) now recognizes an informal microflora between his Potonieisporites and Grandispora Microfloras.

The sequence of microfloras above the Lycospora Microflora and beneath the upper Nuskoisporites Microflora is, in ascending order: Grandispora, unnamed, and Potonieisporites (= Stage 1 of Evans, and lower Nuskoisporites of Balme).

## V CORRELATION

The problems of recognizing boundaries are amplified when the correlation of the Australian Silesian is attempted.

Campbell & McKellar (1969), in considering the macrofaunas, noted that there is little evidence of faunal relationships during the Namurian apart from the occurrence of Cravenoceras in New South Wales. The provincial nature of the fauna is due to the establishment of a cool to cold water fauna. There is a close relationship between the Australian and Argentinian faunas. Species in common include the brachiopods Levipustula levis Maxwell, Alispirifer laminosus Campbell and Beecheria aff. patagonica Amos, the trilobite Australosutura gardneri (Mitchell), and the polyzoans Fistulamina frondescens Crockford, Penestella anodosa Campbell, Polypora neerkolensis Crockford, P. septata Campbell, and Rhombopora bifurcata Campbell (Amos & Sabattini, 1969; Campbell & McKellar, 1969). The Argentinian fauna is much closer to Australian faunas than it is to faunas from the northern part of South America (Campbell & McKellar, op. cit.).

Although the Australian Levipustula levis fauna has twelve genera which also occur overseas, Campbell & McKellar (1969, p.105) concluded that it was not possible to draw any safe conclusions with regard to correlation. The Levipustula levis fauna has always been considered to be Westphalian. This conclusion is based on the range of the brachiopod genus Lissochonetes, which first appears in the Westphalian (W3a) of Belgium; its close relative, Eolissochonetes, occurs in the Pennsylvanian Morrowan and Desmoinesian in North America (McKellar, 1965). Campbell (1961) noted that Levipustula occurs in the Westphalian B and basal Westphalian C in the Franco-Belgian coalfield and in Britain.

Böger & Fiebig (1963) also described three species of Levipustula from the Silesian of west Germany, the combined range of which is from Westphalian A to the Westphalian B/C boundary (Flöz Ägir). The oldest known species is L. piscariae (Waterlot), from the oberes Sarnsbank-Nieveau, i.e., at the Namurian/Westphalian A boundary. Although this evidence strongly supports a Westphalian age for the Levipustula levis fauna the picture is complicated by the presence of the levis fauna beneath the occurrence of Cravenoceras in the Kempsey district (Lindsay, 1969); however, the area is structurally complex.

The only occurrence of conodonts is in a structurally deformed and metamorphosed area 160 km northwest of Brisbane, Queensland (Palmieri, 1969). The fauna contains European and North American species of Namurian and Westphalian age, indicating that marine sedimentation was taking place during this time in eastern Australia.

The palynological evidence is tenuous. There is no direct correlation between the Grandispora and Potonieisporites microfloras and those of the northern hemisphere. The upper limit is set at the incoming of Evans' (1969) Stage 2, equivalent to the upper of Balme's (1964) Nuskoisporites microflora.

The flexibility of correlation of the microfloras is shown by the various ages assigned to the Potonieisporites microflora: post-Carboniferous (Helby, 1969 a,b) to Namurian (Helby & McClung, 1973). This microflora occurs above beds yielding the post-Levipustula levis fauna within the Isaacs Formation (Helby, 1969b).

Palynology has been used to correlate the subsurface Silesian but the information remains unpublished.

## VI CARBONIFEROUS/PERMIAN BOUNDARY

The provincial nature of the fauna and flora in the late Carboniferous and early Permian makes the recognition of the Carboniferous/Permian boundary extremely difficult.

Traditionally the boundary has been taken at the incoming of the pelecypod Eurydesma and of the Glossopteris flora and the extinction of the Rhacopteris flora (Evans, 1969; Jones et al., in press). However, Runnegar (1969) has pointed out that Eurydesma and Rhacopteris overlap in sections at Cranky Corner, 16 km north of Gosford (Seaham Formation) and at Cania in the Yarrol Trough, Queensland (Youlambie Conglomerate).

Palynologists have endeavoured to recognize a change in the microfloras similar to that in the macrofloras. Balme (1964) tentatively suggested that this change could be the incoming of the Nuskoisporites Assemblage but, as Evans (1969, p. 42) pointed out, the incoming of this assemblage is within the upper range of Rhacopteris.

Evans (op. cit.) considered that this interval, between the introduction of monosaccate pollen and the appearance of the Glossopteris Flora, is Late Carboniferous.

Palynologically the Carboniferous/Permian boundary has been taken at the base of Evans' Stage 2, characterized by the first appearance of striate bisaccate pollen and new forms of Apiculatisporis, Lophotriletes, and certain cingulate mesosporoids (Norvick, 1971, unpubl.).

The boundary is poorly defined palaeontologically in Europe. Jongmans & Gothan (1937) remarked that Schmidt had also noticed Gothan's observation that Callipteris conferta and Medlicottia first appeared at about the same time. Jongmans & Gothan considered that the first appearance of Medlicottia in the Krasnofimsk Formation of Russia marked the base of the Permian.

This suggests that the first appearance of Callipteris conferta may also mark the beginning of the Permian; "however according to Gothan Callipteris is very much influenced by the facies of floral association" (Jongmans, 1952) and it is questionable whether it is restricted to the early Permian.

Helby (1966) suggested that Callipteris conferta may be represented by Illinites unicus in dispersed spore assemblages. "Illinites unicus is a characteristic component of the assemblage replacing the Lycospora Microflora in Western Europe, North America, and the USSR" (Helby, 1969b, p. 11).

Although Illinites has not been found in Australia, Helby considered the microfloral change from the "Lycosporoid" microflora of Balme (1964) to the Potonieisporites Microflora as being a comparable change and that the incoming of Potonieisporites marks the top of the Carboniferous. However, this scheme also suggests that some of the Pennsylvanian of North America, nearly all the Uralian of the USSR, and much of the Stephanian C of Western Europe are also post-Carboniferous.

There are obvious problems associated with using the incoming of Callipteris conferta to mark the top of the Carboniferous; furthermore the translation of this boundary into Australian terms, when none of the above index species are present, is fraught with danger. This is highlighted by the recent suggestion (Helby & McClung, 1973) that the Potonieisporites microflora, rather than signifying post-Carboniferous, is in fact Namurian.

At the present time there is no satisfactory solution to recognizing the Carboniferous-Permian boundary; however, the incoming of the Glossopteris flora approximates the boundary in Australia.

### TECTONICS

The continent, except for the Tasman Geosyncline, was subjected to mainly vertical movements during the Silesian. In the Western Australian intracratonic basins (Bonaparte Gulf, Canning, and Carnarvon Basins), this resulted in uplift, erosion, and rejuvenation of source areas leading to the accumulation of coarse terrigenous sediments, marking the beginning of a depositional episode extending into the Late Permian. Gentle folding and fairly extensive faulting along the flanks of these basins probably persisted into Permian times.

In the Tasman Geosyncline, intensive diastrophic movements occurred during the Silesian, together with the formation of volcanic and plutonic rocks. These movements affected the Neranleigh Fernvale Group (including the Brisbane Metamorphics), and represent the later phases of the Kanimblan Orogeny (Green, 1973). Both in the Yarrol Trough and the northern part of the Tamworth Trough, there is a break between the earliest Namurian and Westphalian. In the lower Hunter Valley, however, marine sedimentation continued, but it was accompanied by widespread volcanicity and the introduction of glaciogene sediments.

The faunas became provincial and most genera were new and endemic. This has been attributed to the considerable change in conditions brought about by the relative movement of the continent into a near polar position.

Towards the end of the Silesian nonmarine sedimentation occurred over large areas of Central and Western Australia; in the east, terrestrial sediments interdigitate with volcanics and glaciogene sediments including varve-like sequences; marine faunas are rare and are restricted to polyzoans, molluscs, and a few endemic brachiopods.

In the Early Permian there was a transgression with widespread marine deposits in both eastern and Western Australia.



## CONCLUSIONS

The geology of the Silesian in Australia is complicated by continent-wide tectonic activity and the onset of glaciation which caused faunal and floral provincialism.

The pattern of sedimentation changed from west to east: in the western and possibly the central parts, large areas were covered with continental deposits; on the eastern margin volcanics interfingered with clastics and glaciogene deposits which, in turn, interfingered with open marine sediments.

Good biostratigraphic information at the Viséan/Namurian boundary and in the Sakmarian adequately delineates the Silesian. However, the presence of endemic species and the virtual absence of northern hemisphere forms makes correlation with the type section difficult.

## REFERENCES

(The numbers which appear before the references refer to those shown at the top of Fig. 2)

1. AMOS, A.J., & SABATTINI, N., 1969 - Upper Palaeozoic faunal similitude between Argentina and Australia. in Gondwana Stratigraphy, IUGS Symp. 1967, UNESCO Earth Sciences 2, 235-248.
2. BALME, B.E., 1960 - Notes on some Carboniferous microfloras from Western Australia. Compte Rendu 4 eme Congress Avanc. Etud. Stratigr. Carbonifere, 1, 25-31.
3. BALME, B.E., 1964 - The palynological record of Australian pre-Tertiary floras. In Ancient Pacific Floras (10th Pacif. Sci. Congr. Honolulu, 1961), 49-80, 7 pls.
4. BALME, B.E., 1973 - Age of a mixed Cardiopteris-Glossopteris Flora. J. geol. Soc. Aust., 20(1), 103.
5. BANKS, M.R., & CLARKE, M.J., 1973 - Tasmania, Parmeener Supergroup in 3rd Int. Gondwana Symp., Guidebook, Field Trip 1, 23-47.
6. BLACK, L.P., MORGAN, W.R., & WHITE, M.E., 1972 - Age of a mixed Cardiopteris-Glossopteris Flora from Rb-Sr measurements on the Nychum Volcanics, North Queensland. J. geol. Soc. Aust., 19(2), 189-196.
7. BÖGER, H., & FIEBIG, H., 1963 - Die fauna des Westdeutschen Oberkarbons II. Die Articulaten Brachiopoden des Westdeutschen Oberkarbons. Palaeontographica Abt A. 122 (4/6), 111-165, pls. 14-23, 25 text-figs.
8. BROWN, D.A., CAMPBELL, K.S.W., & CROOK, K.A.W., 1968 - The Geological evolution of Australia and New Zealand. Pergamon Press, 1-409.
9. CAMPBELL, K.S.W., 1961 - Carboniferous Fossils from the Kuttung Rocks of New South Wales. Palaeontology, 4(3), 428-474, pl. 53-56.
10. CAMPBELL, K.S.W., 1962 - Marine Fossils from the Carboniferous Glacial Rocks of New South Wales. J. Paleont., 36, 38-52.

11. CAMPBELL, K.S.W., & McKELLAR, R.G., 1969 - Eastern Australian Carboniferous invertebrates, sequence and affinities; in K.S.W. CAMPBELL (Ed.), STRATIGRAPHY AND PALAEOONTOLOGY, Canberra, ANU Press, 77-119.
12. CAMPBELL, K.S.W., & McKELVEY, B.C., 1972 - The geology of the Barrington district, N.S.W. Pacific Geology 5, 7-43, 4 pls. 10 figs.
13. CROWELL, J.C., & FRANKS, L.A., 1971 - Late Palaeozoic glaciation of Australia. J. geol. Soc. Aust., 17(2), 115-155.
14. DICKINS, J.M., & MALONE, E.J., 1973 - Geology of the Bowen Basin. Bur. Miner. Resour. Aust. Bull. 130, 1-154.
15. ENGEL, B.A., 1962 - Geology of the Bulahdelah - Port Stephens district, N.S.W. J. Proc. R. Soc. N.S.W. 95, 197-215.
16. ENGEL, B.E., 1965 - Carboniferous Studies in New South Wales, Australia. D.N. Wadia Commem. Vol., Min. Metall. Inst. India, 196-216.
17. ENGEL, B.A., 1966 - Newcastle, N.S.W. - 1:250 000 Geological Series. Dep. Mines Geol. Surv. N.S.W. explan. Notes SI/56-2.
18. EVANS, P.R., 1969 - Upper Carboniferous and Permian palynological stages and their distribution in eastern Australia in Gondwana stratigraphy, IUGS Symp. 1967, UNESCO Earth Sciences 2, 41-54.
19. FLEMING, P.J.G., 1969 - Fossils from the Neerkol Formation of Central Queensland; in K.S.W. CAMPBELL (Ed.), STRATIGRAPHY AND PALAEOONTOLOGY, Canberra, ANU Press, 264-275.
20. GOSTIN, V.A., & HERBERT, C., 1973 - Stratigraphy of the Upper Carboniferous and Lower Permian Sequence, Southern Sydney Basin. J. geol. Soc. Aust., 20, 49-70.
21. GREEN, D.C., 1973 - Radiometric Evidence of the Kanimblan Orogeny in Southeastern Queensland and the Age of the Neranleigh-Fernvale Group. J. geol. Soc. Aust., 20, 153-160.



22. GULLINE, A.B., 1967,- The first proved Carboniferous Deposits in Tasmania. Aust. J. Sci., 29, 332-333.
23. HELBY, R., 1966 - Sporologische Untersuchungen an der Karbon/  
Perm-Grenze im Pfälzer Bergland. Fortschr. Geol. Rheinld.  
Westf., 13, 645-704.
24. HELBY, R., 1969a - The Carboniferous - Permian Boundary in Eastern Australia: an interpretation on the basis of palynological information. Spec. Publs. geol. Soc. Aust., 2, 69-72.
25. HELBY, R., 1969b - Preliminary palynological study of Kuttung sediments in central eastern New South Wales. Rec. geol. Surv. N.S.W., 11(1), 5-14.
26. HELBY, R., & HERBERT, C., 1971 - Relationships of the Clyde Coal Measures to the Pigeon House Creek Siltstone. Quart. Notes geol. Surv. N.S.W., 5, 1-6.
27. HELBY, R., & McCLUNG, G., 1973 - Microfloral and faunal evidence for dating Australian Late Palaeozoic Glacials. 3rd Int. Gondwana Symposium, Abstracts, 39.
28. JONES, P.J., CAMPBELL, K.S.W., & ROBERTS, J., in press - Correlation chart for the Carboniferous System of Australia. Bur. Miner. Resour. Aust. Bull. 156A.
29. JONGMANS, W.J., & GOTHAN, W., 1937 - Betrachtungen über die Ergebnisse des zweiten Kongresses für Karbonstratigraphie. Compte Rendu. 2 eme Congres Avanc. Etud. Stratigr. Carbonifere, 1, 4-5.
30. KIRKEGAARD, A.G., SHAW, R.D., & MURRAY, C.G., 1970 - Geology of the Rockhampton and Port Clinton 1:250 000 Sheet areas. Rep. geol. Surv. Qld., 38, 1-155.
31. KRAUSEL, R., -1961 - Die Permflora der Südhälfte der Erde und ihre Beziehungen zu anderen Floren, in Recent Advances in Botany. 9th Int. bot. Congr. Montreal 1959, 2, 951-953.

32. LINDSAY, J.F., 1969, - Stratigraphy and structure of the Palaeozoic sediments of the Lower Macleay Region, Northeastern New South Wales. J. Proc. R. Soc. N.S.W., 102, 41-55.
- 
33. LUDBROOK, N.H., 1961 - Permian to Cretaceous subsurface stratigraphy between Lake Phillipson and the Peake and Denison Ranges, South Australia. Trans. R. Soc. S.Aust. 85, 67-80.
34. McKELLAR, R.G., 1965 - An Upper Carboniferous Brachiopod Fauna from the Monto District, Queensland. Publ. geol. Surv. Qld., 328 Palaeont. Pap. 1, 1-15.
35. McKELLAR, R.G., 1967 - The geology of the Cannindah Creek area, Monto district, Queensland. Publ. geol. Surv. Qld., 331, 1-38.
36. McWHAE, J.R.H., PLAYFORD, P.E., LINDNER, A.W., GLENISTER, B.F., & BALME, B.E., 1958 - The Stratigraphy of Western Australia. J. geol. Soc. Aust., 4, part 2.
37. MAMET, B.L., & BELFORD, D.J., 1968 - Carboniferous foraminifera, Bonaparte Gulf Basin, northwestern Australia. Micropaleontology, 14(3), 339-347.
38. NORVICK, M., 1971 - Some Palynological observations on Amerada Thunderbolt No. 1 Well, Galilee Basin, Queensland. Bur. Miner. Resour. Aust. Rec. 1971/53 (unpubl.).
39. OLGERS, F., 1973 - The geology of the Drummond Basin, Queensland. Bur. Miner. Resour. Aust. Bull. 132, 1-76.
40. OSBORNE, G.D., 1922 - Geology and Petrography of the Clarencetown-Paterson District. Proc. Linn. Soc. N.S.W., 47, 161 and 519.
41. PALMIERI, V., 1969 - Upper Carboniferous conodonts from limestones near Murgon, southeast Queensland. Publ. geol. Surv. Qld., 341, Palaeont. Pap. 18, 1-13, pls. 1-7.
- 
42. PICKETT, J.W., 1966 - Lower Carboniferous Coral Faunas from the New England District of New South Wales. Geol. Surv. N.S.W., Mem. Palaeont., 15, 38, pl. 20.

43. PLAYFORD, G., & HELEY, R., 1968 - Spores from a Carboniferous Section in the Hunter Valley, New South Wales. J. geol. Soc. Aust., 15(1), 103-119, pls. 9-11.
44. RATTIGAN, J.H., 1967a - The Balickera Section of the Carboniferous Kuttung Facies. J. Proc. R. Soc. N.S.W., 100(2), 75-84.
45. RATTIGAN, J.H., 1967b - Cyclic Sedimentation in the Carboniferous Continental Kuttung Facies, New South Wales. J. Proc. R. Soc. N.S.W., 100(3), 119-128.
46. RIGBY, J.F., 1973 - Gondwanidium and other similar Upper Palaeozoic genera, and their stratigraphic significance. Publ. geol. Surv. Qld., 350, 1-10, pls 1-3.
47. ROBERTS, J., 1971 - Devonian and Carboniferous brachiopods from the Bonaparte Gulf Basin, northwestern Australia. Bur. Miner. Resour. Aust., Bull. 122, 1-319.
48. ROBERTS, J., & OVERSEY, B., 1973 - The early Carboniferous Palaeogeography of the Southern New England Belt, New South Wales. J. geol. Soc. Aust., 20, 161-174.
49. RUNNEGAR, B.N., 1969 - The Permian Faunal Succession in Eastern Australia. Spec. Publs geol. Soc. Aust., 2, 73-98, pl. 1.
50. STRZELECKI, P. DE, 1845 - Physical Description of New South Wales and Van Diemen's Land. Longmans London, 1-462, pls 1-19.
51. THOMAS, G.A., 1971 - Carboniferous and Early Permian Brachiopods from Western Australia. Bur. Miner. Resour. Aust. Bull. 56, 1-215, pls 1-31.
52. VEEVERS, J.J., & ROBERTS, J., 1968 - Upper Palaeozoic rocks, Bonaparte Gulf Basin of northwestern Australia. Bull. Bur. Miner. Resour. Aust., 97, 1-155.
53. WALKOM, A.B., 1920 - Notes on the correlation of the fossil floras of the Carboniferous in SUSSEILCH, C.A. & DAVID, T.W.E., Sequence, glaciation and correlation of the Carboniferous rocks of the Hunter River district, New South Wales. J. Proc. R. Soc. N.S.W., 53, 282-287.

54. WYATT, D.H., 1968 - Townsville, Queensland 1:250 000 Geological Series. Bur. Miner. Resour. Aust. explan. Notes SE 55/14.
55. WYATT, D.H., PAINE, A.G.L., HARDING, R.R., & CLARKE, D.E., 1970 - Geology of the Townsville 1:250 000 sheet area, Queensland.  
Bur. Miner. Resour. Aust. Rep. 127, 1-85.