


1990/7
c.2



~~CONFIDENTIAL~~
SEE INSIDE
COVER.

Bureau of Mineral Resources, Geology & Geophysics

EMR PUBLICATIONS COMPACTUS
(NON-LENDING-SECTION)



RECORD 1990/7

THE NORTHEAST AUSTRALIAN MARGIN AND ADJACENT AREAS - A BIOSTRATIGRAPHIC REVIEW AND GEOHISTORY ANALYSIS

by

George C.H. Chaproniere, Christopher J. Pigram, Philip A. Symonds, Peter J. Davies.

1990/7
c.2

Information in this report has been obtained by the Bureau of Mineral Resources, Geology and Geophysics as part of the policy of the Government to assist in the exploration and development of mineral resources. It may not be published in any form or used in a company prospectus or statement without the permission in writing of the Director.

Department of Primary Industries and Energy
BUREAU OF MINERAL RESOURCES, GEOLOGY AND GEOPHYSICS

RECORD 1990/7

THE NORTHEAST AUSTRALIAN MARGIN AND ADJACENT AREAS -

A BIOSTRATIGRAPHIC REVIEW AND GEOHISTORY ANALYSIS

by

George C.H. Chaproniere, Christopher J. Pigram, Philip A. Symonds, Peter J. Davies.

(Division of Marine Geosciences and Petroleum Geology)



BMR Continental Margins Program

The Northeast Australian Margin and Adjacent Areas - A Biostratigraphic Review and Geohistory Analysis

Copy No:

To:

This publication is for use of the above organisation only and no part of it can be reproduced or transferred to any third party without the express permission of the Director, Bureau of Mineral Resources, Geology and Geophysics, GPO Box 378, Canberra City, ACT 2601, Australia.

CONTENTS

Summary	iii
Introduction.....	1
Biostratigraphic Summary.....	2
Basin Margin sections	3
Eocene	4
Late Oligocene to Early Middle Miocene.....	5
Post-Middle Miocene	6
Basinal sections	7
Capricorn Basin.....	7
Coral Sea Basin	8
Lord Howe Rise	9
Dredge Material	9
Gravity and Piston Cores	11
Regional Summary.....	11
Eocene	11
Oligocene to Middle Miocene	11
Latest Miocene to Holocene.....	11
Unconformities	12
Middle Eocene.....	12
Early Oligocene.....	12
Late Miocene.....	13
Early Pleistocene.....	14
Minor Unconformities	14
Palaeoenvironments	14
Eocene	14
Late Oligocene to Middle Miocene	15
Post-Middle Miocene	16
Predicted Biostratigraphic Sequence for the Northeast Australian Region.....	17
Platforms	17
Southern Part.....	17
Northern Part	17
Troughs.....	18
Geohistory Analysis.....	18
Introduction	18
Results	19
Papuan Basin	19
Capricorn Basin area	21
Coral Sea Basin	21
Lord Howe Rise	21
Discussion	22
Conclusions.....	22
Acknowledgements.....	24
References.....	25

LIST OF FIGURES

Figure 1. Locality map showing data points used in this study.....	2
Figure 2. Biostratigraphic summary for all localities used in this study.....	3
Figure 3. Composite biostratigraphic sequences	4
Figure 4. Biostratigraphic summary for six gravity and piston cores	10

LIST OF TABLES

Table 1. Biostratigraphic range chart for samples from Borabi-1 well.....	7
---	---

LIST OF PLATES

Plate 1. Well analysis diagram for Anchor Cay-1	
Plate 2. Well analysis diagram for Borabi-1	
Plate 3. Well analysis diagram for Kusa-1	
Plate 4. Well analysis diagram for Pasca-C1	
Plate 5. Well analysis diagram for Uramu-1A	
Plate 6. Well analysis diagram for Aquarius-1	
Plate 7. Well analysis diagram for Capricorn-1A	
Plate 8. Well analysis diagram for Sandy Cape-1-3R	
Plate 9. Well analysis diagram for Wreck Island-1	
Plate 10. Well analysis diagram for DSDP Site 209	
Plate 11. Well analysis diagram for DSDP Site 210	
Plate 12. Well analysis diagram for DSDP Site 287	
Plate 13. Well analysis diagram for DSDP Site 208	
Plate 14. Well analysis diagram for DSDP Site 587	
Plate 15. Well analysis diagram for DSDP Site 588	
Plate 16. Geohistory analysis - Anchor Cay-1	
Plate 17. Geohistory analysis - Anchor Cay-1 - with modelled heatflow	
Plate 18. Geohistory analysis - Anchor Cay-1 - with thick pre-Tertiary section	
Plate 19. Geohistory analysis - Borabi-1	
Plate 20. Geohistory analysis - Kusa-1	
Plate 21. Geohistory analysis - Kusa-1 - with modelled heatflow	
Plate 22. Geohistory analysis - Pasca-C1	
Plate 23. Geohistory analysis - Uramu-1A	
Plate 24. Geohistory analysis - Aquarius-1	
Plate 25. Geohistory analysis - Capricorn-1A	
Plate 26. Geohistory analysis - Sandy Cape-1-3R	
Plate 27. Geohistory analysis - Wreck Island-1	
Plate 28. Geohistory analysis - DSDP Site 209	
Plate 29. Geohistory analysis - DSDP Site 210	
Plate 30. Geohistory analysis - DSDP Site 287	
Plate 31. Geohistory analysis - DSDP Site 208	
Plate 32. Geohistory analysis - DSDP Site 587	
Plate 33. Geohistory analysis - DSDP Site 588	

SUMMARY

A review of the Tertiary biostratigraphy of the northeast Australian offshore region is presented in this report, based on seven wells in the Papuan and Capricorn Basins, six DSDP Sites in the Coral Sea Basin and Lord Howe Rise, and a number of dredges and cores taken on various research cruises in the region.

Three Tertiary sedimentary packages, each separated by unconformities are present in the basin margins of the region, but only two are present in the basins.

The first package ranges in age from middle to late Eocene (Zones P.11 to P.17 - Ta_3), and overlies older rocks, both sedimentary and igneous, ranging in age from Palaeozoic to early Eocene. At basin margin locations this package was deposited in non-marine environments at the southern end of the Marion Plateau, and shallow marine situations north of the Plateau.

Deposition of the second sedimentary package was initiated during the late Oligocene, within Zone N.19/20 in the Coral Sea Basin, in Zone P.21 in Anchor Cay-1, and within Zones N.3 or N.4A elsewhere, and then continuing into the middle Miocene, Zone N.9. Water depths on the basin margins were shallow, with larger foraminiferal-red algal carbonates dominating the late Oligocene and early Miocene. Within the late early and early middle Miocene (Zones N.6 to N.9; lower Tf), growth of coral and *Halimeda* commenced in the Papuan Basin, but elsewhere to the south there is no evidence of coral growth, although corals probably were able to survive on the northern parts of the Queensland Plateau. Along the basin margins, sedimentation ceased in the middle Miocene, until the late Miocene, but within the deeper water areas sedimentation appears continuous to the Holocene, except for some localised, short-lived breaks.

Deposition of the third package was initiated during either the latest Miocene (Zone N.17B) or earliest Pliocene (N.18) on the basin margins and plateaux, continuing to the Holocene. Coral and *Halimeda* are present in basal limestones of this package at least in the Papuan Basin, where this initial carbonate phase is overlain by terrigenous and shallow marine siliciclastic sediments. To the south, carbonate sediments were deposited at depths similar to those in the areas at the present time. There is evidence for a widespread hiatus separating the latest Pliocene (Zone N.21) from the middle Pleistocene.

Using the data from the various drill holes and dredge samples taken in the region, it is possible to predict the probable biostratigraphic succession likely to be encountered in any drilling both within the Great Barrier Reef and any sites either in troughs or on plateaux of the southern margins of the Coral Sea Basin.

Geohistory analyses have been made for all wells and DSDP Sites. These results suggest that heatflow regimes throughout the Tertiary cooled from those prevailing during the formation of the basins. Subsidence rates appear to have been gradual through Eocene to Miocene time, rapidly increasing during the Pliocene and Pleistocene for all basins. Sediments in the Papuan Basin became marginally mature by the end of the Mesozoic, but there is strong evidence that maturity levels increased during the Pliocene and Pleistocene. Within the Capricorn Basin, geohistory modeling indicates that levels in the lower part of Aquarius-1 approached marginal maturity only during the Pliocene or Pleistocene.

INTRODUCTION

Eight wells from the southeastern part of the Papuan Basin (Anchor Cay-1, Borabi-1, Kusa-1, Pasca-A1, Pasca-A2, Pasca-C1, Pasca-C2 and Uramu-1A) and four wells from the southern end of the Great Barrier Reef (GBR) and Capricorn Basin (Aquarius-1, Capricorn-1A, GSQ Sandy Cape-1-3R and Wreck Island-1) provide the data base for this study (Figure 1). In addition Deep Sea Drilling Project (DSDP) Sites in the region (208, 209, 210, 287, 587 and 588) have also been used. All but the DSDP sites were drilled on potential reefal structures established either on basement highs or shelf edges, in relatively shallow water. Of the DSDP sites only 588 penetrated to the Mesozoic; 209 and 210 penetrated the Eocene; 587 the latest Miocene, and 208 and 587 were barren over much of the interval drilled. Sites 588 and 287 are very close to 208 and 210 respectively and so should contain a similar section. Thus the sections available for study provide information for the biostratigraphic sequence likely to be encountered in the sites to be drilled during Ocean Drilling Program (ODP) Leg 133. Also a number of dredges and gravity or piston cores were taken during several cruises in the north-east Australian region have also provided data for this study. Well analysis diagrams (Plates 1 to 15) have been prepared for all wells and DSDP sites. The data for these has been obtained from well completion reports, from compilations by Robertson et al., (1984) Research Australia Ltd., and Flower Doery Buchan Ltd., (1984) and from the DSDP Initial Reports. In addition geohistory analysis has been carried out on all drill sites using input data from the same sources as the well analysis diagrams. As with the biostratigraphy, emphasis has been placed on the Tertiary section; the Mesozoic sections are only from the wells of the Papuan and Capricorn Basins.

The biostratigraphic part of this review is based almost entirely on the Tertiary. In those few sections which penetrated the Mesozoic this part of the column marine faunas are solely represented by a few dinoflagellate species and biostratigraphic studies have been based entirely on studies of palynomorph assemblages. In addition, there is also uncertainty as to the stratigraphic units intercepted. Because the Mesozoic part of the section is of importance for petroleum exploration and must be taken into account in any geohistory analysis, summaries for this part of the section are to be found in the well analysis diagrams (Plates 1 to 15); these summaries are based on the unpublished work of H. Struckmeyer (pers. com., 1990). Detailed biostratigraphic reviews have been left to others (e.g. Davey, 1987, and Morgan, 1989).

As would be expected, the wells penetrated sequences which have been deposited in shallow water marine sites during the Tertiary. Deposition for some of the DSDP sites was initiated in shallow water, but show gradual deepening with time (particularly those on the Lord Howe Rise, which subsided to its present depths in the Palaeogene). Other sites have remained at similar depths throughout their history. The majority of dredges taken in the region have sampled shallow water sequences. Of the cores, most sampled only the Pleistocene, but at three sites the late Pliocene was encountered.

To date foraminiferal faunas only from Borabi-1 have been examined; additional faunal and floral data has been obtained from various reports and publications (e.g. Robertson Research Australia Ltd., and Flower Doery Buchan Ltd., 1984; Lloyd, 1978, 1988; Palmieri, 1971, 1974, 1975 and 1984; Chaproniere, 1983; Kennett and von der Borch, 1985). In addition the Pleistocene planktic foraminiferal biostratigraphy for the area around the Queensland and Marion Plateaux is summarised here and detailed results will be published elsewhere.

The time scales used in this work are based on the following sources: Tertiary (Berggren et al., 1985a, b), Cretaceous (Burger, in prep a), Jurassic (Burger, in prep b) and Triassic (Haq et al., 1989).




Figure 1. Locality map showing data points used in this study. A: Exploration wells, dredge and core sites; B: DSDP drill sites.

BIOSTRATIGRAPHIC SUMMARY

A number of bore holes have been drilled into sites at both deep and shallow water localities. In addition a number of sites have been dredged, mainly in the shallow water parts of some of the marginal shelfal areas and plateaux bounding the western part of the Coral Sea Basin. The various sections are discussed below and an attempt is made to relate the dredge material to these sequences. The planktic zonal scheme of Blow (1969) as modified by Kennett and Srinivasan (1983) together with the East Indies Stage Classification (Adams, 1970, 1984; Jenkins et al., 1985) provide the basis for the Tertiary

biostratigraphy used in this report (Figure 2); for the Quaternary, the zonation of Bolli & Premoli Silva (1973) as modified by Chaproniere (1985a), is utilised. Figure 2 summarises the stratigraphy for all the sections and dredges used in this study. Figure 3 is an idealised sequence for both the basin margins (A) and the deep basin (B) locations.

Basin Margin Sections

In the sequences studied from the southeastern Papuan Basin, Eocene sediments are found to unconformably overlie Mesozoic or older rocks. The Eocene is overlain disconformably by a late Oligocene or early Miocene to early middle Miocene sequence, which in turn is overlain by Pliocene and younger sediments. In the southern sequences, Eocene marine sediments are lacking. Instead late Oligocene unconformably overlies a non-marine sequence of unknown age, but possibly Eocene, the remaining section is similar to that from the Papuan Basin.




Figure 2. Biostratigraphic summary for all localities used in this study. (A) after Berggren et al., (1985a, b); (B) after Blow (1969), Berggren (1969), Kennett and Srinivasan (1983; (C) after Adams (1984), Adams et al., (1986), Chaproniere (1981, 1983) and Jenkins et al., (1985).

Eocene: The identification of levels within the Eocene is dependent on the quality of fauna present. Thus, in some sections (Anchor Cay-1) both middle (Ta_3) and late (Tb) Eocene can be identified, but in others (Kusa-1, Borabi-1) only undifferentiated middle to late Eocene (Ta_3 to Tb). In those sections with a diverse fauna (such as Anchor Cay-1), planktic foraminiferids *Acarinina* and *Morozovella* indicate middle Eocene, and higher up, *Subbotina linaperta* indicates late Eocene. In other sections (e.g. Borabi-1) planktics are absent and the presence of larger foraminiferids *Asterocyclina* and *Discocyclina* indicate a general middle to late Eocene age (Ta_3 to Tb), but the presence of *Pellatispira* (in Anchor Cay-1) indicates late Eocene (Tb). A number of other species (such as *Heterostegina saipanensis*, *Spiroclypeus vermicularis* and *Nummulites pengaronensis*) typical of the middle to late Eocene shallow water carbonate sequences in the adjacent Indo-Pacific area (such as Fiji, Tonga, Saipan and Enewetak Islands) are absent from the sections studied here. These species appear to be tropical forms and their absence may have resulted from cooler water temperatures in the study area at that time.




Figure 3. Composite biostratigraphic sequences for (A) basin margin locations, and (B) for basinal locations for the Coral Sea area of northeastern Australia.

With the exception of the Pasca wells, all wells studied from the Papuan Basin contain an Eocene section which has been referred to the Mendi Group (by Lloyd, 1978), a unit made up of bioclastic calcarenites or limestones. The absence of typical Tb sediments (that

is those with *Pellatispira*) in most of the wells (except for Anchor Cay-1) may be the result of either their removal by the erosion episode marked by the unconformity separating these sediments from the late Oligocene or Miocene sequence, due to the intense recrystallisation zone which sometimes appears to cover this interval, or to the absence of *Pellatispira* due to environmental reasons.

To conclude, all shallow water sections show an earlier deeper water, planktic-rich phase, initiated within the middle Eocene, Ta_3 letter stage, which grades upwards into a shallowing sequence, in which larger benthics may become dominant, by the late Eocene, Tb stage. The degree of completeness of the section is probably controlled by the amount eroded off during the interval represented by the unconformity prior to the deposition of the Darai Limestone, or by the amount of recrystallisation which has destroyed the original fabric of the sediments.

Late Oligocene to Early Middle Miocene: Definite late Oligocene planktic assemblages are present in only one well, Anchor Cay-1, where Zone P.21 planktic faunas from the Puri Limestone overlie the Eocene Mendi Group. Larger foraminiferids appear with the gradual lithologic change marking the transition from Puri Limestone to Darai Limestone. The appearance of both *Heterostegina borneensis* and *Lepidocyclina (Eulepidina) ehippioides*, without *Miogypsina* indicates the presence of lower Te . Adams (1984) correlates lower Te with Zones P.19/20 to N.4A (see Figure 2) and at least one locality containing a Zone P.19/20 planktic fauna is present in the region, but in a deep basinal setting (Site 210), which contrasts with the shallow water sequences in which the larger foraminiferids occur. However, larger foraminiferids typical of lower Te are first recorded from levels within Zones N.3 to N.4A elsewhere (Chaproniere, 1976), and it seems probable that the appearance of larger foraminiferal faunas in the shallow water parts of the region took place within the zonal interval N.3 to N.4A; for this reason I have correlated the levels with lower Te larger foraminiferids to Zones N.3 and N.4A in Figure 2. The FAD (first appearance datum) of *Miogypsina* within the range of *L. (Eulepidina) ehippioides* marks the base of upper Te (Jenkins et al., 1985), and the LAD (last appearance datum) of *L. (Eulepidina)* defines the base of lower Tf , an event which takes place within Zone N.6 (Chaproniere, 1981; Belford, 1984). This sequence of events is found in all the wells studied. In Anchor Cay-1, the presence of *Globigerinoides sicanus* and *Praeorbulina glomerosa curva* above the FAD of *Miogypsina* indicates a level within Zone N.8 (late early Miocene). The stage of evolutionary development of both *L. (Nephrolepidina)* (parameter F of Chaproniere, 1980) and *Miogypsina* (parameter V of Drooger, 1952) is typical of that for lower Tf . The highest *Miogypsina* within the upper part of the Darai Limestone is *M. (Lepidosemicyclina) thecideaformis*, and as this disappears at a level below the LAD of *M. (Miogypsina) antillea* (Adams, 1984), its extinction level cannot be used as evidence for correlation with a level high in lower Tf , which would be equivalent to a correlation with planktic Zone N.10 or higher (Adams, 1984). Thus the youngest larger foraminiferal assemblages within the top of the Darai Limestone in Anchor Cay-1 are no younger than early middle Miocene, Zone N.9. A similar sequence for the Darai Limestone is present in the other wells studied, but the Puri Limestone is absent. However, Lloyd (1978) has identified a Zone N.12 to N.14 planktic fauna above the highest larger foraminiferids in Uramu-1A, and Palmieri (1971, 1984) records a similar occurrence in Wreck Island-1 and GSQ Sandy Cape-1-3R. Lloyd (1978) bases his conclusions for a Zone N.12 to N.14 determination on the overlap of *Gr. (Gr.) cultrata* and *Gr. (Turborotalia) siakensis*; the remainder of the fauna ranges from Zones N.12 to N.18. Though the only other record of Zone N.12 to N.14 from shallow water sections is that from Wreck Island-1 to the south, Zone N.18 to N.22 assemblages are found in all the shallow water sections. The species

siakensis is a homomorph of *N. humerosa*, a form which ranges from Zone N.18 to N.22 (Kennett and Srinivasan, 1983); the two forms are thus very difficult to differentiate. In addition the geophysical logs indicate a major unconformity marking the top of the early to early middle Miocene limestones, with no evidence for a marl phase overlying these limestones. Thus, on balance of probabilities, a better interpretation is that specimens of *humerosa* and not *siakensis* are present, indicating that earliest Pliocene (Zone N.18) directly overlies the early middle Miocene (Zone N.9) in Uramu-1A. Further south, however, (in Wreck Island-1, and sample BMR701370 from the shelf east of Sandy Cape-1-3R) good faunal evidence indicates that a middle Miocene (Zone N.10 to N.14) planktic phase overlies the early middle Miocene larger foraminiferal limestones, as it does in the basinal sections.

In Anchor Cay-1 the Oligocene-Miocene cycle of sedimentation commenced during the Late Oligocene (Zone P.21B) with a deeper water planktic limestone (Puri Limestone), containing a subtropical/tropical planktic foraminiferal assemblage. This grades upwards into a shallow water sequence of larger foraminiferal-algal limestones (Darai Limestone), later in the Late Oligocene (Zones N.3 to N.4A). In the other wells from the Papuan Basin incorporated in this study, this planktic phase is absent, and sedimentation commenced with the larger foraminiferal-bearing limestone, probably at the same time that this facies developed in Anchor Cay-1, within lower Te. This larger foraminiferal assemblage is widespread and is present in all the wells (including Wreck Island-1). Study of faunas from cores in Borabi-1 (Table 1) suggest that the Darai Limestone was deposited in gradually shallowing conditions. The larger foraminiferal sequence is similar to that recorded from northwestern Australia, exhibiting changes believed to have been in response to a gradually shallowing environment (Chaproniere, 1975). In support of this conclusion is the gradual increase in proportion of miliolids in the Borabi section from the top of upper Te, until they are the predominant faunal element prior to the appearance of corals within lower Tf; coralline algae are ubiquitous throughout the section, but appear to become more numerous at the top of the lower Tf part of the well. Similar faunas are found in other wells. Corals are recorded from both the upper Te (Pasca-C1) and lower Tf (Borabi-1, Kusa-1, Uramu-1A) indicating that conditions favourable to the development of corals appear to have occurred at different times in the southeastern Papuan Basin. It is possible that at other locations more than one phase of coral development occurred. Conditions conducive to coral growth did not seem to have developed further south in Wreck Island-1 where the larger foraminiferal faunas seem to be constant throughout the section. In some wells (Sandy Cape-1-3R and Wreck Island-1) the planktic foraminiferal faunal component seems to increase with a corresponding reduction in the larger foraminiferids during the early middle Miocene (Zones N.10-N.12).

Post-Middle Miocene: In the Papuan Basin the Darai Limestone is disconformably overlain by either the Era beds or the Orubadi beds (Stewart and Durkee, 1985, fig. 5). This sequence commences with shallow water carbonates, containing shallow water benthic foraminiferids, rare planktics and corals, which grade into more carbonaceous sandstones and mudstones. Lloyd (1978) and Stewart and Durkee (1985) consider that the Era beds are Pliocene to Pleistocene in age; the rare presence of *Globorotalia tosaensis* (in Anchor Cay-1) and *Calcarina spengleri* (in Pasca-C1) supports this assessment. The same authors consider the Orubadi beds, which underlie the Era beds in the Pasca wells and Uramu-1A, to be latest Miocene and Pliocene lateral equivalents of the Era beds; the presence of *Pulleniatina primalis*, *Globigerinoides obliquus extremus* and *Sphaeroidinellopsis paenedehiscens* with *Sphaeroidinella dehiscentis* (in the Pasca wells) and the overlap of *Dentoglobigerina altispira altispira*, *Globigerinoides obliquus extremus* and *Globoquadrina*

dehiscens (in Uramu-1A) supports this contention. *Alveolinella quoyi*, which ranges from the late Miocene to Recent (Adams, 1984) is present in some wells. However, in Anchor Cay-1 this species occurs above the marker for Zone N.21 (late Pliocene), *Globorotalia tosaensis*; in another section *Calcarina spengleri* (Pliocene to Recent - Adams, 1984) is present. This evidence suggests that presence of *A. quoyi* in the beds overlying the Darai Limestone in the Papuan Basin is indicative of ages of no older than latest Miocene. Lloyd (1978, fig.2) considers that the oldest beds in the wells studied from the Papuan Basin in this report are early Pliocene.

		Acervulina inhaerens Alveolinella quoyi Amphistegina bikiniensis Amphistegina radiata Austrotullina howchini Barrizina proteiformis Borelis pugmaeus Borodinia septentrionalis Cyclolopeus (Cyclolopeus) carpenteri Gypsina plana Heterostegina borneensis Lepidocyclina (Eulepidina) ephippoides Lepidocyclina (Nephrolepidina) howchini praehouchini Lepidocyclina (Nephrolepidina) sumatrensis Martinopecten vertebraalis Mioqupsina (Lepidosemicyclina) thecideseiformis Mioqupsina (Mioqupsina) sp. Operculina complanata Operculinella venosa Planorbulinella larvata Pseudorotalia schroeteriana Sorites marginalis Spiroclipeus margaritatus Indeterminable miliolines Indeterminable rotalines Indeterminable textularines No foraminiferids Bryozoa Coralline algae Corals Echinoids Gastropods Helmeda Molluscs Pelecypods																									
4304.5ft	26	?																								4304.5ft	26
4624.5ft	27																									4624.5ft	27
4628.5ft	28	X	X																							4628.5ft	28
4951.5ft	31	?																								4951.5ft	31
4956.5ft	32																									4956.5ft	32
5294.5ft	35																									5294.5ft	35
5945.0ft	38																									5945.0ft	38
6283.0ft	39																									6283.0ft	39
6291.0ft	41																									6291.0ft	41
6526.0ft	42																									6526.0ft	42
6532.0ft	43																									6532.0ft	43
7005.5ft	44																									7005.5ft	44
7323.0ft	45																									7323.0ft	45
7633.3ft	46																									7633.3ft	46
7638.3ft	47																									7638.3ft	47
7943.3ft	48																									7943.3ft	48
7944.8ft	49																									7944.8ft	49
7948.5ft	50																									7948.5ft	50
8286.0ft	51																									8286.0ft	51
8288.0ft	52																									8288.0ft	52
8294.0ft	53																									8294.0ft	53

Table 1. Biostratigraphic range chart for samples studied from Borabi-1 well.

Basinal Sections

Capricorn Basin area: Two wells (Aquarius-1 and Capricorn-1A) were drilled in the Capricorn Basin (in the southern part of the GBR region), in water depths intermediate between the shallow water sequences and those drilled by the DSDP, effectively linking

events between the two depth zones. Palmieri (1971, 1974, 1975) made detailed faunal and biostratigraphic studies of these wells. In both wells the lowest marine beds overlie a non-marine pre-late Oligocene sequence believed to be Eocene in age (Hekel, 1972; Foster, 1982). These marine beds are late Oligocene (Zones N.3 to N.4A) in age, and initially contain glauconite and larger foraminiferids (*Lepidocyclina*, *Miogysinoides* and *Operculina*), which give way to sediments with a foraminiferal fauna very similar to that from the Puri Limestone to the north. This planktic sequence appears to be fairly complete through to the Pliocene at least, on the evidence given by Palmieri (1971), although a hiatus is placed around Zone N.7/N.8 boundary in Capricorn-1A. Within the early to early middle Miocene section in both wells calcarenites containing larger foraminiferids (*Cycloclypeus*, *Lepidocyclina* and *Operculina*) occur interbedded with the planktic foraminiferal chalks, calcisiltites and marls. Palmieri (1974) believed that larger foraminiferids and iron oxides contained in calcisiltites from the lower part of the section in Aquarius-1 were derived from shelfal depths by slumping, and that calcarenites with a similar fauna were the result of localised shallowing. However, it is possible that all larger foraminiferids in both wells at levels other than at the base of the marine sequence were deposited by slumping from shallow water. Planktic faunas seem to be diverse at all levels above the basal beds within these two wells, and thus the larger foraminiferids seem to be out of place. Fourcade and Butterlin (1988) have linked the occurrence of larger foraminiferids in sediments containing good planktic faunas with slumping and redeposition due to changes in sea level in the Bahamas and a similar explanation may be valid for these occurrences. Palmieri (1974, fig. 2) shows that the highest larger foraminiferids occur within the early middle Miocene in both Aquarius-1 and Capricorn-1A, at the same level as in Wreck Island-1.

Coral Sea Basin: Three sites (Sites 209, 210, 287) from two legs (Legs 21 and 30, respectively) have been drilled in the Coral Sea Basin. Site 209 was drilled on the boundary between the northeastern margin of the Queensland Plateau and the southwestern side of the Coral Sea Basin. Sites 210 and 287 were drilled in close proximity in the central part of the basin.

Burns et al., (1973a) have described the sequence encountered at Site 209. The hole bottomed in a middle Eocene sequence of glauconitic sandy mudstones and silty sandstones, which were probably deposited at fairly shallow depths adjacent to a continental margin. Water depths gradually increased in the late Eocene accompanied by a reduction in terrigenous material, although some sediments contain fossil fragments derived from shallow water. The Eocene section is disconformably overlain by a late Oligocene to middle Miocene sequence, which, in common with the Eocene, was deposited in gradually increasing water depths, but poor core recovery prevented detailed biostratigraphic studies being made (Kennett, 1973). The middle Miocene is disconformably overlain by a probable continuous middle Pliocene to Holocene sequence containing a well preserved, exclusively tropical planktic foraminiferal fauna (Kennett, 1973).

Site 210 (described by Burns et al., 1973b), bottomed in an early Eocene sequence, which was deposited at greater depths than those of Site 209, but above the calcium carbonate compensation depth (CCD). The Eocene is disconformably overlain by an apparently complete late Oligocene to Holocene section. However, most of this part was deposited at depths hovering around the CCD producing parts either barren of planktic foraminiferids, or with dissolution effected faunas. From the late Miocene the sequence is made up of turbidites, sometimes containing reworked shallow water assemblages (Burns et al., 1973b). As noted above Fourcade and Butterlin (1988) have linked similar

occurrences with slumping associated with lowering of sea level.

Site 287, which was a repeated attempt to intersect oceanic basement adjacent to Site 210, has been described by Andrews et al., (1975). The hole bottomed in basalt which is overlain by an early to middle Eocene nannofossil chalk with interbedded cherts. The Eocene is disconformably overlain by a late Oligocene nannofossil ooze, which in turn is disconformably overlain by a nonfossiliferous interval believed to be Miocene. A seemingly continuous Pliocene to Holocene sequence which disconformably overlies the ?Miocene sediments, completes the section. As for Site 210, detailed microfossil studies were not made due to the poor preservation and low diversity of the faunas, believed to be the result of deposition near the CCD.

Lord Howe Rise: Only the three most northerly sites drilled on the Lord Howe Rise are discussed here: Sites 208 (Leg 21), 587 and 588 (Leg 90). Site 587 was drilled south of Lansdowne Bank, which forms the southern margin of the Coral Sea Basin. Sites 208 and 588 were drilled in close proximity on the western side of the Lord Howe Rise, south of Site 587.


Sites 208 and 588 penetrated virtually an identical section. Both intersected a siliceous-calcareous middle Eocene unit, but at Site 208 this unit was drilled through and the hole bottomed in a middle Paleocene to Cretaceous sequence. Thus the middle Eocene disconformably overlies the middle Paleocene, and is disconformably overlain by a late Oligocene to Holocene calcareous sequence. Kennett and von der Borch (1985) stated that the late Oligocene to Holocene section at Site 588 is complete, but at Site 208 a minor disconformity separates the early and middle Miocene (Kennett, 1973). The three sites have apparently remained at bathyal depths since at least the Eocene, although Boersma (1985) noted that at Site 587 shallow water benthic foraminiferids were reworked into deeper water deposits during the late Miocene (Zone N.17A).

Dredge Material

Palmieri (1974, 1984) described a fauna dredged from the Australian continental shelf near Fraser Island (BMR 701370 - Marshall, 1971, 1972) which contained a larger foraminiferal fauna including *Lepidocyclina* (*Nephrolepidina*) sp. with a Zone N.14 planktic fauna (*Globigerina nepenthes* and *Paragloborotalia mayeri*); similar faunas to this occur in nearby GSQ Sandy Cape-1-3R as well as in the deeper water sections such as Capricorn-1A and Aquarius-1. This may be evidence for an increase in water depths during the early middle Miocene (within the Zone N.8 to N.9 interval), with the larger foraminiferal faunas being replaced by planktic assemblages, but slumping of shallow water species into deeper water cannot be dismissed.

Chaproniere (1983) described faunas from four dredge hauls at the northern end of the Queensland Trough in the vicinity of Osprey Reef, at the northwestern corner of the Queensland Plateau, collected during R/V "Sonne" Cruise 16. The oldest sample contained a middle Eocene fauna made up of planktics *Acarinina bullbrooki* and *Globigerapsis ?subconglobata* and larger foraminiferids *Asterocyclina incisuricamerata* and *Operculina pacifica*. In addition samples of late Oligocene (lower Te) and early Miocene (upper Te) were collected. A sample of chalk (So16-03KD-D) collected during the same cruise, in the vicinity of the Osprey Embayment, NW of Osprey Reef, contained a well preserved, high diversity planktic fauna, including *Globorotalia* (*Fohsella*) *kugleri kugleri*, *Globoquadrina*

dehiscens, *Globigerinoides quadrilobatus altiapturus*, *Gds. q. subquadratus* and *Globigerina woodi woodi*, indicating Zone N.4B, basal Miocene.



1 Berggren et al 1985 2 Lamb & Beard 1975 3 Saito 1976 4 Kennett & Srinivasan, 1983 5 Thompson et al 1979 6 Bolli & Premoli Silva, 1973

Figure 4. Biostratigraphic summary for six gravity and piston cores taken in the Queensland and Townsville Troughs. Time scale after Bolli and Premoli Silva (1973) and Chaproniere (1985a); Oxygen isotope scale after Shackleton and Opdyke (1976).

More recently, a number of dredge samples were collected during four cruises (50, 51, 75, 76) of the R/V "Rig Seismic" over the Queensland and Marion Plateaux (Figure 1). Some of these (75DR03, 76DR04 to 76DR09) contained larger foraminiferal faunas (*Lepidocyclina* (*Nephrolepidina*) *howchini*) with values for Parameter F (Chaproniere, 1980, 1983, 1984) typical of Zones N.7 and N.8 (late early Miocene). A number of the late early Miocene samples from the Marion Plateau (e.g. 75DR03-III, 76DR02-15) had borings infilled with late Miocene to early Pliocene and Pleistocene planktic foraminiferal ooze, suggesting that they were collected from immediately below the regional late Miocene or early Pliocene unconformity. Also, some (e.g. 75DR02-III) contained lower Tf larger foraminiferids reworked into a Pliocene (Zones N.18 to 19/20) planktic assemblage. These samples were probably collected above the regional unconformity and indicate that the late early Miocene carbonates were being eroded at that time.

Gravity and Piston Cores

A large number of gravity and piston cores have been collected during four cruises of the "Rig Seismic" and one of the "Sonne" off the northeast Australia region. All but two of these penetrated Holocene or Pleistocene (Zones N.22 to N.23) sediments; two (51GC73 and 51GC74) recovered only the latest Pliocene (Zone N.21) and one (So16-13SL) passed through the Pleistocene into the latest Pliocene. Detailed studies of the planktic foraminiferids have been made for six of these cores (see Figure 4) and 12 biostratigraphic events have been recognised for the middle and late Pleistocene (from within the top of the Matuyama Chron) (Chaproniere, in prep.). All six cores penetrated into the middle Pleistocene *Globorotalia crassaformis* Subzone of Zone N.22 (*sensu* Chaproniere, 1985a), but in one core (So16-13SL) this subzone directly overlies the late Pliocene Zone N.21. In addition the two cores which sampled only the latest Pliocene were sited in the floor of a canyon. The top of this Pliocene unit forms a reflector which can be traced on seismic data into the walls of the canyon. The evidence from the three cores suggests that there is a regional break, between the latest Pliocene and the middle Pleistocene, at least within the areas adjacent to the northeast Australian margin. Until detailed studies are made elsewhere, such as in the DSDP Leg 90 sites, the full extent of this break will not be known.

Regional Summary

Three main sedimentary packages can be recognised throughout the region, with one being confined to the shallow water areas. Each being bracketed by unconformities.

Eocene: An early or middle to late Eocene package overlies basaltic basement in the Coral Sea Basin, but in the Papuan Basin it overlies Jurassic or Cretaceous sediments, and on the Lord Howe Rise, Paleocene siliceous chalks. The topmost beds vary in age from middle to late Eocene, due almost certainly to erosion prior to the deposition of the overlying package in the late Oligocene. The Eocene sequence in all areas except the Capricorn Basin, is fully marine, with the basal sediments containing mainly planktic faunas. Gradually a shallow water component (including larger foraminiferids) displaces the planktic assemblage, indicating shallowing. In the deeper parts there is evidence for an increase in water depths (at Site 209), from the middle to late Eocene. In the Capricorn Basin the non-marine section underlying the marine late Oligocene is believed to be of Eocene age, being a possible correlative of the Eocene oil shales on shore.

Oligocene to Middle Miocene: A late Oligocene to late early or early middle Miocene package directly overlies the Eocene surface. The basal beds of this package vary in age from late Oligocene in the deep sea areas to basal Miocene in some of the shallow water sections. Sedimentation of this package ceased, probably within the basal middle Miocene in the shallow water sections, but in the deep sea continued without break to the present day.

Latest Miocene to Holocene: In shallow water areas the deposition of a third package was initiated either in the latest Miocene or the early Pliocene, and continues to the present day. The basal phase of this package is a carbonate platform sequence passing up into terrigenous muds and sands.

UNCONFORMITIES

There are three major unconformities recognised throughout the region, and at least one minor one of limited extent. The three major ones are: the base of the middle to late Eocene sediment package (middle Eocene); the base of the late Oligocene to early middle Miocene package (early Oligocene); and the base of the Pliocene to Holocene package (late Miocene).

Middle Eocene

This unconformity is found in all locations used in this study with the exception of Aquarius-1 where that part of the section below the late Oligocene marine phase is terrigenous (Figure 2). In the Coral Sea Basin, the sediments overlying this unconformity range from fairly shallow water deposits (at Site 209) to probably bathyal at Sites 210 and 287. At these locations the sediments overlie volcanic basement and so are associated with the formation of the Coral Sea Basin crust. The sediments overlying the unconformity at the Lord Howe Rise sites also are of fairly deep water origin and overlie Paleocene and Cretaceous sediments. This unconformity almost certainly is present in the Capricorn Basin, where it probably separates a terrestrial sequence from Palaeozoic rocks. In the Papuan Basin the unconformity universally separates shallow water sediments from Mesozoic or older rocks. Thus in the southeastern Papuan and Capricorn Basins the unconformity separates sedimentary units from rocks of considerably older ages, and is almost certainly related to the formation of the Coral Sea Basin. The formation of this basin permitted shallow marine or non-marine sedimentation to resume over an old topographic surface in the areas now marginal to the basin; within the basin moderate water depths existed as the basin developed. The youngest sediments underlying the unconformity are middle Paleocene, with late Paleocene and part of the early Eocene sediments absent from the region.

Early Oligocene

The oldest sediments overlying the Eocene sediments are of middle Oligocene age (Zone P.19/20), but these have limited distribution being restricted to Site 210 in the Coral Sea Basin only. The Puri Limestone in Anchor Cay-1 contains a Zone P.21B fauna, but it was in the later Oligocene (Zones N.3 to N.4A) that sedimentation became widespread throughout the region. To the east, at the edge of the Australian Plate in the vicinity of Tonga, basal Oligocene (within Zone P.18) sediments are found underlying late Oligocene (Zone P.21B) (Chaproniere, in press). Thus the smallest time gap represented by the unconformity ranges from earliest Oligocene to early middle Oligocene (approximately 3 ma), although in most places the length of the hiatus was longer, from the within Zone P.18 to within Zone P.21 (earliest Oligocene to late Oligocene) representing an interval of approximately 5 ma. This suggests that in some of the shallow water sections in the eastern Papuan Basin erosion may have removed much or all of the late Eocene.

This unconformity is widespread throughout the region ranging from northwestern Australia (Chaproniere, 1984) east to the Tonga Ridge at the edge of the Australian Plate (Chaproniere, in press); it is also present in deep sea sections in the Tasman Basin and Lord Howe Rise (Kennett and von der Borch, 1985), but appears to be absent in the southern parts of the Australian Plate, where sedimentation appears to have been continuous. Kennett et al., (1972) linked the unconformity to the opening of the seaway

between Australia and Antarctica and the resulting circum-Antarctic current eroding any sediments. However, the apparent limitation of the unconformity to the northern part of the Australian Plate where it is universally present, from west to east, implies a vast current if the hypothesis of Kennett et al., (1972) is accepted. Furthermore, currents not only remove sediment they also deposit it, but early Oligocene sediments are not known in the region, with the exception of some isolated occurrences of Tc shallow-water carbonates in Papua New Guinea (Belford, 1984). There seems little doubt that the mechanism causing the removal of sediments over such a vast area is linked to the northern drift of Australia, but other scenarios must also have been involved. The northward drift of Australia must have had considerable effect on equatorial currents by forcing flows southward along barriers such as the Tonga Ridge, Lord Howe Rise and the eastern edge of the Australia-New Guinea landmass. Such currents may well have been able to erode sediment from shelves, but as noted above, sediment removed from one area should be deposited elsewhere. But where is this sediment? Corrosive bottom waters due to the initiation of Antarctic bottom water, related to the formation of the circum-Antarctic circulation and to the onset of major Antarctic glaciation (Keller et al., 1987) may have removed any sediments eroded from the shelves into deeper waters, but it seems unlikely that all shelfal sediments would have been removed. It is improbable that there was a large fall in sea level, as near continuous sedimentation occurred in shallow water areas both in southeastern Australia and New Zealand over the early Oligocene interval. The elevation of the CCD linked to onset of Antarctic glaciation could also explain removal of calcareous sediments, but those sediments of southeastern Australia and New Zealand referred to above, indicate that it did not rise to shelfal depths. It seems that a combination of events must have interacted during the early Oligocene to cause this major and regional unconformity, but the presence of some early Oligocene within the northern part of the Australian plate is expected, but other than some small isolated occurrences of Tc limestone in Papua New Guinea, it seems totally absent.

Late Miocene

This unconformity is present in the wells from the southeastern Papuan Basin and two bores (Wreck Island-1 and Sandy Cape-1-3R) at the southern end of the GBR (Palmieri, 1971, 1984) as well as being evident on the Marion Plateau (Pigram et al., in prep) and at Sites 209 and 287. Though sedimentation was continuous in most other basinal locations, the presence of reworked shallow water foraminiferids in two of the deeper water Sites 210 and 587 may be evidence for shallowing during the late Miocene.

The oldest sediments overlying the early middle Miocene carbonates are almost certainly Pliocene. It is possible that some of these may be latest Miocene as such sediments have been recorded from the Ok Tedi-Wabag area to the north (Belford, 1984), but this seems unlikely as no late Miocene has been reported from DSDP Sites 209 or 287. To the east, late Miocene (Zone N.17A) sediments have been recorded from the Tongan region (Chaproniere, 1985b; in press). Thus the hiatus spans the interval of Zones N.15 to N.16 representing approximately 3 to 4 ma., but as sediments between Zones N.10 and N.17 appear to be absent from the southeast Papuan Basin, the time interval may have been around 15 ma. For the shelfal areas of northeastern Australia, the time interval may have been around 5 ma for this area, based on the section in Sandy Cape-1-3R bore.

In the Papuan Basin wells the unconformity is overlain by carbonates which may contain corals, *Halimeda*, coralline algae and larger foraminiferids (e.g. in Borabi-1) which may represent short-lived reefal conditions. In some of the wells (e.g. Borabi-1, Kusa-1,

Pasca-A2, Uramu-1A) loss of circulation coincides with the unconformity; this may indicate dissolution channels, possibly the result of development of karst topography on the older Miocene carbonates. There is variation in the age of the carbonates underlying the unconformity ranging from early to early middle Miocene. This variation is almost certainly due to the effects of subaerial erosion, a concept supported by the possible evidence for karst in Borabi-1 and other wells.

Because the unconformity is developed in all shallow water sections studied, it is almost certainly the result of sea level fall, the extent of the fall being considerable (Pigram et al., in prep.).

Early Pleistocene

As noted above much of the early Pleistocene is missing in the only core (So16-13SL) to have penetrated the Pleistocene into the late Pliocene. In this core the FAD of *Pulleniatina obliquiloculata finalis* (dated as 1.3 ma., by Berggren et al., 1985b) directly overlies the LAD of *Globigerinoides quadrilobatus fistulosus* (dated at 1.6 ma., by Berggren et al., 1985b), giving a maximum length of 0.3 ma. for the hiatus. The geographic extent of this hiatus is not known, but its presence in the floor of a canyon adjacent to Geranium Passage, some 500 km south of So16-13SL implies that it may be regional.

Minor Unconformities

Palmieri (1971) records an unconformity separating Zones N.7 and N.8 in Wreck Island-1 and Capricorn-1A, which may be equivalent to a minor hiatus separating the early and middle Miocene in Site 208 (Kennett, 1973). However Condon et al., (1953) and Chaproniere (1984) have recognised an unconformity at a slightly higher level, within Zone N.8, in the Northwest Cape area and elsewhere in the Carnarvon Basin, but it seems absent from sections further north, such as Ashmore Reef-1 (Chaproniere, 1981, 1984). It is possible that these records are of the same unconformity, which though regional, is discontinuous in geographic extent. Also it is very short-lived in time, probably being confined to within Zone N.8. It occurs in both the northwestern and northeastern parts of Australia, in areas of differing tectonic settings. This evidence implies a possible relationship to sea-level fall.

At Site 209 there is a marked hiatus from the top of the middle Miocene to the middle Pliocene (Kennett and von der Borch, 1985), and is followed by a continuous sequence to the Holocene. At Site 287 there is an unconformity between the late Oligocene and the barren ?Miocene, and between the ?Miocene and Pliocene to Holocene sequences. In the absence of faunas this unconformity cannot be accurately dated.

PALAEOENVIRONMENTS

Eocene

The Eocene sediments from wells drilled in the Papuan Basin may have a lower part which contain some planktic foraminiferids the presence of which indicates a connection with an open oceanic environment, together with a low diversity larger foraminiferal

assemblage indicative of shallow water. Higher in the Eocene planktic forms disappear, and a more diverse larger foraminiferal assemblage is present. Coralline algae have been recorded only from the late Eocene (Lloyd, 1978); corals have not been recorded. Anchor Cay-1 differs from the other wells by the more diverse planktic faunas that occur throughout the Eocene section which are indicative of water depths greater than those for the other wells studied, although the presence of larger foraminiferids at the top suggest shallowing. Thus all sections studied indicate a marked shallowing in the Late Eocene, but the absence of corals and the low diversity of larger foraminiferids suggests that temperatures were not tropical.

DSDP sites in the region generally indicate bathyal water depths throughout the Eocene, but at Site 209 there is evidence for deposition on a continental margin during the late middle Eocene, followed by gradual deepening during the late Eocene. The faunas have a close affinity with temperate regions (Burns and Andrews, 1973).

Late Oligocene to Middle Miocene

As noted above the only faunas examined for this report are from Borabi-1, where the larger foraminiferid assemblages are associated with coralline algae, generally nodular forms, suggesting shallow depths, high in the euphotic zone. That depths continued to change throughout the deposition of the Oligocene-Miocene carbonates is shown by the gradual change in faunal composition, from one dominated by rotalines to one dominated by miliolines; in most of the wells from the Papuan Basin, the top of the sequence is marked by the presence of corals, indicating minimum depths of 25m or less. Thus water depths gradually shallowed with time. It is possible that increasing water temperature may have influenced the time of coral development, but as coral development occurred earlier in some wells (Pasca - where corals are present within upper Te) than that found in Borabi-1 (lower Tf), temperature control seems unlikely; instead factors such as water depth and substrate may have controlled coral distribution. Anchor Cay-1 differs from the other wells in that planktic foraminiferids occur throughout the sequence, with larger foraminiferids appearing at the top of the section; these assemblages indicate greater water depths than those seen elsewhere, probably below the euphotic zone at the base of the sequence. Planktics gradually give way to larger foraminiferids and coralline algae, indicating gradual shallowing to depths within the euphotic zone, but not shallow enough to permit development of corals. Thus in the southeastern Papuan Basin, water depths appear to have gradually shallowed throughout the Oligocene-Miocene period, finally reaching depths at which coral development could take place at some locations.

Similar faunal assemblages to those found in the Papuan Basin wells are found in Wreck Island-1 and Sandy Cape-1-3R. The larger foraminiferal associations in the lower part of the sequence contain *Lepidocyclina* (*Eulepidina*) *ephippioides*, *L. (Nephrolepidina) sumatrensis/howchini* and *Miogypsina*; even though *Cycloclypeus* occurs in both sequences, other nummulitids *Heterostegina borneensis* and *Spiroclypeus margaritatus* have not been recorded. Higher in the section in Sandy Cape-1-3R, miliolids *Austrotrillina howchini*, *Flosculinella bontangensis* and *Marginopora vertebralis* appear (Palmieri, 1984). Planktic foraminiferids are present throughout the Wreck Island-1 section, but are present only in the upper parts in Sandy Cape-1-3R; coralline algae and corals have not been recorded from either bore, but almost certainly, coralline algae would be present. Thus, depths seem to have been within the euphotic zone throughout the late Oligocene to early middle Miocene, within an oceanic setting most of the time, but the presence of miliolids in Sandy Cape-1-3R suggests the development of localised areas of increased salinity. The presence

of glauconite in the bottom parts of the sections implies warm temperate to subtropical temperatures, but the presence of larger foraminiferids suggests at least subtropical conditions.

The basal marine beds in both Capricorn-1A and Aquarius-1 are glauconitic muds containing planktic foraminiferids as well as a *Lepidocyclina* (*Nephrolepidina*) - *Miogypsina* (*Miogypsinoidea*) assemblage (Palmieri, 1974), similar to that found at the base of the Wreck Island-1 sequence. This fauna indicates subtropical, shallow water, oceanic conditions. The presence of diverse planktic faunas without larger foraminiferids in the overlying section indicates greater water depths, probably below the euphotic zone. A more accurate palaeodepth estimation will have to await the description of the smaller benthic foraminiferids. Within the early and early middle Miocene, Palmieri (1974) reports levels containing larger foraminiferids interspersed with planktic intervals; he explains some of these occurrences of larger foraminiferids as resulting from slumping from adjacent shelfal areas, but others he explains by localised shallowing events. Without examination of the smaller benthic foraminiferids it is difficult arrive at any firm conclusions, but it seems that slumping from shallow waters, could explain all occurrences without recourse to proposing local shallowing events.

Within the Coral Sea Basin water depths throughout the Oligocene and Miocene remained close to the CCD (Kennett and von der Borch, 1985).

Post-Middle Miocene

In the southeastern part of the Papuan Basin the early or middle Miocene carbonates are locally overlain by a thin limestone which in Borabi-1 contains corals, coralline algae, *Halimeda* and a typically Pliocene or younger larger foraminiferal assemblage *Alveolinella quoyi*, *Marginopora vertebralis*, *Acervulina inhaerens* and *Gypsina plana*. This limestone is succeeded by a sequence of carbonaceous sands and marls containing shallow water benthic foraminiferids and some planktic forms. Elsewhere, the Miocene carbonates are succeeded by carbonaceous sands and marls containing diverse planktic faunas. Thus the beds overlying the early to middle Miocene carbonates show evidence of an initial reefal development within the Pliocene, which were soon swamped by a terrigenous-derived sequence in which depths varied; generally there was a connection to open oceanic conditions.

In the southern GBR area shallow water Pliocene calcarenites were succeeded by Pleistocene reefal sediments in Wreck Island-1 (Palmieri, 1974). In Sandy Cape-1-3R the middle Miocene is overlain by mixed carbonate and terrigenous sediments often containing coralline algae, corals, planktic and larger benthic foraminiferids. Within the Capricorn Basin planktic foraminiferids are very common in the Pliocene sediments, but lack of samples prevented study of the higher levels in Aquarius-1 and Capricorn-1A wells (Palmieri, 1974). However, it is likely that sedimentation continued through the Pleistocene at depths similar to those existing at these sites at the present day.

Within the Coral Sea Basin sedimentation continued throughout the Pliocene to the Holocene under similar conditions to the present day.

PREDICTED BIOSTRATIGRAPHIC SEQUENCE FOR THE NORTHEAST AUSTRALIAN REGION

From the above discussion and using the hypothesis for the control of reef development by the northward drift of Australia proposed by Davies et al. (1987), it is possible to predict the biostratigraphic sequence expected for the marginal platforms and troughs in the offshore area of northeastern Australia.

Platforms

Although DSDP Site 209 was drilled on the northeastern margin of the Queensland Plateau, no other drilling has taken place on either the Queensland or Marion Plateaux. Evidence for the predicted stratigraphic section comprising both plateaux comes from dredges and cores, which were taken mainly at the margins, as well as correlations with the few data points in the area, such as Aquarius-1, Capricorn-1A, Sandy Cape-1-3R and Wreck Island-1 wells for the southern end, and Anchor Cay-1 for the northern parts. The basement for the Queensland Plateau is believed to be a subsided extension of the Tasman Fold Belt (Ewing et al., 1970). Nothing is known for the Marion Plateau, but evidence from wells at the southern end of the plateau suggest that Mesozoic igneous or volcanic rocks and slightly metamorphosed sandstones and shale (Carlsen and Wilson, 1968a, b) may form the upper part of the basement sequence.

Southern Part: The biostratigraphic section on highs built-up on Mesozoic sandstones and shales or igneous rocks forming the basement of the Marion Plateau would be expected to be similar to that encountered in Wreck Island-1; that is the basal marine section would be a late Oligocene section directly overlying non-marine sediments of possible Eocene age. These older marine sediments would be glauconitic and contain larger benthic as well as planktic forms. Sedimentation probably continued into the early middle Miocene, with sediments composed mainly of larger foraminiferal limestones. Given the probable latitudinal position of this part (Marion Plateau) at this time (Davies et al., 1987; Feary et al., in prep.), it is unlikely that water temperatures would have been sufficiently warm to permit coral growth. A marked unconformity spanning the late Miocene is expected, with sedimentation recommencing in the early Pliocene with shallow water bioclastic deposits. At this time warmer temperatures, due to Australia's more northern position than during the earlier Miocene, may have permitted the initiation of reef growth, which has continued to the present day. Areas adjacent to the plateau may have been deeper throughout much of the Tertiary resulting in a sequence similar to that of the Capricorn Basin, that is with continuous planktic faunas with shallow water debris derived from adjacent shallow water areas.

Northern Part: The Queensland Plateau would be expected to exhibit an identical biostratigraphy to the Marion Plateau, but may differ in one respect. Shallow water Eocene marine sediments have been recorded in the vicinity of Osprey Reef (Chaproniere, 1983), indicating the possibility of such an occurrence elsewhere on the plateau, instead of the non-marine sequence found further south. Following the hypothesis advanced by Davies et al., (1987) it is possible that the slightly warmer water temperatures prevailing in this part of the area during the late Oligocene and early Miocene may be reflected in a more diverse larger foraminiferal assemblage, especially higher in the early Miocene, than that recorded in Wreck Island-1.

Troughs

As noted above the oldest marine sediments recorded from the Coral Sea Basin (at Site 287) are early or middle Eocene, directly overlying oceanic crust. In areas of continental crust (such as the Queensland and Townsville Troughs - Symonds and Davies, 1988), however, Mesozoic (possibly Cretaceous) sandstones and shales have been recorded from the wells in the Capricorn Basin (Carlsen and Wilson, 1968a, b), where they are overlain by a possible Paleocene to Eocene non-marine sequence (Hekel, 1972; Foster, 1982). The Eocene is disconformably overlain by a sequence of late Oligocene to Holocene pelagic sediments, deposited close to the CCD. The water depths in the Queensland and Townsville Troughs are much shallower than those of the Coral Sea Basin, and dissolution effects, if encountered, should be minimal. Thus the oldest marine sediments within the troughs could be expected to be early or middle Eocene carbonates of possible shallow water origin. The late Eocene should be disconformably overlain by late Oligocene sediments made up of pelagic sediments mixed with shallow water debris derived from adjacent shallow water areas. It is difficult to assess whether a continuous sequence from late Oligocene to Holocene is present, or whether disconformities in the late Miocene should be present. Evidence from the Capricorn Basin wells, drilled in shallower water depths than those within the troughs, suggests that a continuous section could be expected. Levels of concentration of shallow water derived fossils may be evidence for sea level fluctuations.

GEOHISTORY ANALYSIS

Introduction

Geohistory analyses for all wells and DSDP sites used in this study (Plates 16-30) were undertaken using the method described by Falvey and Deighton (1982) and Williamson and Pigram (1986). This method models subsidence and thermal maturation from data extracted from these well sections. Input data has largely been obtained from well completion reports, although, for the Papuan Basin wells, data has also been extracted from other sources, such as Burns and Bein (1980) and Stewart and Durkee (1985), and for the Capricorn Basin from Ericson (1976); reliability of the analysis is dependent upon the quality of this data. With the exception of Kusa-1 (drilled in 1977) and Sandy Cape-1-3R (drilled in 1980) all wells are old being drilled prior to 1970. Because of the age of these wells, temperature data is lacking for almost all locations except for some from the Papuan Basin, and even here temperature logs are limited to the upper part of the well. Also, of these locations the temperature logs appear to have been run to locate the position of cement during casing operations and appear to be distorted due to the exothermic reaction of setting cement and thus have no value for heatflow estimations. Vitrinite reflectance data is available for only two sections (Anchor Cay-1 and Kusa-1), and porosity data is lacking for all sections. For this discussion, vitrinite reflectance values in the range 0.6-0.8 are considered marginally mature, 0.8-1.3 represents the peak generation window, and values greater than 1.3 correspond to the dry gas-condensate window. As thermal conductivity is a major parameter, the lack of these data has required that the analyses depend on modelling to obtain these parameters. However, the results do have some implications for the oil prospectivity of the region, in particular the southern Papuan Basin, and provide a good basis for an analysis of the relative prospectivity of basins within the region.

At unconformities it is necessary to attempt to evaluate the amount of sedimentary section removed by erosion and the time interval involved, and based on these figures, to make an estimate of the amount of sediment removed. For the first of these figures it is appropriate to estimate the time at which sedimentation ended and erosion commenced; the second of these is constrained by the quality of the faunal or floral evidence. The ages used have been taken from Figure 2.

Heatflow modelling for the well locations in proximity to the rifted margins was undertaken using the method of Sleep (1971). This modelling shows that the heatflow rose to the time of breakup and then slowly decayed to the present day. For well and DSDP sites that are axial to seafloor spreading ridges (that is basins within transform margins such as the Capricorn Basin) numerical modelling was undertaken to ascertain the heatflow history at these locations. Present day heatflow values for the region are based on Cull and Conley (1983, fig. 4).

Heatflow within the region has been controlled by two rifting episodes: the first in the early Mesozoic, which defined the east-west trend of the northern margin of Australia (Pigram and Panggabean, 1984); the second in the Middle or Late Cretaceous, which preceded to the opening of the Coral Sea Basin; peak palaeoheatflow related to the latter regime occurred at the time of Coral Sea Basin breakup about 63Ma (magnetic anomaly 27) (Symonds et al., 1984; Weissel and Watts, 1979). Subsequently the general heatflow within the region has declined. The Cretaceous/Paleocene heatflow peak occurred during a period of time represented by a major unconformity, and so would not be shown on the palaeoheatflow plots; for this reason an artificial point has been inserted at 63Ma so that this heatflow event is incorporated into the analysis. Present day heatflow in the region is about 60mW/m² (Cull and Conley, 1983), and this figure (as 1.43 h.f.u.) has been used in the geohistory analyses for all the sites in the Capricorn and Papuan Basins.

Results

Papuan Basin: Three major unconformities are present in the sections studied from the Papuan Basin and each has been modelled with different parameters. The age for the top of the Mesozoic section is poorly constrained because of poor fossil control or uncertainty as to the stratigraphic units represented, which is in contrast to the more closely dated start of Eocene sedimentation. Because of these age constraints this unconformity separating the Cainozoic from the Mesozoic has been modelled using an arbitrary figure of 1000m for the amount of section removed by erosion during the interval prior to the resumption of Tertiary sedimentation. This figure is supported by the findings of Hill and Gleadow (1989) who estimated that ≈800 m of Mesozoic section was removed during the Paleocene erosion event in the Iehi-1 well in the Papuan Fold Belt to the north of the wells studied here. For the break between the Eocene and late Oligocene it has been assumed that sedimentation ceased at the end of Tc letter stage, coinciding with a major fall in sea-level at around 34Ma; shallow water sediments of Td age are virtually absent from the region suggesting that the Td stage coincides with this period of lowered sea-level. Because shallow water carbonates tend to become cemented shortly after deposition, the compaction of the Eocene limestones due to the overlying sediment load is assumed to be zero. Also, because erosion on limestones takes place by solution, it is expected that only a small amount of section has been removed and a figure of 50m is considered a reasonable estimate. The cessation of early middle Miocene sedimentation is estimated to have occurred at 13Ma, coinciding with a sea-level fall at the top of Zone N.11, with sedimentation being resumed at 5Ma. These Miocene sediments are limestones, so similar arguments for assessing the

compaction effects and the amount removed for the Eocene carbonates are used here; because the time interval ranges from 12 to 15Ma (compared with approximately 6Ma between the Eocene and late Oligocene) a figure of 100m has been used for the amount removed by erosion.

Three wells, Anchor Cay-1, Kusa-1 and Borabi-1 form a north-south transect across the Papuan Basin. These geohistory plots show a marked similarity, with rapid subsidence through the early and middle Jurassic leveling out for the remainder of the Mesozoic. As noted above, much of the Cretaceous section is missing, presumably lost by erosion. The Tertiary again was a period of increased subsidence, particularly during the deposition of the post-Miocene section, which is time transgressive to the south. This subsidence phase is related to flexuring of the margin associated with foreland basin development caused by the collision to the north, which formed the New Guinea orogen, and to the subsequent loading by sediments derived from the rapidly uplifted rocks forming the orogen (Pigram et al., 1990).

The vitrinite data for the two wells (Anchor Cay-1 and Kusa-1), which penetrated a significant part of the pre-Tertiary section, show that the Mesozoic and older rocks within this part of the Papuan Basin became marginally mature by the late Jurassic, associated with the early Jurassic rift-related heatflow peak. The dead oil staining observed within the Jurassic section of Anchor Cay-1 is probably related to this period of maturation. These Mesozoic rocks were effectively frozen at this maturation level until the Late Miocene, when rapid subsidence associated with large volumes of clastic sediment entering the Papuan Basin from the New Guinea orogen, causing rapid burial. In all other wells, the computed vitrinite reflectance plots indicate values of less than 0.5.

All five Papuan Basin wells show similar geohistory plots, especially for the Tertiary, with gradual subsidence through the Eocene to Miocene, with the rate of subsidence increasing rapidly during the Late Miocene or Pliocene. Anchor Cay-1 and Kusa-1 are the only wells with a good Mesozoic section, with the former having the best section which may go to the base of the Triassic. Three wells show modelled vitrinite values of 0.5 or greater; in Anchor Cay-1 the 0.5 level was reached within the Late Jurassic (at approximately 2700m), the 0.6 level within the Middle Cretaceous (at approximately 3000m) and the 0.7 level within the Tertiary (at approximately 3300m); in Kusa-1 the 0.5 level was reached by the Middle Cretaceous (at approximately 2800m) and the 0.6 level by the Paleocene (at approximately 3200m); in Pasca-C1, for which there is no vitrinite reflectance data, the 0.5 to 0.7 levels were reached in the Pliocene (below 3100m).

The basement subsidence curves for all wells show a gradually subsiding basement throughout the Mesozoic, with uplift occurring by the Eocene, preceding the opening of the Coral Sea Basin. The rate of basement subsidence was again gradual during the Eocene to Miocene, increasing rapidly in the Late Miocene and Pliocene. The increase in rate of subsidence occurs slightly earlier in Uramu-1A than in the other wells indicating that the event was time transgressive, being later to the south. The basement subsidence curve for Anchor Cay shows some minor uplift during the Late Jurassic, which is probably related to water depth changes induced by possible sea-level fluctuations.

Though the computed iso-vitrinite reflectance plot appears to match the observed data plot for Kusa-1 (Plate 20), this is not the case for Anchor Cay-1, where there is a marked disparity between the two plots (Plate 16). In an attempt to reconcile these differences a number of models were produced. Firstly the palaeoheatflow curve was modified to reflect an increased heatflow peaking at magnetic anomaly 27 (63 Ma),

plateauing until anomaly 22 (51 Ma) during the development of a possible transform plate boundary through the Pandora Trough area, and then declining through the rest of the Cainozoic. This change altered the original model moving the hypothetical curve closer to the vitrinite reflectance values (Plate 17). This model was further refined by doubling the amount of sediment removed by erosion during the Paleocene, and this resulted in both the computed and observed iso-vitrinite plots coinciding in part (Plate 18); increasing the erosion figure by a further 500m resulted in an even greater coincidence of the two plots, but the removal of 2500m of Cretaceous sediments during the Paleocene is considered to be highly improbable. An additional plot was made, where the unconformity between the ?Toro Sandstone and the Koi-Iange Formation was removed, and the sandstone was considered to be part of the Koi-Iange Formation; the figure of 1000m of eroded Cretaceous section was used as for the initial plot. The resulting model was little different to those noted above. To this model an additional heatflow event was placed in the Jurassic, but this too made little difference to the result. Thus the only way the two iso-vitrinite plots could be made to coincide was to add a thickness of sediment and to increase the heatflow regime at this site. However, it is difficult to envisage in excess of 2000m of erosion within the time constraints (Paleocene) within the region and a greater heatflow than used in these models. Even though the two iso-vitrinite plots coincide for Kusa-1, the heatflow was similarly modified to simulate the heatflow pulse due to the opening of the Coral Sea Basin, but because this well was sited further away from the heat source, the maximum heatflow levels were lower than those used for the Anchor Cay-1 model. This resulted in the two iso-vitrinite plots becoming separated (Plate 21), and it was only by using a lower modern day heatflow value of 1.2 h.f.u. that the two plots could be reconciled.

Capricorn Basin area: All locations have been modelled using a constant heatflow throughout time because of their distance from the heat source and is confirmed by modelling. Both Sandy Cape-1-3R and Wreck Island-1 show almost identical plots, but both differ from those for Aquarius-1 and Capricorn-1A, both of which are very similar. All sections show a gradual subsidence through the Eocene to Miocene, with the rate increasing rapidly in the Pliocene. The plots for Aquarius-1 and Capricorn-1A differ mainly in the thickness of sediment, which was sufficient to produce hypothetical (modelled) vitrinite reflectance values in excess of 0.5 in the Pliocene in Aquarius-1. The similarities between the two pairs of wells is to be expected since they penetrated similar stratigraphic sections. Both Aquarius-1 and Capricorn-1A penetrated a similar, poorly dated terrestrial section, which is absent in both Sandy Cape-1-3R and Wreck Island-1.

Coral Sea Basin: All sites were modelled using the same heatflow values as for the Papuan Basin, and with the heatflow being modelled as peaking between magnetic anomalies 27 and 22, gradually cooling until the present day. The plots show that water depths reached close to their present day depths soon after the end of this heatflow event. The sudden drop shown in the plots for the subsidence and decompacted burial curves for DSDP Site 287 is more artificial than real, due to the wide error bars for palaeodepth estimates based on the foraminiferids. All plots show a slow gradual subsidence through the Tertiary, rapidly increasing during the Pliocene.

Lord Howe Rise: The heatflow for these three DSDP Sites was modelled using slightly lower present day values (1.40 h.f.u.) than for the other sites. Also, the heatflow models were based on a heatflow peak at 76Ma (magnetic anomaly 33) because of the earlier opening of the adjacent Tasman Basin, gradually declining to the present day. The heatflow event is based on the rifting and breakup events which separated the Lord Howe Rise from the

Australian continent. The three sites show close similarities for all plots reflecting their similar geological histories.

Discussion

Though the four areas have different geological histories, one marked similarity for them all is the increase in the rate of subsidence in the Late Cainozoic which in the Papuan and Coral Sea Basins reflects flexuring of the margin due to collision and the development of a foreland basin. That a parallel increase in the rate of subsidence occurs in the Capricorn Basin and the southern end of the GBR is more difficult to explain but the timing of the event suggests that it is produced by intraplate stress effects related to tectonic events along the northern margin of Australia. Models of basin subsidence related to intraplate stress (Lambeck, 1983a, b) probably offer the most plausible explanation for associating distant collisional events to enhanced basin subsidence.

CONCLUSIONS

Though no one sequence used in this review provides a complete section, it is possible to construct such a section (Figure 3A, B). The study area spreads over in excess of 10° of latitude resulting in a significant temperature differential north to south which has effected faunal distribution since at least the late Oligocene when the region was well south of its present position. In addition the various sections on which this review is based are from a broad range of water depths, from abyssal depths below the CCD to near present day sea level. For these reasons it has been convenient to discuss the biostratigraphy in sections related by depth.

Marine sedimentation in the Papuan and Coral Sea Basins commenced in the middle Eocene, being linked to the formation of the Coral Sea Basin. In the northeastern Australian area, marine Eocene has been sampled only in the northern part of the Queensland Plateau (Chaproniere, 1983), but on seismic evidence such sediments may form a thin veneer over parts of both the Queensland and Marion Plateaux (Symonds and Davies, 1988). However, on the evidence from the wells drilled in the Capricorn Basin it is possible that non-marine Eocene sediments may be present on the southern parts of the Marion Plateau. To the north sedimentation continued into the late Eocene, and possibly into the basal Oligocene (as it did locally elsewhere in Papua New Guinea). There is evidence that water depths in the basin gradually increased with time, but in the shallow water sequences, depths decreased. Much of the Eocene sequence may have been removed during the early Oligocene erosion event which effected both shallow and deep water sequences. The rocks underlying the northerly sections range in age from Palaeozoic to Cretaceous, whereas in the south, they are probably Palaeozoic.

Sedimentation resumed again in the Late Oligocene, at first localised (during Zone P.21), but later (during Zones N.3 to N.4A) over the entire region. In the shallow water areas, larger foraminiferal-coralline algal limestones were deposited over the Eocene sediments. This shallow water sedimentation continued into the early and early middle Miocene; to the north water temperatures were sufficiently warm to permit coral development, but to the south, temperatures remained subtropical. In the north at least two phases of coral development took place, the first (in Pasca-C1) during upper Te (early Miocene) and the other (in Uramu-1A, Borabi-1 and Kusa-1) during lower Tf (late early or

early middle Miocene). In some sections (Anchor Cay-1) the late Oligocene sediments are dominated by planktic forms, grading upwards into larger foraminiferal carbonates. Thus all shallow water sections show evidence of shallowing. In the deeper water sequences depths remained stable. In the shallow sections, the early or early middle Miocene is truncated by a disconformity not present in the deeper water areas. The Miocene carbonates appear to have been exposed to subaerial weathering in most of the shelfal areas as seen in the wells of the southeastern Papuan Basin and in samples from the edge of the Marion Plateau. In some areas, particularly in the north, this erosional episode possibly resulted in the formation of solution channels (resulting in loss of drilling fluids). This hiatus appears to span an interval of approximately 15 ma., from at least within the early middle and including most of the late Miocene.

During the latest Miocene or early Pliocene sedimentation commenced in the shallow water areas over the eroded middle Miocene surface. However, in deeper water regions, sedimentation was continuous at most sites from the late Oligocene to the Holocene. In the Papuan Basin the initial sediments were dominated by carbonates, possibly reefal locally, but were soon overwhelmed by clastics. To the south carbonate sedimentation continued throughout the Pliocene to Holocene with only minor amounts of clastic material.

Three major and at least two minor unconformities are present in the sequence. The oldest is pre-middle Eocene and is present throughout the region, being probably related to the formation of the Coral Sea Basin. An early Oligocene unconformity is also universally present and in basinal areas is probably related to the initiation of corrosive Antarctic bottom waters linked with major glaciation in Antarctica (Keller et al., 1987), whilst in shallow areas it is due to erosion by currents resulting from major changes in oceanic circulation patterns due to the northward movement of Australia (Kennett et al., 1972). The third major unconformity represents most of middle and late Miocene time and effected only the shallow water areas, resulting in probable subaerial erosion and karst topography; this was almost certainly related to sea level fall. A short-lived, discontinuous, but nevertheless widespread unconformity is present within the Zone N.7 or N.8 (late early Miocene) interval, and again probably related to sea level fluctuation.

In the northern part of the area at least two periods of coral development occurred during the Miocene, and one in the Pliocene; to the south corals are definitely present only in the Pliocene to Holocene section. Thus in the northern area water temperatures during the early and early middle Miocene were warm enough to sustain coral growth with water depth and the presence of suitable substrate being the sole controls on reefal development. Because the latest early and earliest middle Miocene was a time when temperatures peaked in the region, it is possible that some reefal growth may have occurred at some locations in the southern part of the area reviewed.

Geohistory analyses have been made for all wells and DSDP Sites. These results suggest that heatflow regimes throughout the Tertiary cooled from those prevailing during the formation of the basins. Subsidence rates appear to have been gradual through Eocene to Miocene time, rapidly increasing during the Pliocene and Pleistocene for all basins. Sediments in the Papuan Basin became marginally mature by the end of the Mesozoic, but there is strong evidence that maturity levels increased during the Pliocene and Pleistocene. Within the Capricorn Basin, geohistory modeling indicates that levels in the lower part of Aquarius-1 approached marginal maturity only during the Pliocene or Pleistocene. Though the four areas have different geological histories, one marked similarity for them all is the increase in the rate of subsidence in the Late Cainozoic which in the Papuan and Coral Sea Basins reflects flexuring of the margin due to collision and the development of a foreland

basin. That a parallel increase in the rate of subsidence occurs in the Capricorn Basin and the southern end of the GBR is more difficult to explain but the timing of the event suggests that it is produced by intraplate stress effects related to tectonic events along the northern margin of Australia. Models of basin subsidence related to intraplate stress (Lambeck, 1983a, b) probably offer the most plausible explanation for associating distant collisional events to enhanced basin subsidence.

ACKNOWLEDGEMENTS

We gratefully acknowledge the help and advice given by Mr Michael Swift for the geohistory modelling, Ms Heike Struckmeyer for permitting us to use her interpretations of the Papuan Basin Mesozoic stratigraphy and Mr Samir Shafik for advice with interpretation of nannofossil biostratigraphic data. The drafting was carried out in the BMR Cartographic Services Unit by Mr Brian Pashley, Ms Debra O'Donoghue and Mr Larry Hollands.

REFERENCES

- Adams, C.G., 1970: A reconsideration of the East Indian Letter Classification of the Tertiary. *Bulletin of the British Museum (Natural History), Geology*, 19, 85-137.
- Adams, C.G., 1984: Neogene larger foraminifera, evolutionary and geological events in the context of datum planes. In: N. Ikebe and R. Tsuchi, Pacific Neogene datum Planes. Contributions to Biostratigraphy and Chronology. University of Tokyo Press. 47-67.
- Adams, C.G., Butterlin, J., and Samanta, B.K., 1986: Larger Foraminifera and events at the Eocene/Oligocene boundary in the Indo-E West Pacific region. In: C. Pomeroy and I. Premoli Silva (Eds), Terminal Eocene Events. 237-252. Elsevier, Amsterdam.
- Andrews, J.E., Packham, G., et al., 1975: Site Reports, Sites 285 to 289. *Initial Reports of the Deep Sea Drilling Project*, 30, 25-398. U.S. Government Printing Office, Washington.
- Armstrong, B.J., 1969: Phillips Australian Oil Company, Pasca C-1 Well Completion Report, 1-75.
- Belford, D.J., 1984: Tertiary foraminifera and age of sediments, Ok Tedi-Wabag, Papua New Guinea. *Bureau of Mineral Resources, Australia, Bulletin* 216, 1-52.
- Berggren, W.A., 1969: Paleogene biostratigraphy and planktonic foraminifera of northern Europe. In: P. Brönnimann and H.H. Renz (Eds), *Proceedings of the First International Conference on Planktonic Microfossils*, I, 121-160.
- Berggren, W.A., Kent, D.V., and Flynn, J.J., 1985a: Jurassic to Paleogene: Part 2. Paleogene geochronology and chronostratigraphy. In: N.J. Snelling (Ed.), *The Chronology of the Geological Record. The Geological Society of London, Memoir* 10, 141-195.
- Berggren, W.A., Kent, D.V., and van Couvering, J.A., 1985b: The Neogene: Part 2. Neogene geochronology and chronostratigraphy. In: N.J. Snelling (Ed.), *The Chronology of the Geological Record. The Geological Society of London, Memoir* 10, 211-250.
- Blow, W.H., 1969: Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. In: P. Brönnimann and H.H. Renz (Eds), *Proceedings of the First International Conference on Planktonic Microfossils*, I, 199-421.
- Blow, W.H., 1979: The Cainozoic Globigerinida. Vols. 1-3, pp. 1-1413. E.J. Brill, Leiden.
- Bolli, H.M., 1957: Planktonic foraminifera from the Oligocene - Miocene Cipero and Lengua Formations of Trinidad, B. W. I. *United States National Museum Bulletin* 215, 97-131.
- Bolli, H.M., 1966: The planktonic foraminifera in well Bodjonegoro-1 of Java. *Eclogae Geologicae Helvetiae*, 59, 449-465.
- Bolli, H.M., and Premoli Silva, I: 1973: Oligocene to Recent planktonic foraminifera and stratigraphy of the Leg 15 sites in the Caribbean Sea. In: N.T. Edgar, J.B. Saunders, et al., 1973, *Initial Reports of the Deep Sea Drilling Project*, 15, 475-497. U.S. Government Printing Office, Washington.
- Boersma, A., 1985: Biostratigraphy and biogeography of Tertiary bathyal foraminifers: Tasman Sea, Coral Sea, and on the Chatham Rise Deep Sea Drilling Project, Leg 90). In: J.P. Kennett and C.C. von der Borch et al., *Initial Reports of the Deep Sea Drilling Project*, 90, 1493-1517. U.S. Government Printing Office, Washington.
- Bukry, D., 1973: Low-latitude coccolith biostratigraphic zonation. In: N.J. Edgar, J.B. Saunders, and others, *Initial Reports of the Deep Sea Drilling Project*, 15, 685-703. U.S. Government Printing Office, Washington.
- Bukry, D., 1975: Coccolith and silicoflagellate stratigraphy, northwestern Pacific Ocean, Deep Sea Drilling Project Leg 32. In: R.L. Larson, R. Moberly, et al., *Initial Reports of the Deep Sea Drilling Project*, 32, 677-701. U.S. Government Printing Office, Washington.


- Burger, D., 1973: Spore zonation and sedimentary history of the Neocomian, Great Artesian Basin, Queensland. *Geological Society of Australia, Special Publication* 4, 87-118.
- Burger, D., 1988: Early Cretaceous environments in the Eromanga Basin; palynological evidence from GSQ Wyandra-1 corehole. *Association of Australasian Palaeontologists, Memoir* 5, 173-186.
- Burger, D., in prep a: Australian Phanerozoic Timescales: 9. Cretaceous: Geochronological charts and explanatory text. *Bureau of Mineral Resources, Australia, Record* 1989/39.
- Burger, D., in prep b: Australian Phanerozoic Timescales: 8. Jurassic: Geochronological charts and explanatory text. *Bureau of Mineral Resources, Australia, Record* 1989/38.
- Burns, B.J., 1977: Well Completion Report, Kusa-1, Papuan Basin, P.N.G., 1-200.
- Burns, B.J., and Bein, J., 1980: Regional geology and hydrocarbon potential of the Mesozoic of the western Papuan Basin, Papua New Guinea. *Apea Journal*, 1980, 1-15.
- Burns, R.E., and Andrews, J.E., 1973: Regional aspects of deep sea drilling in the southwest Pacific. In: R.E. Burns, J.E. Andrews, et al., *Initial Reports of the Deep Sea Drilling Project*, 21, 897-906. U.S. Government Printing Office, Washington.
- Burns, R.E., Andrews, J.E., van der Lingen, G.J., Churkin, M., jnr., Galehouse, J.S., Packham, G.H., Davies, T.A., Kennett, J.P., Dumitrica, P., Edwards, A.R., and von Herzen, R.P., 1973a: Site 209. In: R.E. Burns, J.E. Andrews, et al., *Initial Reports of the Deep Sea Drilling Project*, 21, 333-367. U.S. Government Printing Office, Washington.
- Burns, R.E., Andrews, J.E., van der Lingen, G.J., Churkin, M., jnr., Galehouse, J.S., Packham, G.H., Davies, T.A., Kennett, J.P., Dumitrica, P., Edwards, A.R., and von Herzen, R.P., 1973b: Site 210. In: R.E. Burns, J.E. Andrews, et al., *Initial Reports of the Deep Sea Drilling Project*, 21, 369-440. U.S. Government Printing Office, Washington.
- Cameron, D.H., 1975: Carbon and carbonate analyses, Leg 30. In: J.E. Andrews, G. Packham, et al., *Initial Reports of the Deep Sea Drilling Project*, 30, 687-688.
- Carlsen, C.T., and Wilson, T.C., 1968a: Gulf-A.O.G. Capricorn No. 1A well completion report. (Unpublished).
- Carlsen, C.T., and Wilson, T.C., 1968b: Gulf-A.O.G. Aquarius No. 1 well completion report. (Unpublished).
- Caron M., 1985: Cretaceous planktonic foraminifera. In: H.M. Bolli, J.B. Saunders and K. Perch-Nielson (Eds.), *Plankton Stratigraphy*, 17-86. Cambridge University Press, Cambridge.
- Chaproniere, G.C.H., 1975: Palaeoecology of Oligo-Miocene larger Foraminifera, Australia. *Alcheringa*, 1, 37-58.
- Chaproniere, G.C.H., 1976: A new rock-stratigraphic unit from the Carnarvon Basin, western Australia. *BMR Journal of Australian Geology and Geophysics*, 1, 171-174.
- Chaproniere, G.C.H., 1980: Biometrical studies of Early Neogene larger Foraminifera from Australia and New Zealand. *Alcheringa*, 4, 153-181.
- Chaproniere, G.C.H., 1981: Late Oligocene to early Miocene planktic Foraminifera from Ashmore Reef No.1 Well, northwest Australia. *Alcheringa*, 5, 103-131.
- Chaproniere, G.C.H., 1983: Tertiary larger foraminifera from the northwestern margin of the Queensland Plateau, Australia. *Bureau of Mineral Resources, Australia, Bulletin* 217, 61-57.
- Chaproniere, G.C.H., 1984: Oligocene and Miocene larger Foraminifera from Australia and New Zealand. *Bureau of Mineral Resources, Australia, Bulletin* 188, 1-98.

- Chaproniere, G.C.H., 1985a: Late Neogene and Quaternary planktonic foraminiferal biostratigraphy and paleobathymetry of dredge samples from the southern Tongan Platform (Cruise L5-82-SP). In: D.W. Scholl and T.L. Vallier (Eds.), *Geology and Offshore Resources of Pacific Island Arcs - Tonga Region. Circum-Pacific Council for Energy and Mineral Resources Earth Science Series*, 2, 131-140. Circum-Pacific Council for Energy and Mineral Resources, Houston.
- Chaproniere, G.C.H., 1985b: Late Tertiary and Quaternary foraminiferal biostratigraphy and paleobathymetry of cores and dredge samples from Cruise KK820316 Leg 2. In: T.M. Brocher (Ed.), *Geological Investigations of the Northern Melanesian Borderland. Circum-Pacific Council for Energy and Mineral Resources Earth Science Series*, 3, 103- 122. Circum-Pacific Council for Energy and Mineral Resource, Houston, Texas.
- Chaproniere, G.C.H., in prep: Middle Pleistocene to Holocene planktic foraminiferal biostratigraphy of the Coral Sea, offshore Queensland, Australia.
- Chaproniere, G.C.H., in press: Late Eocene to Pleistocene foraminiferal biostratigraphy and paleobathymetry of dredge samples from the southern Tonga Platform (Cruise L3-84-SP). In: P.F. Ballance, R.H. Herzer and T.L. Vallier (Eds.), *Contributions to the Geology of the Tonga and Lau Regions of the Southwest Pacific. Circum-Pacific Council for Energy & Mineral Resources, Earth Science Series*, __, __ - __. Circum-Pacific Council for Energy and Mineral Resources, Houston, Texas, U.S.A.
- Condon, M.A., Johnstone, D., Perry, W.J., and Crespin, I., 1953: The Cape Range Structure, Western Australia, 1st edition. *Bureau of Mineral Resources, Australia, Bulletin* 21.
- Cull, J.P., and Conley, D., 1983: Geothermal gradients and heat flow in Australian sedimentary basins. *BMR Journal of Australian Geology and Geophysics*, 8, 329-337.
- Davey, R.J., 1987: Palynological zonation of the early Cretaceous and late Jurassic of Papua New Guinea. *Geological Survey of Papua New Guinea, Memoir* 13, 1-53.
- Davies, P.J., Symonds, P.A., Feary, D.A., and Pigram, C.J., 1987: Horizontal plate motion: a key allocyclic factor in the evolution of the Great Barrier Reef. *Science*, 238, 1697-1700.
- Derrington, S.S., 1960: H.B.R. No.1 Bore Wreck Island, Queensland. *Bureau of Mineral Resources, Australia, Petroleum Search and Subsidy Acts Publication* 4, 1-15.
- Dettmann, M.E., 1986: Early Cretaceous palynoflora of subsurface strata correlative with the Koonwarra Fossil Bed, Victoria. *Association of Australasian Palaeontologists, Memoir* 3, 79-110.
- Dettmann, M.E., and Playford, G., 1969: Palynology of the Australian Cretaceous: a review. In: K.S.W. Campbell (Ed.), *Stratigraphy and Palaeontology. Essays in honour of Dorothy Hill*, 174-210. Australian National University Press, Canberra.
- Drooger, C.W., 1952: Study of American Miogypsinidae. Vonk & Co., Zeist.
- Ericson, E.K., 1976: Capricorn Basin. In: R.B. Leslie, H.J. Evans & C.L. Knight, (Eds.), *Economic Geology of Australia and Papua New Guinea*, 3, Petroleum. *Australian Institute of Mining and Metallurgy Monograph*, 7, 464-473.
- Ewing, M., Hawkins, L.V., and Ludwig, W.J., 1970: Crustal structure of the Coral Sea. *Journal of Geophysical Research*, 75, 1953-1962.
- Falvey, D.A., and Deighton, I., 1982: Recent advances in burial and thermal geohistory analysis. *APEA Journal*, 22, 65-81.
- Feary, D.A., Davies, P.J., Pigram, C.J., and Symonds, P.A., in press: Climatic evolution and control on carbonate deposition in northeast Australia. *Global and Planetary Change*.
- Foster, C.B., 1982: Illustrations of early Tertiary (Eocene) plant microfossils from the Yaamba Basin, Queensland. *Geological Survey of Queensland, Publication*, 381, 1-33.


- Fourcade, E., and Butterlin, J., 1988: Reworked and redeposited larger foraminifers on slopes and in basins of the Bahamas, Leg 101. In: J.A. Austin, jnr., W. Schlanger, et al., *Proceedings of the Ocean Drilling Program, Scientific Results*, 101, 47-61.
- Grimes, K.G., 1982: Stratigraphic drilling report: GSQ Sandy Cape 1-3R. *Geological Survey of Queensland, Record Series*, 1981/45, 1-40.
- Haq, B.U., Hardenbol, J., and Vail, P.R., 1988: Mesozoic and Cenozoic chronostratigraphy and cycles of sea-level change. In: *Sea-Level Change - An Integrated Approach. Society of Economic Paleontologists and Mineralogists, Special Publication 42*, 71-108.
- Hekel, H., 1972: Pollen and spore assemblages from Queensland Tertiary sediments. *Geological Survey of Queensland, Publication*, 355, 1-34.
- Helby, R.J., Morgan, R., and Partridge, A.D., 1987: A palynological zonation of the Australian Mesozoic. *Memoir of the Association of Australasian Palaeontologists* 4, 1-94.
- Helby, R.J., and Partridge, A.D., 1976: Palynological analysis of core samples from the Mesozoic in Anchor Cay-1. Esso Australia Ltd., Palaeontological Report 1976/20, 1-3. (Unpublished).
- Hill, K.C., and Gleadow, A.J.W., 1989: Uplift and thermal history of the Papuan Fold Belt, Papua New Guinea: Apatite fission track analysis. *Australian Journal of Earth Sciences*, 515-539.
- Ingram, B.S., and Morgan, R., 1988: The development and status of the Mesozoic palynostratigraphy of the North West Shelf, Australia. In: P.G. Purcell & R.R. Purcell (Eds.), *The North West Shelf, Australia: Proceedings of the Petroleum Exploration Society of Australia Symposium, Perth 1988*, 581-590.
- Jenkins, D.G., Bowen, D.Q., Adams, C.G., Shackleton, N.J., and Brassil, S.C., 1985: The Neogene: Part 1. In: N.J. Snelling (Ed.), *The Chronology of the Geological Record. Geological Society of London, Memoir* 10, 199-210.
- Jenkins, D.G., and Srinivasan, M.S., 1985: Cenozoic planktonic foraminifers from the equator to the sub-Antarctic of the southwest Pacific. In: J.P. Kennett and C.C. von der Borch et al., *Initial Reports of the Deep Sea Drilling Project*, 90, 795-834. U.S. Government Printing Office, Washington.
- Keller, G., Herbert, T., Dorsey, R., D'Hondt, S., Johnsson, M., and Chi, W.R., 1987: Global distribution of late Paleogene hiatuses. *Geology*, 15, 199-203.
- Kennett, J.P., 1973: Middle and late Cenozoic planktonic foraminiferal biostratigraphy of the Southwest Pacific - DSDP Leg 21. In: R.E. Burns, J.E. Andrews, et al., *Initial Reports of the Deep Sea Drilling Project*, 21, 575-639. U.S. Government Printing Office, Washington.
- Kennett, J.P., Burns, R.E., Andrews, J.E., Churkin, M., jnr., Davies, T.A., Dumitrica, P., Edwards, A.R., Galehouse, J.S., Packham, G.M., and van der Lingen, G.J., 1972: Australian-Antarctic continental drift, palaeocirculation changes and Oligocene deep-sea erosion. *Nature (Physical Science)*, 239, 51-55.
- Kennett, J.P., and Srinivasan, M.S., 1983: Neogene Planktonic Foraminifera. A Phylogenetic Atlas, 1-265. Hutchinson Ross, Stroudsburg, Pennsylvania.
- Kennett, J.P., and von der Borch, C.C., 1985: Southwest Pacific Cenozoic paleoceanography. In: J.P. Kennett and C.C. von der Borch et al., *Initial Reports of the Deep Sea Drilling Project*, 90, 1493-1517. U.S. Government Printing Office, Washington.
- Kennett, J.P., von der Borch, C.C., et al., 1985: *Initial Reports of the Deep Sea Drilling Project*, 90, 1-744. U.S. Government Printing Office, Washington.
- Lambeck, K., 1983a: The role of compressive forces in intracratonic basin formation and mid-plate orogenies. *Geophysical Research Letters*, 10, 845-848.
- Lambeck, K., 1983b: Structure and evolution of the intracratonic basins of central Australia. *Geophysical Journal of the Royal Astronomical Society*, 74, 843-886.

- Lloyd, A.R., 1968: Palaeontological Report, Uramu No.1A Well, Papua. In: A.A. Slanis & R.K. Netzel, Phillips et al. *Uramu No.1A, Well Completion Report*, 1-4.
- Lloyd, A.R., 1978: An outline of the Tertiary paleontology and stratigraphy of the Gulf of Papua, Papua New Guinea. *Proceedings of the Regional Conference on the Geology and Mineral Resources of SE Asia*, 43-54.
- Lloyd, A.R., 1988: The Geology and Hydrocarbon Potential of Southern Papua New Guinea. Unpublished Report. 1-270.
- McDonagh, G.P., and Knuth, B.W., 1968: Phillips Australian Oil Company, Borabi No.1 Well Completion Report, 1-17.
- Marshall, J.F., 1971: Phosphatic sediments on the eastern Australian upper continental slope. *Bureau of Mineral Resources, Australia, Record* 1971/59.
- Marshall, J.F., 1972: Morphology of the east Australian continental margin between 21° S and 33° S. *Bureau of Mineral Resources, Australia, Record* 1972/70.
- Martini, E., 1971: Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: A. Farinacci (Ed.), *Proceedings of the Second Planktonic Conference, Roma 1970*, 739-785. Edizioni Tecnoscienza, Roma.
- Morgan R., 1989: Overview and hybrid palynology zonation for the late Mesozoic of Papua New Guinea. Unpublished Report for Papex, 1-18.
- Oppel, T.W., 1969: Tenneco-Signal Anchor Cay number 1, offshore Queensland, Well Completion Report, 1-15.
- Palmieri, V., 1971: Tertiary subsurface biostratigraphy of the Capricorn Basin. *Geological Survey of Queensland, Report* 52, 1-18.
- Palmieri, V., 1974: Correlation and environmental trends of the subsurface Tertiary Capricorn Basin. *Geological Survey of Queensland, Report* 86, 1-14.
- Palmieri, V., 1975: Planktonic Foraminiferida from the Capricorn Basin, Queensland. *Geological Survey of Queensland, Palaeontological Papers*, 36, 1-47.
- Palmieri, V., 1984: Neogene Foraminiferida from GSQ Sandy Cape 1-3R bore, Queensland: a biostratigraphic appraisal. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 46, 165-183.
- Partridge, A.D., 1976: The geological expression of eustasy in the early Tertiary of the Gippsland Basin. *APEA Journal*, 16, 73-79.
- Pigram, C.J., Davies, P.J., Feary, D.A., Symonds, P.A., and Chaproniere, G.C.H., 1990: Controls on Tertiary carbonate platform evolution in the Papuan Basin: New play concepts. *Proceedings of the First Papua New Guinea Petroleum Symposium*, 185-195.
- Pigram, C.J., Davies, P.J., Feary, D.A., Symonds, P.A., and Chaproniere, G.C.H., in prep.: Absolute magnitude of Miocene sea-level fall from carbonate platforms on the Marion Plateau, NE Australia.
- Robertson Research Australia Ltd., and Flower Doery Bushan Ltd., 1984: Petroleum potential of the Papuan Basin, Papua New Guinea. Report prepared for the Geological Survey of Papua New Guinea. 4 volumes text and appendices, 2 volumes maps and seismic sections.
- Roth, P.H., 1978: Cretaceous nannoplankton biostratigraphy and oceanography of the northwestern Atlantic Ocean. In: W.E. Benson, R.E. Sheridan, et al., *Initial Reports of The Deep Sea Drilling Project*, 44, 731-759. U.S. Government Printing Office, Washington.
- Shackleton, N.J., and Opdyke, N.D., 1976: Oxygen isotope and paleomagnetic stratigraphy of Pacific core V28-239: Late Pliocene to latest Pleistocene. In: R.M. Cline & J.D. Hays (Eds.), *Investigations of Late Quaternary Paleooceanography and Paleoclimatology. Memoir of the Geological Society of America*, 145, 449-464.
- Sissingh, W., 1977: Biostratigraphy of Cretaceous calcareous nannoplankton. *Geologie en Mijnbouw*, 56, 433-440.

- Slanis, A.A., and Netzel, R.K., 1968: Phillips Australian Oil Company, Uramu 1A Well Completion Report, 1-18.
- Sleep, N.H., 1971: Thermal effects of the formation of Atlantic continental margins by continental breakup. *Geophysical Journal of the Royal Astronomical Society*, 25, 325-350.
- Stainforth, R.M., Lamb, J.L., Luterbacher, H., Beard, J.H., and Jeffords, R.M., 1975: Cenozoic planktonic foraminiferal zonation and characteristics of index forms. *The University of Kansas Paleontological Contributions, Article 62*, 1-425.
- Stewart, W.D., and Durkee, E.F., 1985: Petroleum potential of the Papuan Basin. *Oil and Gas Journal*, 1985,
- Stover, L.E., and Evans, P.R., 1973: Upper Cretaceous-Eocene spore-pollen zonation, offshore Gippsland Basin. *Geological Society of Australia, Special Publication 4*, 55-72.
- Stover, L.E., and Partridge, A.D., 1973: Tertiary and Late Cretaceous spores and pollen from the Gippsland Basin, southeastern Australia. *Proceedings of the Royal Society of Victoria*, 85, 237-286.
- Symonds, P.A., and Davies, P.J., 1988: Structure, stratigraphy, evolution and regional framework of the Townsville Trough and Marion Plateau region. *Bureau of Mineral Resources, Australia, Record 1988/48*, 1-47.
- Symonds, P.A., Fritsch, J., and Schluter, H., 1984: Continental margin around the western Coral Sea Basin: Structural elements, seismic sequences and petroleum geological aspects. In: S.T. Watson, (Ed.), Transactions of the Third Circum-Pacific Energy and Mineral Resources Conference, Hawaii, 143-252. American Association of Petroleum Geologists, Tulsa.
- Weissel, J.K., and Watts, A.B., 1979: Tectonic evolution of the Coral Sea Basin. *Journal of Geophysical Research*, 84, 4572-4582.
- Williamson, P.E., and Pigram, C.J., 1986: Geohistory, basin subsidence and maturation in the Bass Basin. In: R.C. Glenie (Ed.), *The Second South-Eastern Australian Oil Exploration Symposium*, 285-302.




1 Berggren et al., (1985a, b), Burger, (in prep.a)
2 Blow (1969, 1979), Breggen (1969)
3 Boli (1957, 1966), Stainforth et al., (1975),
Boli & Premoli-Silva (1973), Caron (1985)
4 Adams (1984), Adams et al., (1986), Chaproniere
(1981), (1983), Jenkins et al., (1985)
5 Bukry (1973, 1975), Martini (1971), Roth (1978), Sissingh (1977)
6 Burger (1973), 1988, in prep. a), Dettmann (1986),
Dettmann & Playford (1969), Helby & Morgan (1987),
Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976),
Stover & Evans (1973), Stover & Partridge (1973)
Compiled from:
Oppel (1969); Helby & Partridge (1976) Inner neritic 0-20m Outer neritic 100-200m
Robertson Research & Flower Doery Mid neritic 20-100m Slope 200m+
Bucha (1984), Lloyd (1988)
Record 1990/7
TOC: Total Organic Carbon (%)
St: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)
TMAX: Temperature at which S2 signal is max (°C)
PI: Production index (S1-S2)
HI: Hydrogen index (mg hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock) (S1-S2)
VR: Vitrinite reflectance
20/00/32



(1) Time scale after Burger (in prep. b.)
(2) after Helby, Morgan & Portridge (1987)

Latitude: 9° 26' 29.43"S Date drilled: Feb-May, 1969
Longitude: 144° 03' 31.00"E Company: Tenneco



Compiled from:
Oppel (1969), Burns & Bein (1980)
Lloyd (1988), Robertson Research &
Flower Doery Buchan (1984)
Struckmeyer (pers.com. 1990)

Inner neritic 0-20m
Mid neritic 20-100m
Slope 200m+

Record 1990/7

TOC: Total Organic Carbon (%)
St: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)
PI: Production index (S1-S2)
HI: Hydrogen index (mg hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock)(S1+S2)
VR: Vitrinite Reflectance
20/00/32

Chronostratigraphy	Age/ Depth Correlation	Depth (m) Drill Sub- sea	Formation	Two-way time	Lithology
		10	Seafloor		
		200			
		400			
		616	Era Beds		
		707		0.80	
		800			
		1000		1.03	
		1200			
		1220		1.21	
		1340		1.31	Bioclastic limestone
		1400			
		1510			
		1560		1.41	Dolomitic bioclastic limestone
		1600			
		1800	Borabi		
		2000	Reef		Bioclastic limestone
		2095		1.61	
		2150			
		2200	Limestone		
		2340		1.75	
		2400			
		2580		1.88	
		2600			
		2800	Mendi Group		Bioclastic limestone
		2820		1.98	
		2835			
		2879	?Maril Shale	2.00	Dark grey pyritic shale
		TD			
		3000			
		3200			
		3400			
		3600			
		3800			

1 Berggren et al., (1985a, b), Burger, (in prep.a)	5 Bukry (1973), 1975), Martini (1971), Roth (1978), Sissingh (1977)	Compiled from:	Inner neritic 0-20m	Outer neritic 100-200m	TOC:Total Organic Carbon(%)	TMAX:Temperature at which S2 signal is max (°C)
2 Blow (1969, 1979), Breggren (1969)	6 Burger (1973), 1988, (in prep. a), Dettmann (1986),	McDonagh & Knuth (1968)	Mid neritic 20-100m	Slope 200m+	StPyrolysis free-hydrocarbon signal	Pt:Production index (S1-S2)
3 Bolli (1957, 1966), Stainforth et al., (1975),	Dettmann & Playford (1969), Helby & Morgan (1987),	Robertson Research & Flower Doery			(mg S2 hydrocarbons/g rock)	Hi:Hydrogen index (ma hydrocarbons/g organic carbon)
Bolli & Premoli-Silva (1973), Caron (1985)	Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976),	Buchan (1984)			S2:Pyrolysis kerogen signal	GP:Genetic potential (Kg hydrocarbons/ton rock)(S1+S2)
4 Adams (1984), Adams et al., (1986), Chaproniere	Stover & Evans (1973), Stover & Partridge (1973)				(mg S2 hydrocarbons/g rock)	
(1981), (1983), Jenkins et al., (1985)						20/00/33

(1) Time scale after Burger (in prep. b.)
(2) after Helby, Morgan & Portridge (1987)

Compiled from:
McDonagh & Knuth (1968)
Robertson Research & Flower Doery
Buchan (1984)
Struckmeyer (pers.com. 1990), Lloyd (1988)


Inner neritic 0-20m	Outer neritic 100-200m
Mid neritic 20-100m	Slope 200m+


TOC: Total Organic Carbon (%)
S1: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)
 PI: Production index (S1-S2)
 HI: Hydrogen index (ma hydrocarbons/g organic carbon)
 GP: Genetic potential (Kg hydrocarbons/ton rock)(S1+S2)



20/OQ/33

[illegible]

Latitude: 8° 42' 28.52" S Date drilled: Feb-April, 1977
Longitude: 144° 03' 02.88" E Company: Esso



Latitude: 8° 42' 28.52" S Date drilled: Feb-April, 1977
Longitude: 144° 03' 02.88" E Company: Esso






(1) Time scale after Burger (in prep. b.)
(2) after Helby, Morgan & Portridge (1987)

Compiled from:
Burns (1977); Burns & Bein (1980); Lloyd (1988)
Struckmeyer (pers.com. 1990)
Robertson Research & Flower Doery Buchan (1984)
Record 1990/7

Inner neritic 0-20m
Mid neritic 20-100m
Outer neritic 100-200m
Slope 200m+

TOC: Total Organic Carbon (%)
S1: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)
P1: Production index (S1-S2)
H1: Hydrogen index (ma hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock)(S1+S2)
VR: Vitrinite reflectance
20/OA/34



1 Berggren et al., (1985a, b). Burger, (in prep. a)
2 Blow (1969, 1979), Berggren (1969)
3 Bolli (1957, 1966), Stainforth et al., (1975),
Bolli & Premoli-Silva (1973), Caron (1985)
4 Adams (1984), Adams et al., (1986), Chaproniere
(1981), (1983), Jenkins et al., (1985)
5 Bukry (1973), 1975, Martini (1971), Roth (1978), Sissingh (1977)
6 Burger (1973), 1988, in prep. a), Dettmann (1986),
Dettmann & Playford (1969), Helby & Morgan (1987),
Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976),
Stover & Evans (1973), Stover & Partridge (1973)

Compiled from: Armstrong (1969)
Lloyd (1988); Robertson Research &
Flower Doery Buchan (1984)
Struckmeyer (pers.com. 1990)
Inner neritic 0-20m
Mid neritic 20-100m
Outer neritic 100-200m
Slope 200m+
Record 1990/7

TOC: Total Organic Carbon (%)
St: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)
TMAX: Temperature at which S2 signal is max (°C)
P1: Production index (S1-S2)
H1: Hydrogen index (ma hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock) (S1+S2)
20/09/35

Berggren et al., (1985a, b), Burger, (in prep.a)	5 Bukry (1973), 1975, Martini (1971), Roth (1978), Sissingh (1977)	Compiled from: Lloyd (1968,1988)	Inner neritic 0-20m	Outer neritic 100-200m	TOC:Total Organic Carbon(%)		TMAX:Temperature at which S2 signal is max (°C)
2 Blow (1969, 1979), Breggren (1969)	6 Burger (1973), 1988, in prep. a, Dettmann (1986).	& Netzel (1968)	Mid neritic 20-100m	Slope 200m+	StPyrolysis free-hydrocarbon signal		Pt:Production index (S1-S2)
3 Bolli (1957, 1966), Stainforth et al., (1975), Bolli & Premoli-Silva (1973), Caron (1985)	Dettmann & Playford (1969), Helby & Morgan (1987), Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976), Stover & Evans (1973), Stover & Partridge (1973)	Robertson Research & Flower Doery Buchan (1984); Struckmeyer (pers.com.)			(mg S2 hydrocarbons/g rock)		Hl:Hydrogen index (ma hydrocarbons/g organic carbon)
4 Adams (1984), Adams et al., (1986), Chaproniere (1981), (1983), Jenkins et al., (1985)		Record 1990/7			S2:Pyrolysis kerogen signal (mg S2 hydrocarbons/g rock)		GP:Genetic potential (Kg hydrocarbons/ton rock)(S1+S2) 20/OQ/36

AGE (MA)	Magnetost­rati­graphy	ERA	SYSTEM	(1) STAGE	BIOSTRATIGRAPHY (2)				AGE (MA)
					Dinoflagellate zones	Spore-Pollen zones			
						West	East&South		
130		MESOZOIC	JURASSIC	BERRIASIAN	<i>K. wismaniae</i>	<i>Microcaryophytes</i>	<i>B. eneabbaensis</i>	<i>C. australiensis</i>	130
131				<i>P. iehiense</i>					
TITHONIAN				<i>D. jurassicum</i>	<i>R. watheroensis</i>		<i>R. watheroensis</i>	135.5	
				<i>D. montgomeryi</i>					
KIMMERIDGIAN				<i>C. perforans</i>				140	
				<i>D. swanense</i>					
OXFORDIAN				<i>W. clathrata</i>	<i>M. florida</i>		<i>M. florida</i>	140	
				<i>W. spectabilis</i>					
CALLOVIAN				<i>R. aemula</i>				150	
				<i>W. digitata</i>					
BATHONIAN				<i>W. indolata</i>	<i>C. cooksoniae</i>		<i>C. cooksoniae</i>	160	
				<i>C. halosa</i>					
BAJOCIAN				<i>D. caddaense</i>	upper <i>D. complex</i>		<i>D. complex</i>	170	
					lower				
AALENIAN					<i>C. turbatus</i>		<i>C. turbatus</i>	180	
TOARCIC								190	
PLIENS-BACHIAN					<i>C. torosa</i>		<i>C. torosa</i>	200	
SINEMURIAN				210					
HETTANGIAN				220					
RHAETIAN		<i>A. reducta</i>		230					
NORIAN		<i>M. crenulatus</i>	<i>P. crenulatus</i>	240					
CARNIAN			<i>C. rotundus</i>	250					
LADINIAN		<i>S. speciosus</i>	<i>S. speciosus</i>	260					
ANSIAN		<i>S. quadrifidus</i>	<i>S. quadrifidus</i>	270					
SCYTHIAN		<i>A. parvispinosus</i>		280					
		<i>T. playfordii</i>	<i>A. tenuispinosus</i>						
TATARIAN		<i>P. samoilovichii</i>	<i>P. samoilovichii</i>	290					
		<i>L. pellucidus</i>	<i>L. pellucidus</i>						
		<i>P. microcorpus</i>	<i>P. microcorpus</i>	300					

(1) Time scale after Burger (in prep. b.)
(2) after Helby, Morgan & Portridge (1987)

Latitude: 7°48'22.8"S Date drilled: Nov-Dec 1967
Longitude: 144°41'39.4"E Company: Phillips Australia

Chronostratigraphy	Age/ Depth Correlation	Depth (m) Drill	Sub- sea	Formation	Two-way time	Lithology	Depositional environment					Remarks	Geochemical data							Depth (m)					
							Locustrine	Estuarine	Inner	Mid	Outer		Neritic	Neritic	Neritic	Slope	TOC	S1	S2		TMAX	P1	H1	GP	
																								0	
																									200
																									400
																									600
																									800
																									1000
																									1200
																									1400
																									1600
																									1800
																									2000
																									2200
																									2400
																									2600
																									2800
																									3000
																									3200
																									3400
																									3600
																									3800


Compiled from:
Lloyd (1968,1988) & Netzel (1968)
Robertson Research & Flower Doery
Buchan (1984); Struckmeyer (pers.com.)




Inner neritic 0-20m
Mid neritic 20-100m
Outer neritic 100-200m
Slope 200m+

Record 1990/7

TOC: Total Organic Carbon (%)
St: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)
P1: Production index (S1-S2)
H1: Hydrogen index (mg hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock)(S1+S2)
20/OQ/36






1 Berggren et al., (1985a, b), Burger, (in prep.)
2 Blow (1969, 1979), Berggren (1969)
3 Bolli (1957, 1966), Stainforth et al., (1975),
Bolli & Premoli-Silva (1973), Caron (1985)
4 Adams (1984), Adams et al., (1986), Chaproniere
(1981), (1983), Jenkins et al., (1985)
5 Bukry (1973), 1975, Martini (1971), Roth (1978), Sissingh (1977)
6 Burger (1973), 1988, in prep. a), Dettmann (1986),
Dettmann & Playford (1969), Helby & Morgan (1987),
Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976),
Stover & Evans (1973), Stover & Partridge (1973)
Compiled from:
Carlsen & Wilson (1968 a)
Palmieri (1971, 1974, 1975)
Ericson (1976), Hekel (1972)
Record 1990/7
Inner neritic 0-20m
Mid neritic 20-100m
Outer neritic 100-200m
Slope 200m+
TOC: Total Organic Carbon (%)
St: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)
TMAX: Temperature at which S2 signal is max (°C)
PI: Production index (S1-S2)
HI: Hydrogen index (mg hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock) (S1-S2)
20/00/38

AGE (MA)	Magnetic polarity	ERA	System	EPOCH	(1) STAGE	International Foraminiferal zonation (2)	(3)	Calcareous nannofossils (4)	(5)	Palynoflora (6)	AGE (MA)
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N23	<i>Globobulimina calida</i>	NN15	NN15		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N22	<i>Globobulimina truncatulinoides</i>	NN14	NN14		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N21	<i>Globobulimina tosaensis</i>	NN13	NN13		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N19/20	<i>Globobulimina micrantha</i>	NN12	NN12		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N18	<i>Globobulimina margaritae</i>	NN11	NN11		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N17	<i>Neoglobobulimina duterrei</i>	NN10	NN10		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N16	<i>Neoglobobulimina decussata</i>	NN9	NN9		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	10.5	<i>G. menardi</i>	NN8	NN8		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	SERRAVALIAN	<i>G. mayeri</i>	NN7	NN7		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N12	<i>G. fashi robusta</i>	NN6	NN6		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N11	<i>G. fashi lobata</i>	NN5	NN5		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N10	<i>G. fashi fashi</i>	NN4	NN4		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N9	<i>Præobulimina glomerata</i>	NN3	NN3		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N8	<i>Globobulimina insueta</i>	NN2	NN2		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N7	<i>Catapsydrax stainforthi</i>	NN1	NN1		
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N6	<i>Catapsydrax dissimilis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N5	<i>Globobulimina kugleri</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	N4	<i>Globobulimina cipoensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P22	<i>Globobulimina opima</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P21	<i>Globobulimina ampliapertura</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P19/20	<i>Cassigerina chipolensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P18	<i>Pseudohastigerina micra</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P17	<i>Globobulimina cerroazulensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P16	<i>Porticulasphera seminivoluta</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P15	<i>Truncatuloides rahri</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P14	<i>Orbulinoides beckmanni</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P12	<i>Morozovella lehneri</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P11	<i>Globigerina subglobulata</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P10	<i>Hantkenina aragonensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P9	<i>Acarina pentacamerata</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P8	<i>Morozovella aragonensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P7	<i>Morozovella formosa</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P6	<i>M. subbotinae</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P5	<i>Morozovella edgari</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P4	<i>Morozovella velascoensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P3	<i>Planorbulina pseudomenardi</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P2	<i>Planorbulina pusilla</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P1	<i>Morozovella uncinata</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P0	<i>Subbotina trinidadensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-1	<i>Subbotina pseudobulimina</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-2	<i>Globigerina eugubina</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-3	<i>Abathomphalus mayaroensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-4	<i>Gansserina gansseri</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-5	<i>Globotruncana aegyptiaca</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-6	<i>Globotruncana havanensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-7	<i>Globotruncana calcarata</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-8	<i>Globotruncana ventricosa</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-9	<i>Globotruncana elevata</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-10	<i>Dicarinella asymetrica</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-11	<i>Dicarinella concavata</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-12	<i>Dicarinella primitiva</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-13	<i>Helvetoglobotruncana helvetica</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-14	<i>Whiteinella archeocretacea</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-15	<i>Rotalipora cushmani</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-16	<i>Rotalipora reicheli</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-17	<i>Rotalipora brotzeni</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-18	<i>Rotalipora appenninica</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-19	<i>Rotalipora ticiensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-20	<i>Rotalipora subticinensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-21	<i>Ticinella breggiensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-22	<i>Ticinella primula</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-23	<i>Ticinella bejovensis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-24	<i>Hedbergella gorbachikae</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-25	<i>Globigerinella algeriana</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-26	<i>Schackina cabri</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-27	<i>Globigerinella blowi</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-28	<i>Hedbergella sigali</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-29	<i>Globigerina hoterivica</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-30	minute planktic foraminiferids				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-31	<i>Capionellids</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-32	<i>Capionellites</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-33	<i>Capionellopsis</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-34	<i>Capionella elliptica</i>				
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-35					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-36					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-37					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-38					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-39					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-40					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-41					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-42					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-43					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-44					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-45					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-46					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-47					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-48					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-49					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-50					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-51					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-52					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-53					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-54					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-55					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-56					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-57					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-58					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-59					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-60					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-61					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-62					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-63					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-64					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-65					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-66					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-67					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-68					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-69					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-70					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-71					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-72					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-73					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-74					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-75					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-76					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-77					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-78					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-79					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-80					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-81					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-82					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-83					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-84					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-85					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-86					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-87					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-88					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-89					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-90					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-91					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-92					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-93					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-94					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-95					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-96					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-97					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-98					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-99					
2.4		Quaternary	Quaternary	Quaternary	Quaternary	P-100					

Latitude: 24° 42'00"S Date drilled: Oct 1980
Longitude: 153° 15'30"E Company: Geol. Survey of Queensland



Depositional environment							Well: Sandy Cape 1-3R							PLATE 8			
Locustine	Estuarine	Inner	Neritic	Mid	Neritic	Outer	Neritic	Slope	Remarks	Geochemical data							Depth (m)
										TOC	S1	S2	TMAX	P1	H1	GP	0
																	100
									Corals with bethic forams (<i>Alv. quoyi</i> ; <i>Calc. spengleri</i>)								
									Bethics more numerous								
									base <i>Pu. primalis</i>								
									Planktics common								200
									Planktics forams (<i>Ga. nepenthes</i>) with rare larger foraminiferids (<i>Operculina</i>)								
									Some recrystallisation								
									<i>N. howchini</i> , <i>Au. howchini</i> <i>F. bontangensis</i>								300
									<i>M. thecideaformis</i> Rare planktics and larger foraminiferids (<i>Eulepidina</i> sp. , <i>Cyc. eidae</i> <i>N. sumatrensis</i>)								400
																	500
									<i>Oper. victoriensis</i> Restricted foram fauna								600
									Barren of spores/pollen								
																	700
																	800

AGE (MA)	Magnetic Polarity Anomaly	ERA	Period	System	EPOCH	(1) STAGE	(2)	(3) International Foraminiferal zonations	East India letter class (4)	(5) Calcareous nannofossils	Palynoflora (6)		AGE (MA)
											Dinoflagellates	Spore-pollen	
1	2	2A	3	3A	3A	3A	N23	<i>Globigerinella calida</i>	Qh	QN15	NN1		
2	2A	3	3A	3A	3A	3A	N22	<i>Globorotalia truncatulinoides</i>	Qh	QN15	NN19		
3	2A	3	3A	3A	3A	3A	N21	<i>Globorotalia tosaensis</i>	Qh	QN15	NN16		
4	2A	3	3A	3A	3A	3A	N19/20	<i>Globorotalia miocenica</i>	Qh	QN15	NN12		
5	2A	3	3A	3A	3A	3A	N18	<i>Globorotalia margaritae</i>	Qh	QN15	NN11		
6	2A	3	3A	3A	3A	3A	N17	<i>Neoglobobulimina duterlei</i>	Qh	QN15	NN11		
7	2A	3	3A	3A	3A	3A	N16	<i>Neoglobobulimina doostensis</i>	Qh	QN15	NN9		
8	2A	3	3A	3A	3A	3A	N15	<i>G. menardi</i>	Qh	QN15	NN8		
9	2A	3	3A	3A	3A	3A	N14	<i>G. mayeri</i>	Qh	QN15	NN7		
10	2A	3	3A	3A	3A	3A	N13	<i>G. toshi robusta</i>	Qh	QN15	NN6		
11	2A	3	3A	3A	3A	3A	N12	<i>G. toshi robusta</i>	Qh	QN15	NN5		
12	2A	3	3A	3A	3A	3A	N11	<i>G. toshi robusta</i>	Qh	QN15	NN4		
13	2A	3	3A	3A	3A	3A	N10	<i>G. toshi robusta</i>	Qh	QN15	NN3		
14	2A	3	3A	3A	3A	3A	N9	<i>Præobulimina glomerata</i>	Qh	QN15	NN2		
15	2A	3	3A	3A	3A	3A	N8	<i>Globigerinella insueta</i>	Qh	QN15	NN1		
16	2A	3	3A	3A	3A	3A	N7	<i>Catapsydrax stainforthi</i>	Qh	QN15	NN1		
17	2A	3	3A	3A	3A	3A	N6	<i>Catapsydrax dissimilis</i>	Qh	QN15	NN1		
18	2A	3	3A	3A	3A	3A	N5	<i>Globorotalia kugleri</i>	Qh	QN15	NN1		
19	2A	3	3A	3A	3A	3A	N4	<i>Globorotalia kugleri</i>	Qh	QN15	NN1		
20	2A	3	3A	3A	3A	3A	P22	<i>Globigerina ciperoensis</i>	Qh	QN15	NN1		
21	2A	3	3A	3A	3A	3A	P21	<i>Globorotalia opima</i>	Qh	QN15	NN1		
22	2A	3	3A	3A	3A	3A	P19/20	<i>Globigerina amplapertura</i>	Qh	QN15	NN1		
23	2A	3	3A	3A	3A	3A	P18	<i>Cassigerinella chipolensis</i>	Qh	QN15	NN1		
24	2A	3	3A	3A	3A	3A	P17	<i>Pseudohastigerina micra</i>	Qh	QN15	NN1		
25	2A	3	3A	3A	3A	3A	P16	<i>Globorotalia cerroazulensis</i>	Qh	QN15	NN1		
26	2A	3	3A	3A	3A	3A	P15	<i>Particulasphera seminivoluta</i>	Qh	QN15	NN1		
27	2A	3	3A	3A	3A	3A	P14	<i>Truncatoloides rohri</i>	Qh	QN15	NN1		
28	2A	3	3A	3A	3A	3A	P13	<i>Orbulinoides beckmanni</i>	Qh	QN15	NN1		
29	2A	3	3A	3A	3A	3A	P12	<i>Morozovella lehneri</i>	Qh	QN15	NN1		
30	2A	3	3A	3A	3A	3A	P11	<i>Globigeropsis subcicloglobata</i>	Qh	QN15	NN1		
31	2A	3	3A	3A	3A	3A	P10	<i>Hankenerina aragonensis</i>	Qh	QN15	NN1		
32	2A	3	3A	3A	3A	3A	P9	<i>Acarina pentacamerata</i>	Qh	QN15	NN1		
33	2A	3	3A	3A	3A	3A	P8	<i>Morozovella aragonensis</i>	Qh	QN15	NN1		
34	2A	3	3A	3A	3A	3A	P7	<i>M. subbotina</i>	Qh	QN15	NN1		
35	2A	3	3A	3A	3A	3A	P6	<i>M. subbotina</i>	Qh	QN15	NN1		
36	2A	3	3A	3A	3A	3A	P5	<i>M. subbotina</i>	Qh	QN15	NN1		
37	2A	3	3A	3A	3A	3A	P4	<i>M. subbotina</i>	Qh	QN15	NN1		
38	2A	3	3A	3A	3A	3A	P3	<i>M. subbotina</i>	Qh	QN15	NN1		
39	2A	3	3A	3A	3A	3A	P2	<i>M. subbotina</i>	Qh	QN15	NN1		
40	2A	3	3A	3A	3A	3A	P1	<i>M. subbotina</i>	Qh	QN15	NN1		
41	2A	3	3A	3A	3A	3A	P0	<i>M. subbotina</i>	Qh	QN15	NN1		
42	2A	3	3A	3A	3A	3A	P-1	<i>M. subbotina</i>	Qh	QN15	NN1		
43	2A	3	3A	3A	3A	3A	P-2	<i>M. subbotina</i>	Qh	QN15	NN1		
44	2A	3	3A	3A	3A	3A	P-3	<i>M. subbotina</i>	Qh	QN15	NN1		
45	2A	3	3A	3A	3A	3A	P-4	<i>M. subbotina</i>	Qh	QN15	NN1		
46	2A	3	3A	3A	3A	3A	P-5	<i>M. subbotina</i>	Qh	QN15	NN1		
47	2A	3	3A	3A	3A	3A	P-6	<i>M. subbotina</i>	Qh	QN15	NN1		
48	2A	3	3A	3A	3A	3A	P-7	<i>M. subbotina</i>	Qh	QN15	NN1		
49	2A	3	3A	3A	3A	3A	P-8	<i>M. subbotina</i>	Qh	QN15	NN1		
50	2A	3	3A	3A	3A	3A	P-9	<i>M. subbotina</i>	Qh	QN15	NN1		
51	2A	3	3A	3A	3A	3A	P-10	<i>M. subbotina</i>	Qh	QN15	NN1		
52	2A	3	3A	3A	3A	3A	P-11	<i>M. subbotina</i>	Qh	QN15	NN1		
53	2A	3	3A	3A	3A	3A	P-12	<i>M. subbotina</i>	Qh	QN15	NN1		
54	2A	3	3A	3A	3A	3A	P-13	<i>M. subbotina</i>	Qh	QN15	NN1		
55	2A	3	3A	3A	3A	3A	P-14	<i>M. subbotina</i>	Qh	QN15	NN1		
56	2A	3	3A	3A	3A	3A	P-15	<i>M. subbotina</i>	Qh	QN15	NN1		
57	2A	3	3A	3A	3A	3A	P-16	<i>M. subbotina</i>	Qh	QN15	NN1		
58	2A	3	3A	3A	3A	3A	P-17	<i>M. subbotina</i>	Qh	QN15	NN1		
59	2A	3	3A	3A	3A	3A	P-18	<i>M. subbotina</i>	Qh	QN15	NN1		
60	2A	3	3A	3A	3A	3A	P-19	<i>M. subbotina</i>	Qh	QN15	NN1		
61	2A	3	3A	3A	3A	3A	P-20	<i>M. subbotina</i>	Qh	QN15	NN1		
62	2A	3	3A	3A	3A	3A	P-21	<i>M. subbotina</i>	Qh	QN15	NN1		
63	2A	3	3A	3A	3A	3A	P-22	<i>M. subbotina</i>	Qh	QN15	NN1		
64	2A	3	3A	3A	3A	3A	P-23	<i>M. subbotina</i>	Qh	QN15	NN1		
65	2A	3	3A	3A	3A	3A	P-24	<i>M. subbotina</i>	Qh	QN15	NN1		
66	2A	3	3A	3A	3A	3A	P-25	<i>M. subbotina</i>	Qh	QN15	NN1		
67	2A	3	3A	3A	3A	3A	P-26	<i>M. subbotina</i>	Qh	QN15	NN1		
68	2A	3	3A	3A	3A	3A	P-27	<i>M. subbotina</i>	Qh	QN15	NN1		
69	2A	3	3A	3A	3A	3A	P-28	<i>M. subbotina</i>	Qh	QN15	NN1		
70	2A	3	3A	3A	3A	3A	P-29	<i>M. subbotina</i>	Qh	QN15	NN1		
71	2A	3	3A	3A	3A	3A	P-30	<i>M. subbotina</i>	Qh	QN15	NN1		
72	2A	3	3A	3A	3A	3A	P-31	<i>M. subbotina</i>	Qh	QN15	NN1		
73	2A	3	3A	3A	3A	3A	P-32	<i>M. subbotina</i>	Qh	QN15	NN1		
74	2A	3	3A	3A	3A	3A	P-33	<i>M. subbotina</i>	Qh	QN15	NN1		
75	2A	3	3A	3A	3A	3A	P-34	<i>M. subbotina</i>	Qh	QN15	NN1		
76	2A	3	3A	3A	3A	3A	P-35	<i>M. subbotina</i>	Qh	QN15	NN1		
77	2A	3	3A	3A	3A	3A	P-36	<i>M. subbotina</i>	Qh	QN15	NN1		
78	2A	3	3A	3A	3A	3A	P-37	<i>M. subbotina</i>	Qh	QN15	NN1		
79	2A	3	3A	3A	3A	3A	P-38	<i>M. subbotina</i>	Qh	QN15	NN1		
80	2A	3	3A	3A	3A	3A	P-39	<i>M. subbotina</i>	Qh	QN15	NN1		
81	2A	3	3A	3A	3A	3A	P-40	<i>M. subbotina</i>	Qh	QN15	NN1		
82	2A	3	3A	3A	3A	3A	P-41	<i>M. subbotina</i>	Qh	QN15	NN1		
83	2A	3	3A	3A	3A	3A	P-42	<i>M. subbotina</i>	Qh	QN15	NN1		
84	2A	3	3A	3A	3A	3A	P-43	<i>M. subbotina</i>	Qh	QN15	NN1		
85	2A	3	3A	3A	3A	3A	P-44	<i>M. subbotina</i>	Qh	QN15	NN1		
86	2A	3	3A	3A	3A	3A	P-45	<i>M. subbotina</i>	Qh	QN15	NN1		
87	2A	3	3A	3A	3A	3A	P-46	<i>M. subbotina</i>	Qh	QN15	NN1		
88	2A	3	3A	3A	3A	3A	P-47	<i>M. subbotina</i>	Qh	QN15	NN1		
89	2A	3	3A	3A	3A	3A	P-48	<i>M. subbotina</i>	Qh	QN15	NN1		
90	2A	3	3A	3A	3A	3A	P-49	<i>M. subbotina</i>	Qh	QN15	NN1		
91	2A	3	3A	3A	3A	3A	P-50	<i>M. subbotina</i>	Qh	QN15	NN1		
92	2A	3	3A	3A	3A	3A	P-51	<i>M. subbotina</i>	Qh	QN15	NN1		
93	2A	3	3A	3A	3A	3A	P-52	<i>M. subbotina</i>	Qh	QN15	NN1		
94	2A	3	3A	3A	3A	3A	P-53	<i>M. subbotina</i>	Qh	QN15	NN1		
95	2A	3	3A	3A	3A	3A	P-54	<i>M. subbotina</i>	Qh	QN15	NN1		
96	2A	3	3A	3A	3A	3A	P-55	<i>M. subbotina</i>	Qh	QN15	NN1		
97	2A	3	3A	3A	3A	3A	P-56	<i>M. subbotina</i>	Qh	QN15	NN1		
98	2A	3	3A	3A	3A	3A	P-57	<i>M. subbotina</i>	Qh	QN15	NN1		
99	2A	3	3A	3A	3A	3A	P-58	<i>M. subbotina</i>	Qh	QN15	NN1		
100	2A	3	3A	3A	3A	3A	P-59	<i>M. subbotina</i>	Qh	QN15	NN1		
101	2A	3	3A	3A	3A	3A	P-60	<i>M. subbotina</i>	Qh	QN15	NN1		
102	2A	3	3A	3A	3A	3A	P-61	<i>M. subbotina</i>	Qh	QN15	NN1		
103	2A	3	3A	3A	3A	3A	P-62	<i>M. subbotina</i>	Qh	QN15	NN1		
104	2A	3	3A	3A	3A	3A	P-63	<i>M. subbotina</i>	Qh	QN15	NN1		
105	2A	3	3A	3A	3A	3A	P-64	<i>M. subbotina</i>	Qh	QN15	NN1		
106	2A	3	3A	3A	3A	3A	P-65	<i>M. subbotina</i>	Qh	QN15	NN1		
107	2A	3	3A	3A	3A	3A	P-66	<i>M. subbotina</i>	Qh	QN15	NN1		
108	2A	3	3A	3A	3A	3A	P-67	<i>M. subbotina</i>	Qh	QN15	NN1		
109	2A	3	3A	3A	3A	3A	P-68	<i>M. subbotina</i>	Qh	QN15	NN1		
110	2A	3	3A	3A	3A	3A	P-69	<i>M. subbotina</i>	Qh	QN15	NN1		
111	2A	3	3A	3A	3A	3A	P-70	<i>M. subbotina</i>	Qh	QN15	NN1		
112	2A	3	3A	3A	3A	3A	P-71	<i>M. subbotina</i>	Qh	QN15	NN1		
113	2A	3	3A	3A	3A	3A	P-72	<i>M. subbotina</i>	Qh	QN15	NN1		
114	2A	3	3A	3A	3A	3A	P-73	<i>M. subbotina</i>	Qh	QN15	NN1		
115	2A	3	3A	3A	3A	3A	P-74	<i>M. subbotina</i>	Qh	QN15	NN1		
116	2A	3	3A	3A	3A	3A	P-75	<i>M. subbotina</i>	Qh	QN15	NN1		
117	2A	3	3A	3A	3A	3A	P-76	<i>M. subbotina</i>	Qh	QN15	NN1		
118	2A	3	3A	3A	3A	3A	P-77	<i>M. subbotina</i>	Qh	QN15	NN1		
119	2A	3	3A	3A	3A	3A	P-78	<i>M. subbotina</i>	Qh	QN15	NN1		

Latitude: 24° 42'00"S
Longitude: 153° 15'30"E

Date drilled: Oct 1980
Company: Humber Barrier Reef Oils

[illegible]

AGE (MA)	Magnetic polarity	Period	System	EPOCH	(1) STAGE	International Foraminiferal zonations	Calcareous nannofossils	Palynoflora (6)	AGE (MA)
						(2)	(3)	(4)	
0		Q	NEOGENE	PLIOCENE	CALABRIAN	N23	<i>Globigerina calida</i>	ON15	0
2		2A			PIACENZIAN	N22	<i>Globorotalia truncatulinoides</i>	ON14	2
3		3A			ZANCLIAN	N19/20	<i>Globorotalia trossensis</i>	ON12	3
4		4A			MESSINIAN	N17	<i>Globorotalia margaritae</i>	ON10	4
5		5A			TORTONIAN	N16	<i>Neoglobobulimina dutreuxi</i>	ON9	5
10		10A					<i>Neoglobobulimina acostaensis</i>	ON7	10
15		15A			SERRAVALLIAN	N12	<i>G. menardi</i>	ON6	15
20		20A					<i>G. mayeri</i>	ON5	20
25		25A					<i>G. foshi robusta</i>	ON4	25
30		30A					<i>G. foshi foshi</i>	ON3	30
35		35A			LANGHIAN	N8	<i>Proorbulina glomerata</i>	ON2	35
40		40A			BURDIGALIAN	N6	<i>Globigerinella insueta</i>	ON1	40
45		45A					<i>Catapsydrax stainforthi</i>	ON1	45
50		50A					<i>Catapsydrax dissimilis</i>	ON1	50
55		55A					<i>Globorotalia kugleri</i>	ON1	55
60		60A					<i>Globigerina ciperoensis</i>	ON1	60
65		65A					<i>Globorotalia opima</i>	ON1	65
70		70A					<i>Globigerina ampliapertura</i>	ON1	70
75		75A					<i>Cassigerinella chipolensis</i>	ON1	75
80		80A					<i>Pseudohastigerina micra</i>	ON1	80
85		85A					<i>Globorotalia cerroazulensis</i>	ON1	85
90		90A					<i>Paricardosphaera seminivolata</i>	ON1	90
95		95A					<i>Truncorotaloides rohrri</i>	ON1	95
100		100A					<i>Orbulinoides beckmanni</i>	ON1	100
105		105A					<i>Morozovella lehneri</i>	ON1	105
110		110A					<i>Globigerinella subglobulata</i>	ON1	110
115		115A					<i>Hantkenina aragonensis</i>	ON1	115
120		120A					<i>Acarinina pentacamerata</i>	ON1	120
125		125A					<i>Morozovella aragonensis</i>	ON1	125
130		130A					<i>Morozovella formosa</i>	ON1	130
135		135A					<i>M. subbotinae</i>	ON1	135
140		140A					<i>Morozovella edgari</i>	ON1	140
145		145A					<i>Morozovella velascoensis</i>	ON1	145
150		150A					<i>Planorbulina pseudomenardii</i>	ON1	150
155		155A					<i>Planorbulina pusilla</i>	ON1	155
160		160A					<i>Morozovella uncinata</i>	ON1	160
165		165A					<i>Subbotina trinidadensis</i>	ON1	165
170		170A					<i>Subbotina pseudoduloides</i>	ON1	170
175		175A					<i>Globigerina eugubina</i>	ON1	175
180		180A					<i>Abathomphalus mayaroensis</i>	ON1	180
185		185A					<i>Gansserina gansseri</i>	ON1	185
190		190A					<i>Globotruncana aegyptiaca</i>	ON1	190
195		195A					<i>Globotruncana havanensis</i>	ON1	195
200		200A					<i>Globotruncana calcarata</i>	ON1	200
205		205A					<i>Globotruncana ventricosa</i>	ON1	205
210		210A					<i>Globotruncana elevata</i>	ON1	210
215		215A					<i>Dicarinella asymetrica</i>	ON1	215
220		220A					<i>Dicarinella concavata</i>	ON1	220
225		225A					<i>Dicarinella primitiva</i>	ON1	225
230		230A					<i>Marginalinella sigali</i>	ON1	230
235		235A					<i>Helvetoglobotruncana helvetica</i>	ON1	235
240		240A					<i>Whiteinella archaeocretacea</i>	ON1	240
245		245A					<i>Rotalipora cushmani</i>	ON1	245
250		250A					<i>Rotalipora reicheli</i>	ON1	250
255		255A					<i>Rotalipora brotzeni</i>	ON1	255
260		260A					<i>Rotalipora appenninica</i>	ON1	260
265		265A					<i>Rotalipora ticinensis</i>	ON1	265
270		270A					<i>Rotalipora subticinensis</i>	ON1	270
275		275A					<i>Brithenella breggiensis</i>	ON1	275
280		280A					<i>Ticinella primula</i>	ON1	280
285		285A					<i>Ticinella bejaouensis</i>	ON1	285
290		290A					<i>Hedbergella gorbachikae</i>	ON1	290
295		295A					<i>Globigerinelloides algeriana</i>	ON1	295
300		300A					<i>Schackoina cabri</i>	ON1	300
305		305A					<i>Globigerinelloides blowi</i>	ON1	305
310		310A					<i>Hedbergella sigali</i>	ON1	310
315		315A					<i>Globigerina hoterivica</i>	ON1	315
320		320A					minute planktic foraminiferids	ON1	320
325		325A					<i>Capionellids</i>	ON1	325
330		330A					<i>Capionellites</i>	ON1	330
335		335A					<i>Capionellopsis</i>	ON1	335
340		340A					<i>Capionella elliptica</i>	ON1	340
345		345A						ON1	345
350		350A						ON1	350
355		355A						ON1	355
360		360A						ON1	360
365		365A						ON1	365
370		370A						ON1	370
375		375A						ON1	375
380		380A						ON1	380
385		385A						ON1	385
390		390A						ON1	390
395		395A						ON1	395
400		400A						ON1	400
405		405A						ON1	405
410		410A						ON1	410
415		415A						ON1	415
420		420A						ON1	420
425		425A						ON1	425
430		430A						ON1	430
435		435A						ON1	435
440		440A						ON1	440
445		445A						ON1	445
450		450A						ON1	450

1 Berggren et al., (1985a, b), Burger, (in prep. a)
2 Blow (1969, 1979), Berggren (1969)
3 Bolli (1957, 1966), Stainforth et al., (1975),
Bolli & Premoli-Silva (1973), Caron (1985)
4 Adams (1984), Adams et al., (1986), Chaproniere
(1981), (1983), Jenkins et al., (1985)
5 Bukry (1973), 1975, Martini (1971), Roth (1978), Sissingh (1977)
6 Burger (1973), 1988, in prep. a), Dettmann (1986),
Dettmann & Playford (1969), Helby & Morgan (1987),
Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976),
Stover & Evans (1973), Stover & Partridge (1973)

Latitude: 15° 56.19'S Date drilled: Nov 1971-Jan 1972
Longitude: 152° 11.27'E Company: Deep Sea Drilling Project

Chronostratigraphy	Age/Depth Correlation	Depth (m) Drill	Sub-sea	Formation	Two-way time	Lithology
		0	1428	Seafloor		
		11		Unit 1		
		23				
		27				
		41				
		45				
		50				
		54				
		100		Unit 1		
		110				
		133				
		151				
		157			0.42	
		200		Unit 2		
		243				
		250				
		282				
		291				
		300				
		309				
		318		Unit 3		
		338				
		344				
		350				
		350				
		400				
		450				

Compiled from:
Burns, Andrews, et al. (1973)
Jenkins & Srinivasan (1985),
Kennett (1973),
Kennett & von de Borch (1985)

Inner neritic 0-20m
Mid neritic 20-100m
Outer neritic 100-200m
Upper bathyal 200-500m
Lower bathyal 500-2000m


Record 1990/7

Depositional environment							Well: DSDP Site 209							PLATE 10	
							Geochemical data							Depth (m)	
Locustrine	Estuarine	Inner	Neritic Mid	Neritic Outer	Neritic Upper	Neritic Lower	Remarks	TOC	S1	S2	TMAX	P1	H1	GP	
							base <i>Gr. truncatulinoides</i>	0.0							0
							Temp 4.84±0.05°C	0.0							
							base <i>Gr. tosaensis</i>								
							base <i>Gds. conglobatus</i> , <i>Gr. crassaformis</i>	0.0							50
							Temp 7.1±0.2°C								
							Estimated velocity is 1500 m/sec								
							base <i>Gr. kugleri</i>								100
								0.1							
							base <i>Ga. anguliseturalis</i>	0.0							150
							Estimated velocity is 2500 m/sec								200
							<i>Cycl. reticulatus</i> <i>Ret. bisecta</i>	0.1							250
								0.2							
															300
							Estimated velocity is 4000 m/sec	0.3							
								0.4							
								0.4							350
															400
															450

TOC: Total Organic Carbon (%)
S1: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)
P1: Production index (S1-S2)
H1: Hydrogen index (mg hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock) (S1+S2)
20/00/41


1 Berggren et al., (1985a, b), Burger, (in prep.)
2 Blow (1969, 1979), Breggren (1969)
3 Boli (1957, 1966), Stainforth et al., (1975),
Boli & Premoli-Silva (1973), Caron (1985)
4 Adams (1984), Adams et al., (1986), Chaproniere
(1981), (1983), Jenkins et al., (1985)
5 Bukry (1973, 1975), Martini (1971), Roth (1978), Sissingh (1977)
6 Burger (1973), 1988, in prep. a), Dettmann (1986),
Dettmann & Playford (1969), Helby & Morgan (1987),
Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976),
Stover & Evans (1973), Stover & Partridge (1973)



TOC: Total Organic Carbon(%)
 S1: Pyrolysis free-hydrocarbon signal
 (mg S2 hydrocarbons/g rock)
 S2: Pyrolysis kerogen signal
 (mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)
 PI: Production index (S1-S2)
 HI: Hydrogen index (mg hydrocarbons/g organic carbon)
 GP: Genetic potential (Kg hydrocarbons/ton rock)(S1+S2)

20/OQ/42



1 Berggren et al., (1985a, b). Burger, (in prep. a)

2 Blow (1969, 1979). Berggren (1969)

3 Bolli (1957, 1966). Stainforth et al., (1975)

4 Bolli & Premoli-Silva (1973). Caron, (1985)

5 Adams (1984). Adams et al., (1986). Chaproniere (1981), (1983). Jenkins et al., (1985)

6 Bukry (1973, 1975). Martini (1971). Roth (1978). Sissingh (1977)

7 Burger (1973). 1988, in prep. a). Dettmann (1986).

8 Dettmann & Playford (1969). Helby & Morgan (1987).

9 Helby et al., (1987). Ingram & Morgan (1988). Partridge (1976).

10 Stover & Evans (1973). Stover & Partridge (1973)

Compiled from: Andrews, Packham, et al., (1975)

Inner neritic 0-20m

Mid neritic 20-100m

Outer neritic 100-200m

Upper bathyal 200-500m

Lower bathyal 500-2000m

Abyssal 2000m +

Record 1990/77

TOC: Total Organic Carbon (%)

Si: Pyrolysis free-hydrocarbon signal (mg S2 hydrocarbons/g rock)

S2: Pyrolysis kerogen signal (mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)

P1: Production index (S1-S2)

H1: Hydrogen index (ma hydrocarbons/g organic carbon)

GP: Genetic potential (Kg hydrocarbons/ton rock) (S1+S2)

20/OQ/43

AGE (MA)	Magnetic polarity	ERA	PERIOD	System	EPOCH	(1) STAGE	International Foraminiferal zonations	Calcareous nannofossils	Palynoflora (6)	AGE (MA)
(2)	(3)	(4)	(5)	Dinoflagellates	Spore-pollen					
0										
1										
2										
3										
4										
5										
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
16										
17										
18										
19										
20										
21										
22										
23										
24										
25										
26										
27										
28										
29										
30										
31										
32										
33										
34										
35										
36										
37										
38										
39										
40										
41										
42										
43										
44										
45										
46										
47										
48										
49										
50										
51										
52										
53										
54										
55										
56										
57										
58										
59										
60										
61										
62										
63										
64										
65										
66										
67										
68										
69										
70										
71										
72										
73										
74										
75										
76										
77										
78										
79										
80										
81										
82										
83										
84										
85										
86										
87										
88										
89										
90										
91										
92										
93										
94										
95										
96										
97										
98										
99										
100										
101										
102										
103										
104										
105										
106										
107										
108										
109										
110										
111										
112										
113										
114										
115										
116										
117										
118										
119										
120										
121										
122										
123										
124										
125										
126										
127										
128										
129										
130										
131										
132										
133										
134										
135										
136										
137										
138										
139										
140										
141										
142										
143										
144										
145										
146										
147										
148										
149										
150										

- 1 Berggren et al., (1985a, b), Burger, (in prep.a)
2 Blow (1969, 1979), Berggren (1969)
3 Bolli (1957, 1966), Stainforth et al., (1975),
Bolli & Premoli-Silva (1973), Caron (1985)
4 Adams (1984), Adams et al., (1986), Chaproniere
(1981), (1983), Jenkins et al., (1985)
5 Bukry (1973), 1975, Martini (1971), Roth (1978), Sissingh (1977)
6 Burger (1973), 1988, in prep. a), Dettmann (1986),
Dettmann & Playford (1969), Helby & Morgan (1987),
Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976),
Stover & Evans (1973), Stover & Partridge (1973)

Latitude: 26° 06.61'S Date drilled: Nov 1971-Jan 1972
Longitude: 16° 13.27'E Company: Deep Sea Drilling Project

Chronostratigraphy	Age/Depth Correlation	Depth (m) Drill	Sub-sea	Formation	Two-way time	Lithology
			1545 m	Seafloor		
		0				
		37				
		42				
		85				
		100		Unit 1		
		200				
		203				
		310				
		375				
		400		Unit 1		
		432				
		488				
		500		Unit 2		
		539				
		577		Unit 2		
		594		Unit 2		
		600				
		700				
		TD				

Compiled from:
Burns, Andrews et al. (1973)
Kennett (1973)
Record 1990/7

Inner neritic 0-20m
Mid neritic 20-100m

Outer neritic 100-200m
Upper bathyal 200-500m
Lower bathyal 500-2000m

Depositional environment	Geochemical data	Depth (m)
Locustine	TOC	
Estuarine	S1	
Inner	S2	
Mid	TMAX	
Neritic	P1	
Outer	H1	
Upper bathyal	GP	
Lower bathyal		
Remarks		
		0
Gr. truncatulinoides	0.0	
Gr. tosaensis		
		100
Gr. tumida	0.0	
Velocities of 1500 to 1590 m/sec		
		200
Nq. acostaensis		
		300
Change in benthic foraminiferal fauna	0.0	
Orbulina		
Gds. triloba	0.0	
		400
Velocity of 1990 m/sec		
		500
Velocity of 2210 m/sec	0.1	
		600
Planktic % gradually increases	0.1	
Cr. tenuis, Z. sigmoides		
Velocity of 2230 to 2420 m/sec		
Ark. cymbiformis		
N. frequens		
		700








TOC: Total Organic Carbon (%)
St: Pyrolysis free-hydrocarbon signal
(mg S2 hydrocarbons/g rock)
S2: Pyrolysis kerogen signal
(mg S2 hydrocarbons/g rock)

TMAX: Temperature at which S2 signal is max (°C)
P1: Production index (S1-S2)
H1: Hydrogen index (mg hydrocarbons/g organic carbon)
GP: Genetic potential (Kg hydrocarbons/ton rock) (S1-S2)
20/OQ/44


1 Berggren et al., (1985a, b), Burger, (in prep.a)	5 Bukry (1973), 1975), Martini (1971), Roth (1978), Sissingh (1977)	Compiled from:	Inner neritic 0-20m	Upper bathyal 200-1000m	TOC:Total Organic Carbon(%)	TMAX:Temperature at which S2 signal is max (°C)
2 Blow (1969, 1979), Berggren (1969)	6 Burger (1973), 1988, in prep. a), Dettmann (1986).	Kennett, von der Borch (1985)	Mid neritic 20-100m	Lower bathyal 1000-2000m	SI:Pyrolysis free-hydrocarbon signal	PI:Production index (SI-S2)
3 Bolli (1957, 1966), Stainforth et al., (1975), Bolli & Premoli-Silva (1973), Caron (1985)	Dettmann & Playford (1969), Helby & Morgan (1987), Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976), Stover & Evans (1973), Stover & Partridge (1973)	Boersma (1985)	Outer neritic 100-200m		S1:P2 hydrocarbons/g rock	HI:Hydrogen index (ma hydrocarbons/g organic carbon)
4 Adams (1984), Adams et al., (1986), Chaproniere (1981), (1983), Jenkins et al., (1985)		<i>Record 1990/7</i>			S2:Pyrolysis krogen signal (mg S2 hydrocarbons/g rock)	GP:Genetic potential (Kg hydrocarbons/ton rock)(SI+S2)
						20/OQ/45

Latitude: 26° 06.7'S		Date drilled: Dec 1982				
Longitude: 161° 13.6'E		Company: Deep Sea Drilling Project				
Chronostratigraphy	Age/ Depth Correlation	Depth (m) Drill Sub-sea		Formation	Two-way time	Lithology
		0	1533 m	Seafloor		
		6				
		32		I _A		Oxidised foram/ nannofossil ooze
		50				
		100		I _B		Light coloured foram bearing nannofossil ooze
		105				
		200				
		214				
		245				
		300		I _C		Foram/ nannofossil chalk
		308				
		344				
		400				
		411				
		469		II		Siliceous foram bearing nannofossil chalk and chert
		488 TD				
		500				
		600		II		




1 Berggren et al., (1985a, b), Burger, (in prep.a)	5 Bukry (1973), 1975), Martini (1971), Roth (1978), Sissingh (1977)	Compiled from:	Inner neritic 0-20m	Outer neritic 10-200m	TOC:Total Organic Carbon(%)	TMAX:Temperature at which S2 signal is max (°C)
2 Blow (1969, 1979), Breggren (1969)	6 Burger (1973), 1988, in prep. a), Dettmann (1986)	Kennett,von der Borch (1985)	Mid neritic 20-100m	Bathal 200-2000	S1:Pyrolysis free-hydrocarbon signal	PI:Production index (S1-S2)
3 Bolli (1957, 1966), Stainforth et al., (1975), Bolli & Premali-Silva (1973), Caron (1985)	Dettmann & Playford (1969), Helby & Morgan (1987), Helby et al., (1987), Ingram & Morgan (1988), Partridge (1976), Stover & Evans (1973), Stover & Partridge (1973)	Boersma (1985) Record 1990/7		Abyssal 2000m+	(mg S2 hydrocarbons/g rock)	HI:Hydrogen index (ma hydrocarbons/g organic carbon)
4 Adams (1984), Adams et al., (1986), Chaproniere (1981), (1983), Jenkins et al., (1985)					S2:Pyrolysis kerogen signal (mg S2 hydrocarbons/g rock)	GP:Genetic potential (Kg hydrocarbons/ton rock)(S1+S2)
						20/OQ/46




HORIZON MATURATION PLOT : Anchor Cay-1




HORIZON TEMPERATURE PLOT : Anchor Cay-1



SUBSIDENCE HISTOGRAMS : Anchor Cay-1



PALEOHEATFLOW & TECTONIC SUBSIDENCE PLOT : Anchor Cay-1



COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : Anchor Cay-1

COMPOSITE GEOHISTORY PLOTS



Anchor Cay-1

VS = 1:23,810





GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES : —
ISO-VITRINITE REFLECTANCE : - - -

HORIZON MATURATION PLOT : Anchor Cay-1



HORIZON TEMPERATURE PLOT : Anchor Cay-1



COMPOSITE GEOHISTORY PLOTS

Anchor Cay-1



VS = 1:23,810

GEOHISTORY PLOT





DECOMPACTED BURIAL CURVES : —
ISO-VITRINITE REFLECTANCE : - - -



HORIZON MATURATION PLOT : Borabi-1



HORIZON TEMPERATURE PLOT : Borabi-1



COMPOSITE GEOHISTORY PLOTS


Borabi-1

VS = 1:19,048




GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES : ———
150-VITRINITE REFLECTANCE : - - - - -


HORIZON MATURATION PLOT : Kusa-1




HORIZON TEMPERATURE PLOT : Kusa-1



SUBSIDENCE HISTOGRAMS : Kusa-1



PALEOHEATFLOW & TECTONIC SUBSIDENCE PLOT : Kusa-1



COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : Kusa-1

COMPOSITE GEOHISTORY PLOTS


Kusa-1

VS = 1:23,810






GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES : —
ISO-VITRINITE REFLECTANCE : - - -

HORIZON MATURATION PLOT : Kusa-1



HORIZON TEMPERATURE PLOT : Kusa-1



COMPOSITE GEOHISTORY PLOTS


Kusa-1

VS = 1:23,810




GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES : ---
ISO-VITRINITE REFLECTANCE : ---


HORIZON MATURATION PLOT : Pasca-C1




HORIZON TEMPERATURE PLOT : Pasca-C1



SUBSIDENCE HISTOGRAMS : Pasca-C1



PALEOHEATFLOW & TECTONIC SUBSIDENCE PLOT : Pasca-C1



COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : Pasca-C1

COMPOSITE GEOHISTORY PLOTS

Pasca-C1


VS = 1:28,572

GEOHISTORY PLOT






DECOMPACTED BURIAL CURVES : ————

ISO-VITRINITE REFLECTANCE : - - - - -

HORIZON MATURATION PLOT : Uramu-1A



HORIZON TEMPERATURE PLOT : Uramu-1A



COMPOSITE GEOHISTORY PLOTS



Uramu-1A

VS = 1:23,810





GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES : —
ISO-VITRINITE REFLECTANCE : - - -

HORIZON MATURATION PLOT : AQUARIUS-1



HORIZON TEMPERATURE PLOT : AQUARIUS-1



COMPOSITE GEOHISTORY PLOTS







AQUARIUS-1







VS = 1:19,048

GEOHISTORY PLOT


DECOMPACTED BURIAL CURVES : —
ISO-VITRINITE REFLECTANCE : - - -

COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : AQUARIUS-1










HORIZON MATURATION PLOT : Wreck Island-1



HORIZON TEMPERATURE PLOT : Wreck Island-1










COMPOSITE GEOHISTORY PLOTS

Wreck Island-1


VS = 1:19,048

GEOHISTORY PLOT




DECOMPACTED BURIAL CURVES : ————
ISO-VITRINITE REFLECTANCE : - - - - -




HORIZON MATURATION PLOT : DSDP 210




HORIZON TEMPERATURE PLOT : DSDP 210



SUBSIDENCE HISTOGRAMS : DSDP 210



PALEOHEATFLOW & TECTONIC SUBSIDENCE PLOT : DSDP 210



COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : DSDP 210

COMPOSITE GEOHISTORY PLOTS

DSDP 210



VS = 1:33,333

GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES : —
ISO-VITRINITE REFLECTANCE : - - -





HORIZON MATURATION PLOT : DSDP 287





SUBSIDENCE HISTOGRAMS : DSDP 287

HORIZON TEMPERATURE PLOT : DSDP 287



PALEOHEATFLOW & TECTONIC SUBSIDENCE PLOT : DSDP 287

PRESENT DAY HEATFLOW : 1.4 h.f.u



COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : DSDP 287

COMPOSITE GEOHISTORY PLOTS


DSDP 287

VS = 1:28,572




GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES :
ISO-VITRINITE REFLECTANCE :


HORIZON MATURATION PLOT : DSDP 208




HORIZON TEMPERATURE PLOT : DSDP 208



SUBSIDENCE HISTOGRAMS : DSDP 208



PALEOHEATFLOW & TECTONIC SUBSIDENCE PLOT : DSDP 208



COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : DSDP 208

COMPOSITE GEOHISTORY PLOTS



DSDP 208

VS = 1:19,048





GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES :
ISO-VITRINITE REFLECTANCE :

HORIZON MATURATION PLOT : DSDP 587



HORIZON TEMPERATURE PLOT : DSDP 587



COMPOSITE GEOHISTORY PLOTS

DSDP 587


VS = 1:19,048

GEOHISTORY PLOT






DECOMPACTED BURIAL CURVES : ———
ISO-VITRINITE REFLECTANCE : - - - - -

COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : DSDP 587

HORIZON MATURATION PLOT : DSDP 588



HORIZON TEMPERATURE PLOT : DSDP 588



COMPOSITE GEOHISTORY PLOTS

DSDP 588

VS = 1:19,048

GEOHISTORY PLOT

DECOMPACTED BURIAL CURVES :
ISO-VITRINITE REFLECTANCE :

COMPUTED vs OBSERVED VITRINITE REFLECTANCE PLOT : DSDP 588