



**Australian Government**  
**Geoscience Australia**




# A Review of Surrogates for Marine Benthic Biodiversity

*McArthur, M. A., Brooke, B., Przeslawski, R., Ryan, D.A., Lucieer, V. L., Nichol, S., McCallum, A.W., Mellin, C., Cresswell, I.D., Radke, L.C.*

**Record**

**2009/42**

**GeoCat #  
69733**



# A review of abiotic surrogates for marine benthic biodiversity

GEOSCIENCE AUSTRALIA  
RECORD 2009/42

by

McArthur, M. A. <sup>1</sup>, Brooke, B. <sup>1</sup>, Przeslawski, R. <sup>1</sup>, Ryan, D.A. <sup>1</sup>, Lucieer, V. L. <sup>2</sup>, Nichol, S. <sup>1</sup>,  
McCallum, A.W. <sup>3</sup>, Mellin, C. <sup>4</sup>, Cresswell, I.D. <sup>5</sup>, Radke, L.C. <sup>1</sup>



**Australian Government**  
**Geoscience Australia**



- 
1. Marine and Coastal Environment Group, Geoscience Australia, GPO Box 378, Canberra ACT 2601
  2. Tasmanian Fisheries and Aquaculture Institute, University of Tasmania, Private Bag 49, Hobart, Tas. 7053, Australia.
  3. Marine Invertebrates, Melbourne Museum, GPO Box 666, Melbourne, Vic. 3001, Australia.
  4. School of Earth and Environmental Sciences, The University of Adelaide, Adelaide SA 5005, Australia
  5. Science Director, Wealth from Oceans National Research Flagship, CSIRO, GPO Box 1538, Hobart, Tas. 7001

**Department of Resources, Energy and Tourism**

Minister for Resources and Energy: The Hon. Martin Ferguson, AM MP

Secretary: Mr John Pierce

**Geoscience Australia**

Chief Executive Officer: Dr Neil Williams PSM

© Commonwealth of Australia, 2009

This work is copyright. Apart from any fair dealings for the purpose of study, research, criticism, or review, as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without written permission. Copyright is the responsibility of the Chief Executive Officer, Geoscience Australia. Requests and enquiries should be directed to the **Chief Executive Officer, Geoscience Australia, GPO Box 378 Canberra ACT 2601.**

Geoscience Australia has tried to make the information in this product as accurate as possible. However, it does not guarantee that the information is totally accurate or complete. Therefore, you should not solely rely on this information when making a commercial decision.

**ISBN Hardcopy 978-1-921672-43-9**

**ISBN Web 978-1-921672-44-6**

**GeoCat # 69733**

<p><b>Bibliographic reference:</b> McArthur, M.A., Brooke, B., Przeslawski, R., Ryan, D.A., Lucieer, V. L., Nichol, S., McCallum, A.W., Mellin, C., Cresswell, I.D., Radke, L. C. 2009. A review of abiotic surrogates for marine benthic biodiversity. Geoscience Australia, Record 2009/42. Geoscience Australia, Canberra. 61pp.</p>
---

# Contents

<b>CONTENTS</b> .....	<b>IV</b>
<b>FIGURE LIST</b> .....	<b>V</b>
<b>TABLE LIST</b> .....	<b>V</b>
<b>ACKNOWLEDGEMENTS</b> .....	<b>VI</b>
<b>EXECUTIVE SUMMARY</b> .....	<b>VII</b>
<b>1. INTRODUCTION</b> .....	<b>8</b>
<b>2. MARINE BIODIVERSITY</b> .....	<b>9</b>
2.1. BENTHIC DIVERSITY AT DIFFERENT SCALES .....	9
2.2. BIOTIC DATA .....	10
2.3. DIVERSITY INDICES .....	11
<b>3. PATTERNS OF BENTHIC DIVERSITY AND THEIR DRIVERS</b> .....	<b>12</b>
3.1. SPATIAL VARIABLES.....	14
3.1.1. <i>Latitude</i> .....	14
3.1.2. <i>Longitude</i> .....	15
3.1.3. <i>Depth</i> .....	16
3.2. SEABED SUBSTRATE PARAMETERS .....	16
3.2.1. <i>General Properties of Substrate Types</i> .....	16
3.2.2. <i>Grain Size and Size Variation</i> .....	19
3.2.3. <i>Gravel</i> .....	20
3.2.4. <i>Sand and Mud</i> .....	21
3.2.5. <i>Geophysical Properties and Sediment Fabric</i> .....	23
3.3. HABITAT COMPLEXITY.....	23
3.3.1. <i>Geomorphology</i> .....	23
3.3.2. <i>Spatial Heterogeneity</i> .....	26
3.3.3. <i>Habitat Size</i> .....	27
3.4. DISTURBANCE .....	27
3.5. PRODUCTIVITY .....	28
3.5.1. <i>Primary Productivity</i> .....	28
3.5.2. <i>Organic Carbon</i> .....	29
3.6. OCEANOGRAPHY .....	30
<b>4. UTILITY CONSIDERATIONS</b> .....	<b>32</b>
4.1. TEMPORAL VARIATIONS .....	32
4.2. SURROGACY IN THE DEEP SEA .....	33
4.3. SURROGACY ACROSS SPATIAL SCALES .....	33
4.4. COMBINED EFFECTS OF POTENTIAL SURROGATES ON BENTHIC BIODIVERSITY .....	34
4.5. UTILITY OF DIFFERENT SAMPLING REGIMES.....	35
4.6. MANAGEMENT DECISIONS AT VARIOUS SCALES.....	35
<b>5. SYNTHESIS</b> .....	<b>38</b>
<b>6. REFERENCES</b> .....	<b>40</b>

## Figure List

<b>Figure 3.1.</b> Conceptual map of the drivers of biodiversity in marine systems and some potential surrogates. ....	13
--	----

## Table List

<b>Table 3.1.</b> Key results of surrogacy studies in which species assembly was analysed against abiotic variables. ....	14
<b>Table 3.2.</b> Soft sediment variables considered as potential abiotic surrogates of benthic biodiversity. ....	18
<b>Table 3.3.</b> Faunal size classification and qualitative ‘soft sediment’ relationships ....	19
<b>Table 3.4.</b> Examples of soft-sediment features of various scales that may have surrogacy value. ....	23
<b>Table 4.1.</b> Approaches to classifying regions in the high seas. ....	37

## Acknowledgements

This work has been partly funded through the Commonwealth Environment Research Facilities (CERF) programme, an Australian Government initiative supporting world class, public good research. The work is part of the Surrogates Program in the CERF Marine Biodiversity Hub, a collaborative partnership between the University of Tasmania, CSIRO Wealth from Oceans Flagship, Geoscience Australia, Australian Institute of Marine Science and Museum Victoria.

## Executive Summary

A growing need to sustainably manage marine biodiversity at local, regional and global scales cannot be met by applying the limited existing biological data. Abiotic surrogates of biodiversity are thus increasingly valuable in filling the gaps in our knowledge of biodiversity patterns, especially identification of hotspots, habitats needed by endangered or commercially valuable species and systems or processes important to the sustained provision of ecosystem services. This review examines the use of abiotic variables as surrogates for patterns in benthic assemblages with particular regard to how variables are tied to processes affecting biodiversity and how easily those variables can be measured at scales relevant to resource management decisions.

Currents and wave action affect larval supply, food availability and disturbance regimes and can therefore be used to describe biodiversity patterns over local, regional and global scales. Direct gradient variables such as salinity, oxygen concentration and temperature can be strong predictive variables for larger systems, although local stability of water quality parameters may prevent usefulness of these factors at fine spatial scales.

Biological productivity has complex relationships with benthic biodiversity, and although the development of local and regional models cannot accurately predict outside the range of their biological sampling, remote sensing may provide necessary information. Indeed, interpolated values are available for much of the world's seas, and these are continually being refined by the collection of remote sensing and field data.

Sediment variables are often interconnected with complex relationships to biodiversity. The strength of the relationship between any one sediment variable and biodiversity may depend on the state of another sediment variable in that system. Percentage mud, percentage gravel, rugosity and compaction hold the strongest independent predictive power. Rugosity and the difference between gravel and finer sediments can be established using acoustic methods, but to quantify grain-size and measure compaction, a sample is necessary.

Pure spatial variables such as latitude, longitude and depth are not direct drivers of biodiversity patterns but often correspond with driving gradients that they can be of some use in prediction. In such cases it would be better to identify what the spatial variable is acting as a proxy for so boundaries for that variable are not overlooked. The utility of these potential surrogates will vary across spatial scales, quality of data, and management needs, and the continued focus on surrogate research will address the need of marine scientists and resource managers worldwide for accurate and robust predictions of species distribution patterns.

# 1. Introduction

Marine biodiversity is of direct benefit to society as a food source, potential pharmacopoeia (Hunt & Vincent 2006), stabilizer of inshore environments (Jie et al. 2001) and regulator of atmospheric processes (Murphy & Duffus 1996). Marine biodiversity provides indirect benefits to society through ecological stability (Menge et al. 1999) and benthic-pelagic coupling (Ponder et al. 2002) which contribute to self-sustaining marine ecosystems. Marine biodiversity also has recreational, aesthetic and intrinsic value (Ponder et al. 2002; Wilson 1994).

Describing marine biodiversity can be challenging, particularly for habitats with difficult-to-sample biota. Using the relationships between abiotic factors and biota to predict patterns of biodiversity offers a promising alternative to biological sampling. The idea that physical and chemical properties of a system can act as surrogates of biodiversity is intuitive, but it is only recently that the relationships between organisms and their environment have been studied in sufficient detail and tested with appropriate mathematical tools to allow estimates of biodiversity to be made based on environmental data alone (Meynard & Quinn 2007). Such surrogacy relationships can be based on physical or chemical properties of a habitat (abiotic surrogacy) or on a defined taxonomic group (biotic surrogacy).

The use of surrogates to assess diversity is particularly important if a natural resource management decision must be made based on minimal biological information. In the application of surrogacy to management questions, consideration must be given to the degree to which present-day benthic environments have been and will be impacted by human activities occurring over the last century. Benthic ecosystems have been negatively impacted by over-fishing (Jackson 2008), bottom trawling and dredging (Pauly et al. 2005), pollution of coastal waters (Halpern et al. 2008), aquaculture and introduced species (Galil 2000; Jackson 2008; Pauly et al. 2005) and human-induced climate change (Bindoff et al. 2007). The combination of these direct and indirect human impacts on the marine environment is inducing unprecedented changes in marine ecosystems, and further loss of biodiversity is likely (Jackson 2008). Logistical constraints may preclude biological sampling in some impacted or vulnerable areas, and abiotic surrogacy research may provide a viable means to estimate biodiversity for marine conservation and vulnerability assessment.

The aim of this review is to synthesize available information on benthic biodiversity and its measurement and to explore the function and utility of potential abiotic surrogates. The review is structured in three parts assessing the current status of surrogacy research and its importance to marine resource management. First, marine biodiversity is defined and its measurement discussed. Second, the roles of abiotic factors in shaping biodiversity are examined. Third, considerations relevant to surrogate utility are addressed.

One challenge facing marine scientists and resource managers is to make sufficiently accurate and robust predictions of biodiversity using the more extensive abiotic datasets currently available so those predictions can be applied to resource management strategies. Therefore, we also discuss the importance of surrogacy research to marine resource management.

## 2. Marine Biodiversity

Several levels of complexity can be examined when considering marine diversity. Genetic diversity within a species or population and diversity of functional roles within a community are informative variables to quantify (Gray 1997), but species is the unit of diversity most easily conceptualized and is therefore most commonly considered (Willig et al. 2003). Of the many definitions available, the most widely applied species model is that of a genetically distinct, reproductively isolated population (Mayr 1996). The number of species in a given space is often referred to as the species diversity but ecologists refer to this integer value as species “richness,” denoted as  $S$ , reserving “diversity” for equation-derived values accounting for proportionality of species’ populations (Magurran 2004).

Marine species range in size and complexity from viruses to whales (Snelgrove 2001), and a range of species compete for resources at each level of size and complexity. Bouchet (2006) reported 229 602 described marine species, but estimates of the grand total of marine taxa based on extrapolations from existing data vary widely: Grassle and Maciolek (1992) proposed that ten million species exist in the benthos alone, and May (1992), using conservative assumptions, responded with an estimate of 500 000 benthic species. The range of these estimates indicate much work must be done to reliably estimate the existing richness of the world’s seafloor and even then, describing the species would only be the first step in defining species’ distributions, information critical to marine spatial planning.

### 2.1. Benthic Diversity at Different Scales

Because of the different sampling strategies required to account for diversity at different scales, ecologists divide animals of interest into non-taxonomic size groups: megafauna are large enough to identify by eye; macrofauna describes organisms small enough to require microscope attention for accurate identification and extends down to those retained on a 300  $\mu\text{m}$  sieve; meiofauna pass through 300  $\mu\text{m}$  mesh but are retained on a 44  $\mu\text{m}$  sieve; and microbes (bacteria and protists) can pass through a 44  $\mu\text{m}$  sieve. Microbes may represent the largest component of Earth’s genetic diversity and biomass (Snelgrove 2001).

Each of these ecologically arbitrary size divisions generates a following among ecologists who tend to sample and model their component of interest as an independent unit, with communities therefore rarely treated holistically (Yeom & Adams 2007). As a result of this division of effort and thought, diversity measures for a given system may only take into account the community fraction of interest to the reporting specialist. This approach may ignore important processes at work on other scales (Ponder et al. 2002) but is often necessary to reduce the costs of sampling and processing.

New species of squid and whales are still being discovered (Chivian & Bernstein 2008) but, compared to megafauna, little is known about distributions and biology of most macrofaunal and meiofaunal species due to the difficulty of sampling and sorting material at these scales and the shortage of expert taxonomists (Bouchet 2006). Macrofauna and meiofauna are crucial components of the benthic ecosystem, playing key roles in nutrient cycling, benthic-pelagic coupling, bioturbation, and succession (Rex et al. 2006), and the response of these groups to environmental variables can be different to

that of the megafauna (Edgar 1999). Research on these groups tends to focus on the intertidal (Hourston et al. 2005) or very shallow systems (Dye 2005).

Finding a value for the species richness of an entire system is currently impossible as the shortage of taxonomic expertise and the large number of undescribed taxa preclude accurate identification of all organisms at all scales in all but the most intimately studied systems (Costello et al. 2006).

## 2.2. Biotic Data

There are five options for organism identification available to researchers undertaking surrogacy research: 1) Identify all organisms to the species level, 2) Identify subsets of taxa for which expertise is available to species level, 3) Identify species to a coarse taxonomic level, 4) Identify target species, or 5) Identify functional groups. Biological surrogacy, where the presence of one taxon implies the presence of others is not addressed here. As species is the level at which organisms react to their habitat (Bertrand et al. 2006), the first option is the most comprehensive and will have the greatest power to detect relationships with abiotic variables. Unfortunately, identification to species level is often difficult, time-consuming and costly (Ponder et al. 2002). Use of coarse taxonomic resolution may hinder the quantification of biodiversity and the utility of potential surrogates (Bertrand et al. 2006) and spatial variation becomes less distinct as taxonomic resolution is lowered (Anderson et al. 2005). Several studies have shown that identification at the genus-, family-, and even order-level is sufficient to detect community response to environmental gradients (Wlodarska-Kowalczyk & Kedra 2007). Nevertheless, even at genus level there may be sufficient biological disparity between closely related organisms to dictate contrasting distributions that limit the utility of surrogates based on taxonomic generalizations (Pitcher et al. 2007). The effectiveness of target species analysis depends on the group chosen (Wlodarska-Kowalczyk & Kedra 2007), and the utility of target groups may vary according to the age of a community (Magierowski & Johnson 2006).

The final approach to quantifying diversity involves identifying organism roles within a system. Functional groups describe organisms that share a similar physiological or ecological function e.g.: deposit feeders, bioturbators, predators (Bonsdorff & Pearson 1999). When clearly defined, such groupings can be used as a proxy for diversity and may provide a more practical way to assess potential surrogacy relationships than strict taxonomic approaches because biological function is more closely associated with abiotic variables than taxonomy. The validity of using functional groups has been questioned because analyses based on such divisions may be meaningless without more comprehensive knowledge about life history and biology of marine biota than is currently available for most species (Pearson 2001). In addition, some evidence points to species identity being closely linked to ecosystem services such as bioturbation (Norling et al. 2007). Potential surrogacy relationships may best be examined by incorporating multiple functions related to the environmental variable of interest, particularly for soft sediment communities (Pearson 2001). The choice of functional group divisions will affect interpretation of biodiversity patterns, as well as the magnitude and significance of relationships between abiotic variables and biota (Sanders et al. 2007).

Further investigation into the effects of taxonomic resolution on the ability to detect patterns is needed as they likely vary across ecosystems and communities. The utility of functional groups to surrogacy has not been examined, and future research

should investigate the suitability of functional group perspectives in the development of predictive models of ecological and biological function based on abiotic factors (Sanders et al. 2007).

### 2.3. Diversity Indices

Species richness, also referred to as alpha diversity, increases monotonically with the area or volume sampled (He & Legendre 2002) and with sampling effort (Benkendorff & Davis 2002). Therefore, when sampling effort is even, comparison of the number of species (richness) should be a valid way to assess biological response to impacts or environmental differences between sites. Different species rarely occur in equal numbers at a site, and the proportion that each species contributes to the community can be as important in defining differences between systems as species richness may be more sensitive to environmental change (Gray 1997). Diversity indices seek to turn the number of species and the proportionality of their relative counts in a given area into a single number for ready comparison of samples, sites or ecosystems (Magurran 2004). This often transforms the relatively simple integers of species richness into complex representations of proportionality. Most proportionality-based diversity indices are geared to record a high number in assemblages where a large number of species occur in similar numbers (high evenness) and a low number if a small number of species dominate a sample or system (low evenness) (Magurran 2004).

Sampling cannot be expected to account for every species in every assemblage (Chao et al. 2005). Richness estimators calculate the number of species expected based on the sequence in which they are encountered in a series of samples, using random iterations of sample data through equations that account for richness and proportionality. Richness estimators are used to determine how many species might be expected in an assemblage under differing levels of sampling (Ugland et al. 2003).

The change in species richness along environmental gradients is referred to as beta diversity. Low beta values indicate assemblage equivalency, and high values indicate assemblage disparity (Magurran 2004). Simple measures of beta diversity compare alpha diversity against the species richness of a region (gamma diversity) (Magurran 2004). Others make direct comparisons of species' presence and absence (e.g. Jaccard's coefficient) or species' relative proportions (e.g. Bray-Curtis coefficient) between two systems (Clarke & Warwick 1998).

The values incorporated by diversity and evenness measures only take into account the number of species and their proportions. Patterns described by these indices may be of interest in an environmental impact assessment, but interpretation of the results back to ecological processes is difficult (Clarke & Warwick 2001). An increase in species richness or evenness can be expected immediately after environmental change when resources, previously fully utilized, are suddenly made available to recruiting species and pre-climax community structures (Connell 1978). Therefore, it is difficult to define ecological change at intermediate levels of disturbance or to examine biological responses to natural environmental gradients using diversity information alone (Clarke & Gorley 2006).

Diversity indices can act as tools to identify community change in response to environmental change, or habitat gradients and richness estimates derived from iterative processes can be used to explore assemblage differences graphically through rarefaction curves. Diversity indices may suggest new avenues of investigation, but the index most

appropriate to a particular situation must be chosen and the resulting values considered carefully with regard for the environmental drivers of a system and how that index will account for them (Magurran 2004). As the values generated by many diversity and evenness indices are complex, mathematical representations of integer data usually cannot be used in hypothesis testing (Legendre & Legendre 1998).

### 3. Patterns of Benthic Diversity and Their Drivers

The goal of surrogacy research is to determine which easily measured characteristics best describe the species assemblage of a particular space and time (Moore et al. 1991). These characteristics are then expected to act as predictors of species assemblages in unexplored areas (Franklin 1995). Pitcher (2007) identified grain size, carbonate composition, available space, benthic irradiance, sheer stress, bathymetry, bottom water physical properties, nutrient concentrations and turbidity as abiotic surrogates of biotic distributions on the Great Barrier Reef; but these variables, while useful predictors, may not be the forces driving the patterns they describe.


The influence of abiotic factors on species assemblages is due to the effect they exert on fundamental niches. A species' fundamental niche was defined by Austin *et al.* (1990) as "that hypervolume defined by environmental dimensions within which a species can survive and reproduce." Fundamental niches are rarely fully realized by species because interspecific competition, disease and disturbance events displace individuals and populations, resulting in a reduced occupied hypervolume, often referred to as the realised niche (Austin & Smith 1989). The environmental gradients that describe a species' fundamental niche can be broadly grouped into resource gradients – e.g. chemicals or energy consumed by a species; direct gradients – variables with a physiological influence on a species but not consumed by it – e.g. sediment grain size or temperature; and indirect gradients – variables correlated with direct and resource gradients but with no physiological connection to the species – e.g. depth and latitude (Meynard & Quinn 2007). When niche theory was first proposed, species were expected to exhibit a Gaussian distribution to environmental gradients but skewed distributions are more common in ecological studies as the effects of additional variables express their influence (Karadzic et al. 2003).

Predictions of biodiversity can also be made using neutral theory, an idea with similarities to island biogeography theory (Volkov et al. 2003), in which relative species abundances are assumed to be determined by random immigration to a system (Leigh Jr 2007). Neutral theory is not universal (Leigh Jr 2007) and has been shown to conflict with some empirical data (Dornelas et al. 2006; McGill 2003), thus niche theory remains the most appropriate ecological model on which to base surrogacy research.

The abiotic variables historically ascribed the greatest direct influence over benthic organism distributions are temperature, salinity, oxygen concentration, light availability and sediment composition (Snelgrove 2001). A model of these influences and some of their potential synergies is given in [Figure 1](#), and a range of recent studies that have quantified the extent to which abiotic information accounted for biotic data are noted in [Table 3.1](#). Olabarria (2006) found depth accounted for as much as a quarter of the

variance in benthic diversity in deep systems but, as benthic organisms lack an apparent mechanism for measuring depth, some correlated water quality parameter or seafloor characteristic most likely influences the settlement, recruitment and survival processes that result in the observed depth related patterns. In the same sense, latitude can act as a proxy for a range of gradients of direct importance to benthic organisms regardless of their actual position relative to the equator. Indirect gradients can be location specific, giving them limited value in explaining realised niches (Austin & Smith 1989) but if the correlation between the indirect and the direct drivers are general enough, predictions based on the indirect variables are of value (Moore et al. 1991), particularly when they are easily measured.

Gray (2002) reduced Snelgrove’s list of direct drivers (2001) to productivity, temperature and sediment composition as the dominant variables in determining regional benthic richness, noting that temperature and productivity are often correlated to depth and latitude. Combinations of these driving influences occur with varying spatial and temporal consistency, in turn producing semi-regular patterns of biodiversity. The validity and origin of several identified general benthic biodiversity patterns are the focus of much recent debate. For example, the latitudinal richness gradient, widely accepted as a rule for benthic fauna since the mid twentieth century (Thorson 1957) has recently been shown to be weaker than previously thought (Snelgrove 2001) or entirely incorrect for some taxa or systems (Rex et al. 2005).



**Figure 3.1.** Conceptual map of the drivers of biodiversity in marine systems and some potential surrogates. Black lines indicate relationships between potential surrogates. Bold black lines represent relationships between processes. Gray lines show potential surrogates linked to a process. Dotted lines show potential relationships between surrogates and biological variables. Brown borders enclose abiotic factors. Green borders enclose biological or biophysical factors.

**Table 3.1.** Key results of surrogacy studies in which species assembly was analysed against abiotic variables.

Study	Important/best variables	% variability accounted for	Statistical process
(Post et al. 2006)	% mud, % gravel, disturbance, depth	59	BIOENV
(Passlow et al. 2006)	depth, longitude	25	BIOENV
(Sanders et al. 2007)	depth	65	BIOENV
(Beaman et al. 2005)	Slope, % gravel, %CaCO <sub>3</sub>	75	BIOENV
(Beaman & Harris 2007)	Slope, % gravel, turbidity	62	BIOENV
(Williams et al. 2006)	Depth, latitude, gear type, longitude	44	BIOENV
(Stevens & Connolly 2004)	% mud, distance to ocean	30	Spearman's
(Ellingsen et al. 2005)	Depth, depth	29	Linear regression
(Gogina et al. 2010)	Depth, total organic content	50, 43	BIOENV (also Spearman's, CCA)

### 3.1. Spatial Variables

Although pure spatial variables are not the driving force behind the patterns they can be used to describe, variables such as latitude, longitude and depth can act as strong proxies for factors that influence species richness such as temperature, day length, light penetration and a variety of dispersal variables (Hawkins 2001).

#### 3.1.1. Latitude

Increasing species richness with increasing proximity to the equator is a long recognised biogeographic pattern (Cox & Moore 2005). While this pattern has been documented in the marine environment (Attrill et al. 2001) it has recently been found to be less general than previously thought (Gray 2001). The terrestrial latitudinal richness gradient, first recorded by von Humboldt at the start of the nineteenth century (Willig et al. 2003) and noted by Darwin on the Beagle (Darwin 1839), led researchers to seek matching patterns in marine systems (Stehli et al. 1967). Particularly strong latitudinal signals are evident in the molluscan fauna on the coastal shelves of North America (Gage 1996) and among deep sea macrofauna (Piepenburg et al. 2002).

Hawkins (2003) and Willig (2003) each cite thirty hypotheses to explain latitudinal richness gradients. These were categorised by Mittelbach (2007) into ecological, historical and evolutionary groups. The ecological hypotheses concentrate on the different adaptive challenges faced by organisms living in different climatic zones: polar and temperate organisms must adapt to environmental conditions (Schemske 2002) while tropical organisms, dealing with less harsh abiotic extremes, adapt to biotic interactions (Crame 2000). Historical explanations concentrate on the age and stability of richness hotspots (Alongi 1990). There is evidence that modern benthic taxa have their origins in the ancient Tethys Sea, radiating into temperate and polar regions during the Miocene (Gage 2004). Evolutionary models incorporate several possible drivers for high rates of speciation including the wide variety of microhabitats available in tropical regions (Rex et al. 2005) and higher rates of molecular evolution (Kerswell 2006).

Attempts to measure and explain the extent of a latitudinal richness gradient in taxa other than molluscs on broad geographic scales have found less evidence for a marine equivalent to the terrestrial pattern (Gray 2001) and brought into question the treatment of data in describing such patterns. For example, Thorson's (1956) pattern of increasing richness in benthic epifauna toward the equator was based only on the 140,000 marine taxa known at the time (Snelgrove 2001).

In contrast to the northern hemisphere, southern hemisphere faunal richness does not show clear latitudinal gradients. Surveys of Antarctic fauna reveal a much higher richness among animals than the depauperate arctic (Brey et al. 1994). Some taxa actually become more speciose towards the poles, and the Latitudinal Gradient Project has been established to examine ecological patterns in a range of systems using latitude as a surrogate for climatic variables in the Ross Sea (Howard-Williams et al. 2006). The age and stability of Antarctic shelf benthos may be a potential contributor to local high species richness (Gray 2002). Box-core sampling of isopods on Australia's south-east slope (Poore et al. 1994) and fish trawls on the south-west slope (Williams & Bax 2001) provide estimates of species and family richness much higher than those of equivalent mid-latitude regions in the northern hemisphere.

Peak richness among bryozoans and some groups of crustaceans occur in the temperate southern latitudes (Barnard 1991; Barnes & Griffiths 2008; Bolton 1996; Kerswell 2006); marine macro-algae and their associated faunal communities, confined to shallow waters, exhibit a bimodal relationship with latitude. Algal richness hotspots occur in both northern and southern temperate regions (Fraser & Currie 1996). This could be due to depressed tropical algal richness caused by direct competition between algae and corals (Snelgrove 2001), but Kerswell (2006) sought a driving mechanism in oceanic currents which, due to their relative direction of rotation in the northern and southern hemispheres, may promote algal propagule dispersal from tropical to temperate regions.

Diversity responses to pure spatial gradients can vary depending on how assemblage data are treated. Ellingsen (2005) determined that latitude could account for variance in richness of molluscs (11.8 % explained), annelids (9.6 %) and crustaceans (13.7 %) in Norwegian shelf benthos, but this influence dropped (4.9 %) when assemblages were considered collectively.

### 3.1.2. Longitude

Although less dramatic than the latitudinal changes in species richness, longitudinal diversity gradients are also evident, and richness hotspots occur in the western regions of both the Pacific and the Atlantic (Jones 1950). The Australasian tropical systems are particularly species rich because of their proximity to the highly diverse Indo-West Pacific region. Coral diversity decreases as one moves eastwards or westwards away from the Indo-West Pacific (Veron 1995). Similar patterns have been found in the cool temperate oceans where bryozoans decrease in diversity away from an Australasian hotspot (Barnes & Griffiths 2008). While significant species turnover (high beta diversity) was measured across the southern Australian coastline, echinoderm and decapod species richness did not vary significantly with longitude (O'Hara & Poore 2000). Kerswell (2006) attributes the algal component of longitudinal richness patterns to propagule distribution caused by oceanic currents, in this case carrying material westward either side of the equator with no equivalent mechanism for eastward dispersal at that latitude.

At a local scale (1-2°) the effects of latitude or longitude on patterns of biodiversity will likely not be observable (Coleman et al. 2007). When observed at local scales, continental patterns are disrupted by other spatial variables (e.g. depth) and environmental ones (e.g. substrate), but general spatial patterns are useful as surrogates of benthic biodiversity at a global scale.

### 3.1.3. Depth

Depth has been a consistently powerful explanatory variable in benthic studies (Nicolaidou & Papadopoulou 1989, Gogina et al. 2010). When generalising from shallow to deep, intertidal and estuarine systems exhibit high biomass and low species richness caused by high productivity and extreme environmental conditions (Edgar 2001), coastal shelves have moderate biomass and species richness (Snelgrove 2001), and the deep sea shows a decrease in biomass and increase in richness (Levin et al. 2001). Peak benthic species richness values have been recorded seaward of the continental rise, excluding the deep sea (Snelgrove 2001). The lower slope and abyssal plains become comparatively depauperate for some groups, and species turnover tends to be high (Paterson et al. 1992). Levin et al. (2001) stated that the deep sea houses greater diversity than coastal shelf systems, although at far lower abundances. Areas with low abundance and high species turnover require greater sampling effort to reliably account for diversity (Etter & Mullineaux 2001).

High relief benthic habitats offer refuge from predators and settlement surfaces not available on flat bottoms. Seagrass beds, reefs, mangrove forests and algal assemblages are home to more species than the adjacent sandy or muddy substrate, accounting for increased richness at a local and regional scale. Seagrasses are poor as a source of food for grazing taxa (Fry 2006), but the vertical structure provided by plants offer habitat for edible epiphytes in addition to being a refuge from predation (Jenkins & Wheatley 1998). Biogenic structures such as worm tubes, mussel beds and hard coral skeletons offer similar habitat advantages (Callaway 2006; Nakamura & Sano 2005; Reise 1981; Tsuchiya & Nishihira 1986).

## 3.2. Seabed Substrate Parameters

### 3.2.1. General Properties of Substrate Types

Numerous studies have provided evidence to show significant differences in the species composition between 'hard' and 'soft' substrata (Beaman et al. 2005; Beaman & Harris 2007; Williams & Bax 2001). For practical purposes, 'soft substrate' is usually defined as detrital mineral or biogenic sediment comprising grains with a mean diameter less than 2 mm, although gravel size fractions are often included (Lewis & McConchie 1994). The term 'hard substrate' is typically used to represent rock outcrops but may include sediments with large grain size (e.g. cobbles, boulders) since these materials can provide a surface that is functionally comparable to bedrock. While the contrast between soft and hard substratum is conceptually simple, defining the boundary between soft and hard substrates can be complex in practice because some rock types are friable or semi-consolidated and may be partly covered by sediment (Ryan et al. 2007b). In addition, because the boundaries between adjacent soft-sediment environments are not always as sharp as those across hard and soft substrate features, the associated boundaries between

biological assemblages may be gradational and spatially complex (Beaman & Harris 2007).

Sediment particle size distribution and composition on the seabed express a strong influence on the morphology and life history of species living in soft sediments (Jones 1950). These variables are determined by complex interactions between local geology, rates of sediment production and supply, actions of bioeroders, current and wave induced bed stress, and slope (Reineck & Singh 1980). Generally, in high-energy areas, coarse sediments (gravel) will predominate, whereas lower energy (depositional) areas are muddy, although there are exceptions (Foster 2001; Hart & Kench 2007; Rees et al. 2007) which have led to a highly complex and variable distribution of seabed sediment types on the shelf and slope.

Although several sediment surrogacy relationships are well documented (Beaman & Harris 2007; Brown et al. 2001; Degraer et al. 2008), the nature and strength of sedimentary surrogates for species composition within soft sediment environments remains a subject of debate (Dye 2006; Inoue et al. 2008; Stevens & Connolly 2004). Soft substrates are home to epifauna and infauna and plant life may include seagrasses (and their epiphytes) or microphytic algae occurring at the sediment-water interface. Hard substrates can act as habitat for epifauna and encrusting or macro-algae, but infauna are excluded.

Many apparent relationships between sediment type and biota remain untested in an experimental sense (Whitlatch 1981) and have been challenged (Snelgrove & Butman 1994). [Table 3.2](#) collates information on the properties of soft-sediment benthic substrates that appear to influence distributions of benthic organisms. More detailed examinations of the relationships between these variables and the biota will follow.

## Abiotic Surrogates Review

**Table 3.2.** Soft sediment variables considered as potential abiotic surrogates of benthic biodiversity.

Parameter	Surrogacy potential	Mechanism	Ease of measurement	Source
Mean grain size	Poor in poorly sorted sediments	N/A – indirect gradient variable.	Needs physical sample. Processing becoming faster with automation.	(Degraer et al. 2008; Diaz et al. 2004)
% mud	Strong	Provides habitat or food for infauna.	As per mean grain size.	(Degraer et al. 2008; Diaz et al. 2004)
% sand	Poor on its own, tends to be inverse to mud	Provides habitat or food for infauna and foraging animals. High permeability offers opportunity for oxygenation of lower horizons.	As per mean grain size.	(Snelgrove & Butman 1994)
% gravel	Strong	Acts as a settlement surface and likely to have large interstices to act as habitat for cryptic species.	Can be differentiated from sand and mud by acoustic means.	(Kostylev et al. 2001)
Sorting	Not established but potentially valuable	Contributes to size of interstices.	As per mean grain size.	(Whitlach 1981)
Skewness	Noted but not quantified	Increased food availability with positive skewness.	As per mean grain size.	(Hogue & Miller 1981)
Compaction/ Porosity	Strong	Contributes to size of interstices.	Requires careful treatment of an undisturbed sample. Cannot be determined from stored material.	(Degraer et al. 2008; Edgar 2001)
Rugosity	Strong	Disturbance in unstable sediments (dunes, ripples, bedforms) depresses richness and biomass. These features may also affect settlement by changing circulation patterns.	Can be identified using acoustic means.	(Kostylev et al. 2001; Thouzeau et al. 1991)

### 3.2.2. Grain Size and Size Variation

Quantitative measures of sediment grain size (e.g. percentage mud, mean grain size) have often been linked to the distribution of infaunal and epifaunal benthic communities (Degraer et al. 2008; Whitlach 1981) (Table 3.3). Research conducted between the 1950s and 1990s concluded that seabed grain-size, measured from grab or dredge samples, exerts strong control on the distribution of benthic assemblages (Ellingsen & Gray 2002; Etter & Grassle 1992; Gray 1974; Rhoads 1974; Sanders 1958; Snelgrove & Butman 1994; Whitlach 1981). Clear influences have been indicated between sediment infauna (meio- and macrofauna) and the physical structure of sediment such as grain-size parameters and the level of sediment compaction (Degraer et al. 2008; Edgar 2001). The influence of grain size properties on epifaunal or demersal species distributions is clear in some environments (Greene et al. 2007a) but is likely to be one of several physical factors, such as seabed sediment mobility, deposit-feeding potential, turbidity and the potential for attachment, that are important driving factors (Snelgrove & Butman 1994).

**Table 3.3.** Faunal size classification and qualitative ‘soft sediment’ relationships after (Edgar 2001).

Classification	Faunal type	Minimum sediment size class	Example faunal groups
Megafauna (> 10 mm)	Epifauna	n/a	Flatfish, gobies, flatheads, dragonets, stingrays, shrimp, holothurians, heart urchins, brittle stars, various crabs.
Macrofauna (0.5-10 mm)	Infauna, epifauna	Very coarse sand	Amphipods, callianassid shrimps, bivalves, polychaetes
Meiofauna (0.063-0.5 mm)	Infauna	Fine sand	Nematodes, harpacticoid copepods, flatworms
Microfauna (< 0.063 mm)	Infauna	Silt, clay	Protozoa, bacteria, fungi

Several summary statistics are derived from standardised techniques that measure the range and abundance of sediment grain size in a sample, including mean grain size, standard deviation of the size range around the mean (sorting) and the symmetry or preferential spread (skewness) to one side of the mean (Blott & Pye 2001).

Mean grain size measurements alone do not take into account particle size variation (sediment sorting) or particle shape (e.g. disc or rod), and are therefore most useful for describing highly unimodal sediment grain size distributions (Lewis & McConchie 1994). Direct comparison of mean grain size may be misleading as a poorly sorted sediment of mud and gravel could produce the same value as a well sorted sand despite representing an entirely different habitat. The mean grain size value is the most commonly reported measure of sediment physical properties because it is relatively easy to measure and it is conceptually simple to represent a sample with one value that captures a basic seabed property, but the information lost in generating the mean may decrease its potential as a biodiversity surrogate (Snelgrove & Butman 1994).

Sorting, the standard deviation of grain size distribution, is a potentially valuable surrogate because it indicates the amount of interstitial space available within a sediment for benthic organisms to use as habitat (Gray 1974). For example, sedimentary environments that are poorly sorted, such as cobble and boulder fields, may incorporate several orders of magnitude difference in sediment grain size, forming complex three dimensional benthichabitat (Motta et al. 2003). Grain size range also determines the

amount of space between particles. In heterogenous sediments, interstices between large particles may be completely filled with smaller grains.

Despite the potential for sorting as a habitat indicator, Whitlatch (1981) could not establish any correlation between sediment sorting and species diversity in benthic deposit feeders in a study of animal-sediment patterns in an intertidal setting. Sorting remains a parameter that few studies have considered in the context of the potential for explaining infaunal or epifaunal distribution patterns.

The skewness of a grain size distribution is rarely reported in studies of soft-sediment benthic habitats. The utility of this parameter lies in it being a measure of the tail, or extremes, of the grain size distribution. A distribution with strong positive skewness (+1 to +3) has an excess of fine-grained particles relative to a normal distribution with the same mean and strong negative skewness (-3 to -1) reflects an excess of coarse grains (Folk 1974). In the case of a positively skewed distribution for sand-dominated sediment, the fine fraction may include sediment of organic origin that could represent an important food source for infauna. This was observed (but not quantified) by Hogue and Miller (1981) in the shallow subsurface of sand ripples where silt represents a potential food source for nematodes.

### 3.2.3. Gravel

Sediments dominated by size fractions greater than 2 mm are classed as 'gravel,' incorporating granules (2 - 4 mm), pebbles (4 - 64 mm), cobbles (64 - 256 mm) and boulders (> 256 mm) (Wentworth 1922). These deposits have the greatest degree of three dimensional structure, permeability and the largest interstitial spaces offering shelter from predators and strong currents. In the English Channel, Brown et al. (2001) found presence or absence of gravel to be the most important variable influencing community composition. On Georges Bank in the northwest Atlantic, Thouzeau (1991) found significant changes in species richness, total abundance and biomass with differing sediment type, with higher values being associated with gravel substrates. Kostylev *et al.* (2001) also recognised the importance of gravel lags on the Scotian Shelf and subsequently identified strong links to scallop distribution and stock abundance, readily defined by high acoustic backscatter (Kostylev et al. 2003). Hard substrate fauna may recruit to sparsely scattered cobbles or shell debris large and stable enough to be suitable for permanent colonisation (Auster & Langton 1999). Very coarse sediments are generally acoustically distinct from finer material (e.g. muds and sands) and are readily mapped by acoustic mechanisms (Kostylev et al. 2003).

#### 3.2.3.1 Boulders and Cobbles

Boulder fields are characterised by high densities of organisms due to their provision of refuge at a range of scales and possibly due to their effects on local hydrodynamics and grain size (Smith & Otway 1997). Boulders provide a complex microhabitat comprising upper surfaces (dominated by algae and providing demersal fish habitat), lower surfaces (dominated by sponges, bryozoans and ascidians), and soft sediment habitats beneath (Motta et al. 2003). The large interstices provided by semi-stationary boulders have been shown to be important sheltering and aggregation sites for numerous fish (Greene et al. 2007a) which use boulders to feed from and as a shelter from predators. Certain species also use the sheltered undersides of boulders to deposit egg masses (e.g. gastropods in

Przeslawski & Davis 2007), thereby making boulders important in reproduction and larval recruitment. Boulder and cobble aggregations also provide significant interstitial habitat for a range of crypto-benthic communities (Patzner 1999). Motta et al. (2003) showed that the sediments underneath boulders were affected by complex hydrodynamic conditions, and incorporate a high percentage of organic matter even in relatively coarse sediments, such as gravel, where levels of organic material are often thought to be low.

Cobble-dominated sediments are typically undisturbed by fair-weather wave-generated currents and most tidal currents. Consequently, these sediments are functionally similar to 'hard' substrate in that they are available for colonisation by attaching organisms (Brown et al. 2001) and also provide cryptic shelter; however, they may be periodically mobilised by storms (Lindholm et al. 2001).

### **3.2.3.2 Granules and Pebbles**

Granule to pebble-sized deposits comprise rock fragments or biogenic calcium carbonate and are often hydrodynamically mobilised. These finer-grained gravels provide important continental shelf habitats that influence the distribution of a range of benthic biota (Beaman et al. 2005). As with larger-grain material, granules and pebbles form a highly permeable substrate with interstitial cavities and 3D-microstructure that is large enough to host a range of infauna, depending on the degree of sediment sorting.

Some organisms (molluscs, corals, green and red calcareous algae) generate gravel-fraction material through the production of calcareous skeletons. In areas of high calcium fixation these organisms contribute to the formation of a gravel-dominated habitat such that a measure of carbonate gravel is a potentially valuable surrogate for some benthic communities. Granule to cobble-sized coral rubble and shingle are important habitats which influence larval settlement on and adjacent to coral reefs (Mumby & Harborne 1999). Similarly, rhodolith beds are recognised as habitat with high local infaunal and epifaunal biodiversity (Hinojosa-Arango & Riosmena-Rodriguez 2004) where faunal settlement is enhanced (Culliney 1974) and three-dimensional microhabitats exist for many surficial organisms (Foster 2001). While high gravel proportion can act as a useful abiotic surrogate (Auster & Langton 1999; Greene et al. 1995; Kostylev et al. 2001; Thouzeau et al. 1991), small gravel percentages in otherwise sandy or muddy samples appear to have little bearing on biodiversity patterns (Post et al. 2006).

### **3.2.4. Sand and Mud**

Sandy habitat is a widely recognised term used to describe expanses of relatively bare (devoid of obvious epifauna) sediment, mainly quartz or carbonate, comprising grains that range from 63  $\mu\text{m}$  to 2 mm, featuring variable sediment sorting properties and often characterised by ripple or dune bedforms (Ryan et al. 2007a). Sand dominated sediments are easily mobilised as bedload in moderate to high energy continental shelf settings, resulting in the development of a dynamic bedform dominated seafloor (Hemer 2006). Large epibenthic and demersal species tend to occur in low abundance or as ephemeral visitors, a feature attributable to the absence of shelter, high sediment mobility, and lack of a firm substrate for use by attaching organisms (Gray 1974). Sandy habitats are important for burrowing organisms and their predators and provide foraging areas for species which may also use nearby firm substrate areas for sheltering purposes (Ross et al. 2007). The bulk of macrofaunal biomass and richness is concentrated in the first few

centimeters of soft sediments (Snelgrove 2001). Below this, oxygen penetration is matched by oxygen consumption, and organisms must be able to use anaerobic respiration or maintain a link to the oxygenated water above to survive (Snelgrove & Butman 1994). Large, loosely packed particles offer greater oxygen penetration than fine, dense muds.

Mud comprises grains including silt (4  $\mu\text{m}$  – 63  $\mu\text{m}$ ) and clay (<4  $\mu\text{m}$ ) (Wentworth 1922). The term often has compositional connotations implying a soft often dark sediment which may contain detrital minerals and have a high organic content (Rhoads 1974). Silty muds resuspend easily and do not aggregate into bedforms such as ripples and dunes (Hemer 2006), although biogenic microstructures may be common (e.g. burrow holes, mounds, traces). Fine muds (clay) can be strongly cohesive due to electrostatic bonds between clay particles, which can make them more resistant to resuspension (Lewis & McConchie 1994).

The high particle surface area to volume ratios available in muddy sediments offer deposit feeders optimal feeding (Taghon 1988). Coarse sediments generated by the removal of fine material by currents offer similar advantages to suspension feeders as food bearing currents are more likely to support their survival (Wildish & Kristmanson 1997). Thus, a measure of percentage volume sand content can offer reciprocal surrogacy value to percentage mud, where gravel proportions are low. Functional associations can be blurred, however, where the grouping of organisms as suspension or deposit feeders is unclear. Whitlach (1981) also noted that intuitive ideas about recruitment to muddy and sandy sediments were confounded by high numbers of deposit and suspension feeding taxa co-occurring in some soft sediments.

Few studies identify percentage sand as a stronger influence on the distribution of benthic communities than percentage mud or gravel so it is likely that sand percentage alone is a relatively poor surrogate (i.e. the difference between mud or gravel and sand is less than the difference between unconsolidated and consolidated sediment). This is a point strongly argued by Snelgrove and Butman (1994) who recognised the importance of looking beyond grain size patterns to also consider sediment dynamics and hydrodynamics in determining organism distributions.

A range of benthic assemblages can occur on sandy or muddy substrates in a small area indicating the significant influence of other physical variables, especially hydrodynamics, in defining community structure (Wildish & Kristmanson 1997). Recent studies have found that the relationship between muddy substrates and benthic taxa appears varied and complex. In the English Channel a combination of factors including mud content and hydrodynamic forces correlated with species distributions in coarse and muddy sediments (Freeman & Rogers 2003). Similarly, in the Arctic, an association of muddy sediments and high total organic carbon content was recognised as an important environmental control on the distribution of macrofaunal communities, particularly in open-sea environments (Denisenko et al. 2003). In contrast, Post et al. (2006) found relatively weak links between mud content and the density of taxa (Spearman rank correlation 0.186) in the Gulf of Carpentaria, whereas the combination of water depth, percent gravel and seabed exposure was found to have stronger explanatory power (correlation 0.599).

### 3.2.5. Geophysical Properties and Sediment Fabric

The roughness of the sea floor is recognised as an important control on habitat type and biodiversity (Greene et al. 2007a). For soft-sediment substrates, seabed roughness can be caused by infaunal bioturbation at the local scale through to large bedforms such as sand waves (Table 3.4). Sedimentary bedforms such as ripples, dunes and sand waves are known to affect the abundance and distribution of benthic organisms by disturbing their habitat (Barros et al. 2004) or by influencing local circulation, larval dispersion and food supply (Vasslides & Able 2008). On Georges Bank, Thouzeau (1991) described a 10-fold decrease in megafauna density on sand dunes compared to more stable sandy bottoms. Kostylev (2001) recorded similar patterns between active bedforms and more stable areas for the Scotian Shelf. Combining parameters that describe bedform patterns and sediment grain size measurements represents a method of capturing information on both the physical and hydrodynamic characteristics of benthic habitats (Kostylev et al. 2001).

The sediment-organism relationships determining benthic assemblage distributions are often complex and involve geological, hydrodynamic, chemical and biological factors that may be difficult to measure. For many settings, sediment properties provide a useful first-order surrogate. The value of these parameters as surrogates can be significantly enhanced when they are employed in combination with other parameters, such as measures of the seabed hydrodynamic regime that better represent the complex ecological processes that control the distribution of benthic organisms.

**Table 3.4.** Examples of soft-sediment features of various scales that may have surrogacy value.

Feature	Linear scale (km)	Relief (m)	Sediment type	Formative process	Potential ecological impact	Example reference
Burrow mounds, holes & small bed irregularities	0.01-1	0.01-1	Mud, sand	Bioturbation; growth of sessile organisms	Local disruption of currents; alteration of surficial grain size; irrigation of sediments	Widdicombe <i>et al.</i> 2003
Sedimentary bedforms (ripples, dunes, sand waves)	0.1-1	0.1-10	Sand, gravel	Near bed current (wave, tidal or ocean current)	Unstable substrate	Thouzeau <i>et al.</i> 1991
Boulder and gravel beds	10-100	1-10	coarse gravel	Wave erosion; Sea level change	Local disruption of currents; potentially semi-consolidated	Green <i>et al.</i> 2007

## 3.3. Habitat Complexity

### 3.3.1. Geomorphology

Seabed geomorphology encompasses the shape and hardness of the seabed at a range of spatial scales and describes the processes behind these occurrences (Kennet 1982). At the broadest spatial scale, geomorphological features are categorical descriptors of the shape

of the seabed (e.g. reefs, ridges, seamounts) that describe structures that range in scale from thousands of km<sup>2</sup> (e.g. basins) to a few hundred m<sup>2</sup> (Harris et al. 2008). Seabed geomorphology can also be described by a range of parameters derived from various types of bathymetric, acoustic, video and seabed sample data. For example, seabed slope and roughness (rugosity) are parameters that can describe the structural complexity of the seabed over regional (100s of km) (Kukowski et al. 2008) to site-specific scales of a few metres (Holmes et al. 2008).

Rugose benthic habitats offer refuge from predators and settlement surfaces not available on flat bottoms. Seagrass beds, reefs, mangrove forests and algal assemblages are home to more species than the adjacent sandy or muddy substrate, accounting for increased richness at a local and regional scale. Seagrasses are poor as a source of food for grazing taxa (Fry 2006), but the vertical structure provided by plants offer habitat for edible epiphytes in addition to being a refuge from predation (Snelgrove 2001). Biogenic structures such as worm tubes, mussel beds and hard coral skeletons offer similar habitat advantages (Callaway 2006; Nakamura & Sano 2005; Reise 1981; Tsuchiya & Nishihira 1986).

Geomorphological parameters form useful surrogates where ecological processes are known or suspected to link seabed structure and the distribution of benthic communities. Seafloor topographic complexity is ecologically important because it provides habitat structure for juvenile and adult animals (Kostylev et al. 2003), plays a role in regulating foraging patterns (Erlandsson et al. 1999) and alters boundary-layer flow over the seabed (Green et al. 1998; Ke et al. 1994). The interaction of flow and substrate heterogeneity affects larval settlement and subsequent population performance because it controls delivery of food, oxygen, and chemical cues (Lenihan 1999). Seabed complexity also influences erosion, transport and deposition of sediment (Widdows et al. 1998) and macrofauna (Urbanski & Szymelfenig 2003). Importantly, as structural complexity shapes the perception and behaviour of benthic organisms it influences higher level processes of population dynamics and community structure (Kostylev et al. 2001). Areas of high species richness are often associated with habitat heterogeneity (Gladstone 2007) - the complexity of habitat can interrupt predator-prey relationships that in a simpler habitat might lead to clear dominance or removal of certain species. Thus, in complex habitats, species may co-exist in a greater diversity than they would in a simpler habitat (Peterson et al. 1998).

The spatial extent to which a geomorphological variable acts as a surrogate for a benthic assemblage or set of species is an important consideration, especially if there is no clearly defined process that explains the link between the physical parameter and distribution of biota (Holmes et al. 2008, Greene et al. 2007c). Links may not be direct, however, and geomorphic parameters may themselves be surrogates for other environmental factors directly controlling ecological processes: sand waves may be a surrogate for wave and current energy, and canyons structure may act as a surrogate for light penetration (Kostylev et al. 2001). Geomorphological surrogacy relationships can range in scale from biogeographic patterns that occur across ocean basins and seamounts (Etter & Mullineaux 2001) through to fine-scale patchiness in shelf benthic community types related to variations in habitat structure produced by the distribution of rocky and unconsolidated seabed (Ryan et al. 2007b). Most often, however, the performance of geomorphological parameters as surrogates is more effective when a combination of variables is considered or when they are considered in concert with sedimentological or oceanographic variables (Post et al. 2006, Greene et al. 2007c).

The delineation of seabed geomorphological features requires bathymetric data with a resolution that matches the scale of the features of interest. Typically, a seabed digital elevation model is derived from bathymetry data and classified using expert interpolation supported by various types of topographic feature algorithms (Lanier et al. 2007). For ocean-basin or global scale assessments of seabed geomorphology, broad-scale compilations of a wide range of data types have been used (Harris et al. 2008). For regional scale studies, a range of data types (hydrographic chart data, single-beam echosounder, multibeam) and resolutions are usually combined (Romsos et al. 2007). For sub-regional to local scales, multibeam sonar data are usually employed (Kostylev et al. 2001), while in shallow, clear-water settings, laser airborne depth sounder systems are often used (Walker et al. 2008). The advantage of multibeam and laser systems is that they can provide high resolution area coverage of accurate bathymetric data. Multibeam systems can also provide backscatter intensity data, which can indicate the texture and hardness of the seabed (Kloser et al. 2007). At regional to local scales, sidescan sonar images are very useful for identifying seabed textural patterns that are produced by both physical (sediment type, roughness) and biological features (macro algae, sessile fauna) (Prada et al. 2008). Local-scale geomorphic features may also be observed directly in towed underwater video footage (Anderson et al. 2007).

In terms of broad patterns in seabed biodiversity and geomorphology, recent studies of seamounts have shown several to be mid-ocean biological hotspots providing hard substrate for attaching organisms and creating a range of unique benthic environments in terms of energy regimes and rates of food supply through the interaction of the seamount topography and ocean currents (Clark et al. 2006). Similarly, structurally complex granite outcrops are characterised by benthic communities distinctly different to those living in the surrounding flat, unconsolidated seabed due to the formation of microhabitats by fractures and joints and enhanced food supply from the current flows around outcrops (Beaman et al. 2005).

Geomorphology often combines with other factors to be used as a successful surrogate. For example, the distribution of benthic communities in the southern Gulf of Carpentaria, Australia, was best explained by the influence on biota of different combinations of depth, bottom disturbance and seabed composition (Post et al. 2006). Similarly, in the northern Great Barrier Reef, seabed slope, in combination with the proportion of gravel and turbidity, is a major factor that influences the observed spatial distribution of megabenthos assemblages (Beaman & Harris 2007). In addition, Holmes (Holmes et al. 2008) used several geomorphological parameters to build predictive models of the distribution of a range of sessile communities in a marine park on the temperate inner and middle shelf of western Victoria, Australia. The best predictors of the main sessile communities in the park, as determined from towed video of the seabed, were found to be bathymetry (2 m grid), hypsometric integral (i.e., a non-dimensional measure of the proportion of an area that is above a certain elevation) (circular window radius 50m and 100m) and local relief (radius 100m) (Holmes et al. 2008). Similarly, quantitative morphological parameters (slope, rugosity, topographic position index) derived from multibeam sonar data collected on the temperate Oregon shelf discriminated a range of seabed structural features (ridge, boulders, cobbles) that form key refugia for several demersal fish species (Whitmire et al. 2007). Broad scale (1-10 km) benthic terrains on the deep-water areas of the continental slope of southeastern Australia, classified using 'soft', 'hard', 'smooth', and 'rough' parameters derived from multibeam sonar data, were shown to be good predictors of the distribution of benthic faunal groups

(Kloser et al. 2007). This approach represents an effective method of mapping seabed habitats and potential patterns of biodiversity in this vast continental slope environment (>150 m – 1000 m) at a scale useful for conservation management (Kloser et al. 2007). However, these parameters alone may not represent good predictors of the distribution of some faunal groups, and other substrate categories such as “mixed” may provide more useful information for use in management decisions (Greene et al. 2007a, b).

### 3.3.2. Spatial Heterogeneity

Spatial heterogeneity is often cited as a diversity driver, with high inshore species richness being promoted by the variety of habitats available on a broad scale and deep water benthic and planktonic richness occurring in spite of low biomass as a result of small scale shifts in sediment or water composition in habitats which otherwise appear homogenous (Snelgrove 2001). Rocky reef habitats are structured across a full spectrum of spatial scales, from sub-millimeter to the order of tens of meters, and these patterns influence the behaviour of benthic organisms, in turn influencing higher level processes of population dynamics and community structure (Knight & Morris 1996). Spatial pattern is difficult to quantify and refers to the spatial character and arrangement, position or orientation of patches within a landscape (Li & Reynolds 1993). Terrestrial landscape ecologists have long studied the effects of spatial pattern on process (With 1997) and have developed a large collection of metrics to describe landscape pattern (O'Neill et al. 1998). These metrics have proved useful for the description of landscape structure and its spatial dynamics over a broad range of spatial and temporal scales (Riitters et al. 1995). Landscape metrics applied to ecological studies fall into two general categories (Gustafson 1998): One quantifies the composition of a map with reference to spatial attributes, and the other category quantifies the spatial configuration of the map requiring spatial information for their calculation. There are many ways to define landscape depending on the phenomenon under consideration. The issue is that a seascape is not necessarily defined by its size but by an interacting mosaic of patches relevant to the habitat under consideration (at any scale) (Harris et al. 2008). Scale in remote sensing is typically defined by pixel resolution but objects have their own inherent scale, and the same type of objects can appear different at different scales (Lillesand & Kiefer 1994). For example, high frequency multibeam sonar data can be processed to < 1 m resolution implying that the minimum object that can be detected is twice the pixel size. In this case patterns in reef systems can likely be detected from individual reefs two meters and greater in size.

A greater understanding of the distribution and complexity of benthic habitats and a common approach to measuring and describing this complexity will provide a spatial framework within which to properly address spatially explicit research and management goals (Kendall et al. 2005). Conservation strategies now frequently consider not only amounts of habitat that must be retained but also the spatial configurations of habitats across depth and exposure gradients (Fortin et al. 2005). The decline of many species has been linked directly to habitat loss and fragmentation. Identifying what characteristics make an area preferentially habitable for particular species has been examined by many landscape ecologists and is being increasingly taken up by marine ecologists to describe patterns of benthic diversity (Barrett et al. 2001). Numerous indices of landscape pattern have been linked to ecological function by recent advances in image processing and GIS technologies. Studies indicate that abundance and species richness of reef fish are related

to habitat diversity (Ault & Johnson 1998), habitat extent (Tolimieri 1998), and habitat complexity (Jones & Syms 1998), but the value of such indices is yet to be incorporated into spatial analysis of marine ecological patterns. It may be true for the marine environment, as for terrestrial studies, that the distinction between what can be measured and patterns relevant to the ecological phenomenon under investigation is sometimes blurred (Levin 1992). Uncertainties in mapping the pattern and extent of marine habitats have, until recently, been behind the rarity of habitat-scale studies in marine ecology (Gray 2000).

### 3.3.3. Habitat Size

The species-area relationship represents one of the earliest quantitative models in biogeography, since a relationship between the number of species and land area was noted as early as 1778 (Brown & Gibson 1983). This was further developed into the equilibrium theory of island biogeography (MacArthur & Wilson 1963). This theory emphasises the importance of island area in combination with isolation, colonization rate and extinction rate as determinants of the equilibrium number of species. Extensions of this model treat species as functionally indistinguishable, only diverging through demographic stochasticity (Bell 2000). Thus, community composition is constantly changing over time as a dynamic balance between colonization and extinction (Kadmon & Allouche 2007).

Evidence for species-area relationships in the marine realm has been demonstrated for marine invertebrate communities associated with boulders (McGuinness 1984), rock walls (Smith & Witman 1999), mussel beds (Witman 1985), coral heads (Abele & Patton 1976) and for coral and reef fishes (Bellwood et al. 2005). Theories based on the species-area relationship have however been criticised for neglecting species-energy relationships (Kalmar & Currie 2006), small-island effect (Lomolino & Weiser 2001) and, above all, the fact that species differ functionally in how they interact with the environment in space and time (Gardner & Engelhardt 2008). The latter assumption forms the basis of the niche theory (Hutchinson 1957) which was recently coupled to the island biogeography theory into a unified general model (Kadmon & Allouche 2007). To successfully characterise a system based on abiotic variables, the area available to form particular habitats must be taken into consideration and the sampling scaled with this in mind (Thrush et al. 2005).

### 3.4. Disturbance

Dayton (1971) established space clearing disturbances as a key variable determining marine assembly structure. Storms and current eddies may contribute to primary space being made available in a system in two fashions: increasing sheer stress at the benthos/water interface, which can remove sediment, algal cover and motile fauna; and mechanical abrasion or damage caused by moving sediment or projectiles (Sousa 2001). Benthic space can also be made available in the wake of acute pollution events (Scanlan et al. 1993), fishing activity (Currie & Parry 1996) and eutrophication (Tett et al. 2007), but the effects tend to be locally focused.

The intermediate disturbance hypothesis of Connell (1978) predicts maximum biodiversity at a frequency of disturbance where recruitment is able to replace lost individuals but inter-specific processes don't have time to exclude species. This response has been recorded in many marine systems (Begon et al. 1996; Gutt & Piepenburg 2003;

Svensson et al. 2007). Disturbance, both anthropogenic and natural, may act as a potential abiotic surrogate for diversity at an appropriate spatial scale and temporal scale (Harris et al. 2008). For example, bed shear stress was found to be of use as an explanatory variable in fish occurrence prediction on the Great Barrier Reef (Pitcher et al. 2007). National indices of disturbance and exposure are currently being developed through Geoscience Australia's GEological and Oceanographic Model of Australia's Continental Shelf (GEOMACS), with general predictions that biodiversity will be highest in areas of intermediate disturbance (Connell 1978).

The stability of the seabed sediment surface exerts a major control on benthic community structure (Newell et al. 1998). Species diversity tends to be highest on stable rocky shores and on cohesive muddy shores, with the more mobile sandy or fine gravel substrates typically showing much lower richness. Sediment stability is dependent on slope, particle size and the degree of water motion on the bed (Bagnold 1963). The shape and roundness of sediment grains are additional properties that determine the stability of a deposit (Lewis & McConchie 1994) but grain shape is difficult to measure and is rarely recorded despite its likely importance. Stability may also be influenced by the presence of biota through biological armouring of the bed and binding of sediment by faunal mucus (Murray et al. 2002). The stability of a sediment surface as a habitat is difficult to quantify, particularly given that one of the key proxies, sediment grain size, is determined on disaggregated samples which have been dislodged from their environment and may have little physical resemblance to what an organism actually encounters (Snelgrove & Butman 1994). Laboratory experiments of near-bed flow and sediment stability in relation to settlement of benthic organisms (St-Onge & Miron 2007), and field verification of the results is required to test their utility as surrogates (Post et al. 2006).

### **3.5. Productivity**

#### **3.5.1. Primary Productivity**

Opposite to patterns observed for terrestrial systems (Currie et al. 2004), high primary productivity in near shore waters tends to promote low species richness (Snelgrove 2001) and high evenness (Hillebrand et al. 2007). In these areas the role of producer tends to be dominated by a small number of species able to monopolise resources under ambient conditions. Corresponding benthic communities are dominated by the taxa best able to use the associated products (Lenihan & Micheli 2001) or withstand periods of anoxia imposed by excess organic input (Dell'Anno et al. 2002). In contrast, oligotrophic waters are often home to low biomass assemblages with high species richness, including a large proportion of endemic taxa (Poore et al. 2008). Coral reefs, areas of high biomass and species richness occurring in oligotrophic waters, are an exception. The symbiosis between coral polyps and their resident zooxanthellae allows higher productivity than would otherwise occur in the ambient conditions and the spatial complexity and diversity of habitats provided by hard corals competing for space and light promotes a high corresponding richness of invertebrate and fish life, in turn supporting a rich community of predators (Cribb et al. 1994). Primary production can be estimated from satellite or airborne spectral analysis of chlorophyll in surface waters (Parmar et al. 2006). While productivity is directly linked to marine biodiversity, the relationship has yet to be fully explored as a predictive surrogate over large scales.

### 3.5.2. Organic Carbon

Detrital matter derived from primary productivity and the wastes of secondary production comprise a valuable resource in the photic zone and, excepting chemosynthetic systems, almost the only energy input to the aphotic zone (Carney 2005; Vetter 1995). This material settles in particles of various sizes, among which larger particles such as faecal pellets (Angel 1984) and marine snow (Alldrege and Silver 1988) are particularly important. It is generally accepted that the flux of particulate organic carbon (POC) from the euphotic zone controls the biomass and abundance of deep-sea benthos. This notion was originally based on observations of high benthic standing-crops beneath productive equatorial and near-shore waters, and low standing-crops underlying oligotrophic gyres (Belyayev et al. 1973; Hessler 1974; Gage and Tyler 1991; Rowe et al. 1991; Blake and Hilbig 1994). However, a direct coupling between pulse-like sedimentation events and the activity of benthic fauna has become clear in recent years (Aberle and Witte, 2003 and references therein). The detailed nature of this coupling remains poorly understood because there have been few studies which combine both types of measurements. One good example is a study by Smith et al. (1997) in the equatorial Pacific, in which strong and significant correlations ( $r^2 > 0.9$ ) were found between both megafaunal- (phototranssects) and macrofaunal (enumerated from box core samples) abundances and annual POC fluxes. An interesting conclusion from this study was that macrofaunal abundance might potentially serve as a proxy (i.e. surrogate) for POC flux in low energy abyssal habitats, implying that the macrofauna themselves are either more widely or more easily measured than POC fluxes (see also Rowe et al. 1991; Cosson et al. 1997).

The main way to directly measure POC fluxes is using sediment traps, and at the time of the Smith et al. (1997) study, approximately 37 records integrating annual time scales existed in the open ocean (Lampitt and Anita 1997). More recently, Seiter et al (2005) drew on particle-trap data from 61 locations, and produced a global map of minimum POC flux to the seafloor which was based on global estimates of diffusive oxygen uptake. This map, and the global map of total organic carbon (TOC) concentrations that underpins it (Seiter et al. 2004), may prove useful in making first order approximations of benthic productivity over broad scales, assuming that benthic communities are not compromised by sediment de-oxygenation. Indeed, the relationships between diversity and POC fluxes (or other productivity proxies) are scale-dependent and may be complicated by other variables that influence diversity including bottom-water oxygen concentration, hydrodynamic regime and the stability of the physical environment (Levin et al. 2001). The study by Smith et al. (1997) of the abyssal equatorial Pacific was unique in the respect that the productivity gradient found there varied independently of other influential factors. Thus, the uni-modal relationships between productivity and diversity expected over broad scales in the deep sea are plausible but are not well substantiated due to: (i) the complexity of interacting factors; and (ii and iii) an inability to both clearly identify mechanisms through which productivity influences deep-sea diversity, and to accurately place available studies in a continuous productivity gradient (Levin et al. 2001).

Proxies of POC fluxes such as TOC, TOC:TN ratios, biochemical markers and pigments in sediment have proven useful in explaining more localized patterns of biodiversity. TOC is undoubtedly the most widely measured of these parameters (Seiter et al. 2005), and, where a consistent and robust method (Galy et al. 2007) has been applied to its measurement, TOC can be a useful surrogate for biomass, deposit-feeding taxa, and

community structure (Gogina et al. 2010). However, its application is limited to interpolations from physical samples (Levin and Gage 1998) as no remote sensing proxy is available. Moreover, correlations between TOC and diversity measures are not always found (Cartes et al. 2002; Danovaro et al. 1995) because a large proportion of TOC in sediment may be refractory and thus resistant to bacterial degradation. Sediment grain size can also affect the amount of biologically available organic matter (OM) in shallow soft sediments (Taghon 1982). Small particles have larger surface area per unit volume than large particles, offering greater habitat for micro-organisms (Fauchald and Jumars 1979; Neira and Hoepner 1994; Petch 1986) and associated organic matter. Some deposit-feeding species use size-specific foraging mechanisms to select and ingest fine sediments (Butman and Grassle 1992; Sebesvari et al. 2006; Thiyagarajan et al. 2005), but both selective and non selective deposit feeders exhibit settlement preferences for sediments with high concentrations of readily available organic carbon (Post et al. 2006; Snelgrove and Butman 1994).

The availability, freshness or quality of OM pertains to the labile fraction, which consists mainly of lipids, carbohydrates, proteins and nucleic acids (Danovaro et al. 1993; 1995; 2001). Several useful biochemical parameters have been derived to describe the lability of OM (Cowie and Hedges 1992; Danovaro et al. 1995; Wakeham et al. 1997; Dauwe et al. 1999), and some of these have proven useful for explaining different diversity indices (Danovaro et al. 1995; Cartes et al. 2002). Such measures, however, often require a high degree of discipline expertise (and advanced techniques), and as such are unlikely to be widely employed in the capacity of surrogates. However, the Chlorin Index (CI) (Schubert et al. 2005) is a simple analytic measurement of OM lability, whose reliability has been demonstrated by comparison to more advanced techniques (e.g. Dauwe Index, total hydrolysable amino acids, and %  $\beta$ -alanine as non-protein amino-acid, and sulfate reduction rates) (Schubert et al. 2005). It is a measure of the amount of chlorophyll (and its degradation products) that could be transformed to phaeophytin, and is expressed as the ratio of the fluorescence intensity of a sediment sample extracted in acetone and subject to HCl treatment to that of the original sediment sample (Schubert et al. 2005). CIs were found to correlate well with an index of track richness developed from photographic stills of seabed sediments, which conveyed differences in the diversity of tracks, faecal casts, burrows and mounds of benthic biota in deep-sea sediments of the Lord Howe Rise (Dundas and Przeslawski 2009). Comparison with this index shows a greater diversity of animal traces, and potentially more metazoan activity, in sediments of apparently higher food quality. CIs thus show promise as an easily measured geochemical surrogate of biodiversity for regions where organic loads are not expected to give rise to significant sediment anoxia.

### 3.6. Oceanography

Oceanographers measure and model variables that directly influence the physiology and behaviour of marine organisms (temperature, salinity, pH), variables affecting productivity (nutrient concentrations, temperature and light intensity: see section on productivity) and the currents that affect larval distributions. Some factors (pH and salinity) vary sufficiently over regional and global scales to show correspondence to biological patterns (Williams & Bax 2001) but are sufficiently uniform at a local scale (with the exception of estuarine systems) to preclude their use in local surrogacy analyses

(Bamber et al. 2008). Dissolved oxygen has been identified as a key predictor of biodiversity in deep sea sediments (Levin & Gage 1998).

In addition to determining local water properties and delivering food and oxygen, ocean currents are important to the dispersal of many marine organisms which, in turn, determines the potential distribution of many benthic taxa. Most larvae and algal propagules spend their early development adrift and must attempt to settle where the prevailing currents take them. With larval periods ranging from hours (e.g., tropical ascidians in Cloney et al. 2006) to four and a half years (e.g., gastropod in Strathmann & Strathmann 2007), the scope for currents to act as a surrogate for potential richness is substantial where both life histories and water movements are well known. The relative rarity of long larval life histories make local currents (~tens of kilometers) more important than regional currents in determining benthic larval supply (Palumbi 2001), but even groups with well known life histories have frustrated attempts to predict geographic assemblies (Shulman & Bermingham 1995). Stevens and Connolly (2004) considered local scale current speed as an abiotic variable in their assessment of surrogates in Moreton Bay, Australia, but found its predictive capacity negligible. In addition to understanding larval supply patterns, the proximity of any given sample to diversity hotspots must be taken into consideration (Bellwood et al. 2005).

Further information on biodiversity patterns as they relate to oceanographic variables can be found in reviews by Hall (1994), Wolanski (2001), and Levin et al. (2001).

## 4. Utility Considerations

### 4.1. Temporal Variations

Knowledge of temporal variation in potential surrogacy relationships is limited. Most of our knowledge of the temporal variations in marine biodiversity comes from intertidal habitats, where some assemblages show pronounced seasonal changes, e.g. nematodes (Hourston et al. 2005), and some other systems show no seasonal variation, e.g. flatworms (Dittmann 1998). Although, we have a good understanding of temporal variation in broad processes such as organic input (Gooday 2002) and numerous studies investigate the relationships between abiotic and biotic factors (see previous sections in this review); we still have limited understanding of how such relationships change over weeks, seasons, and years.

Consideration of temporal variation increases our chances of identifying surrogacy relationships. Some relationships may change or only be detectable during certain times of day, seasons or years, likely due to phenology of biota or inter-annual variations in climate, recruitment, and other processes. Diurnal patterns may play an important role in the detection of surrogacy relationship, particularly regarding sampling mobile organisms. The composition of deep sea demersal fish changes between night and day (Suetsugu & Ohta 2005), and the interpretation of data will therefore depend on the diurnal variation of an assemblage and the timing of sampling.

Seasonal patterns also play an important role in the abundance of many taxa, particularly in shelf waters where benthic communities can be highly regulated by seasonal upwelling and thermal changes. Reproduction of some deep-sea animals are apparently synchronised with the seasonal sinking of phytoplankton blooms at the surface (Turner 2002). Seasonal variation in potential surrogacy relationships may also be related to depth. In the eastern Mediterranean, both abiotic factors and detritus-driven relationships in mesozooplankton vary among seasons in the upper 750 m (Koppelman & Weikert 2007). Between 750-1050 m, differences between seasons were not detected, while deeper waters between 1050-2250 m showed a slightly later seasonal increase in abundance than upper waters, likely due to lag between bloom events and the arrival of their products. No temporal changes in abundance were detected below 2250 m (Koppelman & Weikert 2007). Seasonal variations in abundance and community composition at the bottom may mirror variations noted in zooplankton communities across various depths, but further research is needed to test this hypothesis.

On a larger temporal scale, inter-annual variation has been detected in many communities, including homogenous muddy sand, in which large infauna were neither spatially nor temporally uniform (Parry et al. 2003). The causes underlying such temporal variation remain unknown but may nevertheless affect the detection and utility of surrogacy relationships.

The magnitude of temporal variation may be related to organism size. On the abyssal plain, the abundance of meiofauna and near-surface deposit feeders spiked during the annual pulse of organic input, while the abundance of macrofaunal polychaetes increased much more gradually, lagging behind the pulse (Galeron et al. 2001). In a review of soft-sediment community responses to organic input, Gooday (2002) showed that temporal patterns between biotic and abiotic factors vary according to size classes, specifically that only bacteria and foraminifera show a consistent and direct response to seasonal organic inputs, while populations of larger fauna generally show reduced,

lagged, or unmeasurable responses (Gooday 2002). In contrast, Grove et al. (2006) found the abundance of microbes and meiofauna is not correlated with organic content in Chatham Rise, eastern New Zealand, suggesting that size-dependent temporal patterns in abundance may vary across regions, depths, and habitats. Assuming niches will be filled by functionally equivalent or related organisms through time and assessing biota in terms of functional groups or at coarse taxonomic resolutions may offer a means to circumvent this shortfall.

#### **4.2. Surrogacy in the Deep Sea**

The relationships between abiotic and biotic variables can be quite different in deep and shallow waters. Compared to shallow water fauna, deep water fauna live in darkness at relatively stable temperatures in the absence of phytoplankton and other photosynthetic organisms. Thus, potential surrogates for shallow water biodiversity, such as temperature, wave action, depth (as a proxy for light attenuation), and primary productivity may not apply to deep sea biodiversity. Conversely, factors such as organic composition of sediment may have far more importance in deep-sea environments, in which deposit-feeding species dominate (Levin et al. 2001).

Potential surrogate relationships are far less well-known in deep waters (>1000 m) than in shallow waters, probably due to the comparative difficulty of sampling biota and the lack of knowledge of taxa and life histories. The few studies that do investigate surrogacy in the deep-sea have focused primarily on depth (Brandt et al. 2007), and most are from the northern Atlantic (Levin et al. 2001) or Antarctic regions (Ellingsen et al. 2007) particularly at chemosynthetic or ephemeral habitats such as hydrothermal vents, cold seeps, and whale falls. Increasing efforts are being made to characterise deep water biodiversity as nations seek to manage the resources of their exclusive economic zone.

#### **4.3. Surrogacy across Spatial Scales**

Quantification of spatial scale depends on three characteristics: 1) the size of each sample, 2) the distance between samples, and 3) the total area encompassing all samples (Parry et al. 2003). The importance of spatial and temporal variability has been recognised in terrestrial surrogacy research (Ernault et al. 2006), with broad spatial scales associated with better performance of abiotic surrogates (Sarkar et al. 2005) and increased effectiveness of lower taxonomic resolution (Larsen & Rahbek 2005). Unfortunately, very little is known about differential surrogacy relationships across broad and fine spatial scales in the marine environment. Freshwater studies suggest that broad spatial scales are more useful to detect community patterns than local scales (Paavola et al. 2006), but it is unknown if this applies to marine systems. The influence of energy availability has been shown to differentially affect faunal distributions across spatial scales in the Mediterranean (Gambi & Danovaro 2006), suggesting that the detection of relationships between abiotic and biotic factors is dependent on the spatial scale used. Marine invertebrates show differential variability across spatial scales according to phylum (Anderson et al. 2005), and this variability is likely to be regionally-specific. For example, tropical Australian sponges are best represented at a medium scale of diversity, rather than at local or coarse regional scales (Hooper et al. 2002). The ability to use surrogates effectively at fine spatial scales may be related to the heterogeneity of a given area. Nevertheless, research in the northeast Atlantic found a high level of community

patchiness in a very homogenous soft sediment environment, highlighting the need to consider physical processes that operate on broad spatial scales as well as biological processes that operate on finer scales (Parry et al. 2003). There has been no concentrated effort to understand how spatial scale affects surrogacy relationships in the marine environment.

The effects of spatial scale on surrogacy relationships obviously have important ramifications for the detection of any such relationships (Gambi & Danovaro 2006). Many benthic species are restricted to a very small proportion of sites sampled (e.g. polychaetes and isopods in (Ellingsen et al. 2007), isopods in (Brandt & Schnack 1999), and it would not be practical to use such species to investigate surrogacy relationships except at extremely fine scales. In addition, consideration of spatial scale is crucial not only to evaluate the utility of a potential surrogate, but also to appropriately address the research hypothesis or management strategy. For example, if surrogacy research is needed to better inform the design of protected areas within a bay, basin, or other relatively small area, a regional spatial scale may be too large (Greene et al. 1999).

#### **4.4. Combined Effects of Potential Surrogates on Benthic Biodiversity**

Some physical factors may interact to synergistically or antagonistically affect fitness, fecundity, behaviour, or demography of one or more species (Folt et al. 1999); but until recently most surrogacy research has focused on only one variable at a time. In many aquatic communities, biodiversity and biogeography are best associated with multiple environmental factors and rarely defined according to just one (Dauer et al. 2000; Hagberg et al. 2003; Kennard et al. 2006).

Abundance of certain species may correlate with abiotic variables such as depth and organic carbon flux, but the actual cause of these relationships may be due to a biotic factor such as predator density which is itself directly regulated by abiotic factors (Micheli et al. 2002). Many sessile invertebrates provide habitat and food to other fauna so the presence of these habitat-forming invertebrates directly affects the abundance of other invertebrates and fish (Tissot et al. 2006, but see Jensen & Frederickson 1992; Klitgaard 1995). The presence of infaunal species can change the sediment composition and ecology, thus making it difficult to determine causes and effects (Bremner et al. 2006). In addition, biological processes in separate oceanic zones can affect each other. Benthic-pelagic coupling is well-studied across a variety of systems (Turner 2002), and the contributions of pelagic detritus to benthic processes are extremely variable and affected by several interacting factors, including pelagic community composition, trophic interactions, sinking rates, and oceanographic conditions (Turner 2002). Indeed, the bathymetric zoning of deep-sea fauna has been linked to increasing pressure and decreasing food availability with depth (Carney 2005). Identification of potential surrogacy relationships is only the first step and is relatively straightforward compared to the interpretation of such relationships, due to the linkages between some abiotic factors and physical and biological processes (Figure 1) (Hagberg et al. 2003; Levin et al. 2001). Nonetheless, understanding the causes underlying surrogacy relationships, including interactions between multiple abiotic and biotic variables, will enhance our ability to develop predictive models across a range of systems (Parry et al. 2003).

#### 4.5. Utility of Different Sampling Regimes

Biological sampling can be undertaken through both destructive and non-destructive methods at a broad range of spatial scales. The nature and strength of surrogacy relationships may vary across sampling regimes because each sampling method is associated with discrete optimal habitats and taxa. For example, most grabs target sessile epifauna and small infauna and tend to underestimate larger sedentary animals due their ability to bury deeper than the sampler penetrates (Parry et al. 2003). Tracks, mounds, and burrows can be used to gauge benthic activity (Parry et al. 2003), but the persistence of biological structures is related to currents and disturbance, as tracks will persist far longer in low disturbance environments (e.g. deep sea habitats) (Dundas & Przeslawski 2009, Wheatcroft et al. 1989). Thus, the correlation between tracks and biodiversity will be more immediate in high disturbance environments and detection of associated surrogacy relationships may vary with disturbance rates across regions. To date there has been little research investigating whether sampling methods vary in their suitability to detect relationships between physical surrogates and biodiversity of marine benthic communities. Such research will provide valuable insight into the appropriate methods to use for biological collection, as well as proper interpretation of data analyses and perhaps lead to distinct biodiversity concepts being mated to particular samplers.

Regardless of the sampling method used, species accumulation curves are important to compare diversity among habitats, particularly among large bathymetric gradients. Species accumulation curves display the number of species encountered with increasing sampling effort; an area is considered well-sampled when the curve asymptotes. They are important tools to identify the effectiveness of biodiversity estimates. For example, in a comparison of shallow- and deep-water diversity, even when sampling was conducted for the same duration, diversity may still not be comparable if the species accumulation curve for the deep-water system has not yet reached its asymptote (Levin et al. 2001).

The application of technologies that increase the data returned per unit effort of sampling: e.g. multibeam sonar (Kostylev et al. 2001), acoustic Doppler current profilers (Harris et al. 2008) and airborne remote sensing (Mumby et al. 1997); allow potential surrogates to be nominated and tested. Biological data remains the limiting factor, but targeted surrogacy research is gradually revealing species richness patterns that contribute to our understanding of ecological processes and the identification of effective prediction tools with which marine biodiversity can be managed in an informed fashion.

#### 4.6. Management Decisions at Various Scales

The merit of researching surrogacy relationships is the increased capacity to manage marine resources with confidence that biodiversity concerns have been adequately addressed. This confidence will vary according to how well the information required to meet goals of marine resource managers has been addressed by surrogacy research and by the scale at which a decision is expected to influence environmental outcomes. While biogeographic maps and classifications have been widely applied to natural resource management in terrestrial systems (Vierros et al. 2008), their application to marine resources is still developing. Here, the use of surrogate variables in marine resource management is examined using examples at increasing spatial scales.

Until 2002 most coastal waters of Victoria (Australia) were legally regarded as unreserved public land (Environmental Conservation Council 2000). The Victorian Environment Conservation Council adopted a community, habitat and ecosystem approach to categorising Victorian marine resources and choosing sites for marine protected areas along the Victorian coast (O'Hara 2000). This strategy was chosen because habitat protection was thought to be a sound means of maintaining existing biodiversity (Environmental Conservation Council 2000), defined as “the natural diversity of all life: the sum of all our native flora and fauna, the genetic variation within them, their habitats, and the ecosystems of which they are an integral part,” and because of the difficulty involved in describing the highly endemic local marine flora and fauna at the species level (Poore & O'Hara 2007). Extensive species level surveys and genetic studies were acknowledged as a possible route to defining aspects of biodiversity missed by habitat level characterisation but these could not be applied as widely or quickly as required. An existing model of five coastal bioregions defined by bathymetry and benthic macrofauna distributions was therefore used to establish thirteen Marine National Parks and eleven Marine Sanctuaries protecting 5.3 % of the Victorian coast (Ferns et al. 2003) under the *National Parks Act 1975* (Vic).

Focusing on a broad scale definition of biodiversity allowed the Environment Conservation Council to create a management strategy in a timely fashion at the expense of the ability to manage resources at the species level, although the protected area selection criteria applied to Victoria's marine resources had sufficient flexibility to allow management decisions to take into account individual species with an eye for large, charismatic taxa such as the giant kelp and the weedy seadragon (Ferns et al. 2003).

Terrestrial resource regionalisations have led the way in surrogate-based management of biodiversity at regional scales. It has tended to use some biological attributes along with physical attributes within a classification to produce regions (Ferns 1999). While the vast majority of national level work has been land-based over the past 50 years, marine classification maps of some form have been produced for most of the world's shelf waters (Vierros et al. 2008). Regionalisations have been made using numerical classification techniques or through the synthesis of patterns and processes by experts. Utilizing existing knowledge and classifications, and bringing together experts from each jurisdiction to assimilate and combine existing classifications into single national regionalisations has been the most successful approach for achieving useful outcomes. The Interim Biogeographic Regionalisation for Australia (Thackway & Cresswell 1995) and the Interim Marine and Coastal Regionalisation for Australia (IMCRA 2006) provided sufficient scientific synthesis of known biophysical attributes while also acknowledging and utilizing existing classifications at the State level. Both projects were developed to assist in planning national systems of protected areas, terrestrial and marine respectively, in accordance with several national and international environmental obligations (Cresswell & Thackway 1998). Both of these regionalisations have been updated in recent years and extended through the addition of new information and levels of detail (Heap et al. 2005).

Perhaps the best measure of the usefulness of any classification or regionalisation is not in the exact nature of the boundaries that have been created either by computer or human eye, but rather in their application in biodiversity management. These regionalisations have been adopted and used for a multitude of conservation and sustainable use purposes at national and regional levels, and are now widely accepted as the standard biogeographic framework for use in both scientific and management studies.

In international waters biological sampling has been disparate (Harris et al. 2007b) other than around novel features, such as seamounts (Rowden et al. 2005) or hydrothermal vents (Van Dover 2000). Surrogate-based management of biodiversity is the only practical means to make reasoned decisions about high seas resource management. To fulfil their obligations under the United Nations Convention on the Law of the Sea, nations that use high seas resources are expected to monitor that use and put in place mechanisms to protect marine resources beyond their national boundaries (Ardron et al. 2008), but the biological information required to establish effective networks of protected areas does not yet exist (Harris et al. 2007a). Global regionalisations of high seas resources have a long history but have only recently applied spatial approaches to their classifications (Table 4.1).

**Table 4.1.** Approaches to classifying regions in the high seas (after Vierros et al. (2008)).

<b>Project</b>	<b>Basis of zonation</b>	<b>Reference</b>
Zoogeography of the Sea	Intuitive divisions of faunal zones	(Ekman 1953)
Marine Biogeography	Review of the work of Eckman and others	(Hedgpeth 1957)
Marine Zoogeography	Taxonomic biogeography	(Briggs 1974)
Classification of Coastal and Marine Environments	Spatial approach to Brigg's work	(Hayden et al. 1984)
Large Marine Ecosystems	Bathymetry, hydrography, productivity, trophic dependence	(Sherman & Alexander 1989)
Global Representative System of Marine Protected Areas	Intuitive consideration of biogeographic representations	(Kelleher et al. 1995)
Ecological Geography of the Sea	Surface productivity	(Longhurst 1998)
Ecoregions: the Ecosystem Geography of the Oceans and Continents	Latitudinal belts and patterns of oceanic circulation	(Bailey 1998)
Marine Ecosystems of the World	Review of previous work and intuitive input from experts	(Spalding 2007)
High Seas Marine Protected Areas	Multivariate analysis of depth, slope, sediment thickness, primary productivity, seabed temperature and dissolved oxygen	(Harris & Whiteway 2009)

The 53 713 polygons resulting from Harris & Whiteway (2009) were used to identify hotspots of habitat heterogeneity, potential candidate areas for international protection. Spatial classifications have been used to establish marine protected areas in the high seas (Ardron et al. 2008), but considerable effort and money will be required to police resource use and make high seas marine protected areas more than paper parks (Mossop 2004).

Various different numerical classification techniques have been used in an attempt to reach the holy grail of geographic regionalisation that can explain the distribution of the biotic realm in space and, for the adventurous, time. No single classification methodology has gained universal acclaim. The classification of a set of thematic environmental attributes recorded at a multitude of sites or taken from a uniform grid has been undertaken at a variety of scales as the basis for nature conservation planning and management (Lewis et al. 1991; Lyne & Hayes 2005; Mackey et al. 1988; Mackey et al. 1989; Thackway & Cresswell 1992). To date no numerically produced regionalisation or assessment has been widely adopted for use in biodiversity management. The reasons for this are complex and relate to the significant difficulties in utilising the correct combination of attributes in a classification that will replicate the patterns and processes recognized by humans, as well as institutional and governance complexity in how we manage biodiversity.

## 5. Synthesis

When decisions about the conservation or use of an area must be made, information about the biodiversity values of that area may not exist or be ignored, thereby hindering the development of appropriate resource management plans. This is often the case in marine benthic systems as they are difficult and therefore expensive to study and often have poor historical coverage compared to their terrestrial or pelagic equivalents. Exploring new ways of measuring and understanding the relationship between biota and the physical environment offers an opportunity to address challenges for biodiversity management using existing and easily gathered data. Surrogacy research is a rapidly-growing field, and useful relationships have been identified in several habitats, particularly with regard to sediment parameters. Some variables with direct influence on presence and abundance of benthic species (temperature, salinity, pH) vary over spatial and temporal scales that make them appropriate to apply at regional scales. Their broader influence can be invoked from less directly influential variables that correlate to longer or larger trends (depth, latitude, oceanic currents), but at those scales, confounding factors (larval/propagule supply, proximity to diversity hotspots) must be taken into account.

The use of remote sensing of chlorophyll-*a* as a surrogate of benthic biodiversity is hampered by poor understanding of the relationship between primary productivity and benthic assemblies. The general model of higher benthic richness in areas of lower productivity, an inversion of that observed in terrestrial systems, is too readily confounded by other variables to be confidently applied to management decisions.

The relationship between acoustic data and variables with direct (sediment hardness, rugosity and slope) and indirect (depth) influence on benthic assemblages, coupled with recent advances in our ability to map large areas of the seafloor using acoustic equipment offer the greatest potential for surrogacy based management of marine biodiversity. Ground-truthing of habitat classifications being derived from acoustic

mapping is an important but painstaking process, hampered by gaps in our knowledge including the difficulty of identifying organisms to fine taxonomic levels and the comparative shortage of information about meiofauna, microorganisms and deep sea ecosystems.

Patterns of benthic biodiversity observed in many regions today may not reflect the distribution and diversity of species of just a few decades ago, as overfishing, trawling, introduced species, and climate change have altered habitats. Therefore, physical surrogates of present-day patterns of biodiversity may carry some uncertainty about the degree of impact study areas may have suffered. Impacts affecting areas studied to refine abiotic surrogacy models must be considered before predictions based on those relationships are scaled beyond the impacted areas. Predictions of patterns of biodiversity based on surrogacy relationships may be considered ‘potential’ patterns that may occur in impacted areas when environmental pressures are reduced.

The range of biodiversity concepts and metrics available make choosing an appropriate approach to gathering and interpreting surrogacy data challenging. The results of any management decision should be kept in mind – is the system in question being managed to protect a particular habitat, for maintenance of ecosystem services, to create refugia for genetic diversity, or to insure particular taxa against extinction? – when a biodiversity measure is being considered. A thorough knowledge of the mathematical properties of that measure should inform any subsequent analyses based on it.

Our ability to measure abiotic variables in marine systems and our understanding of the relationships between those variables and biodiversity have increased with our need to make predictions about biodiversity in otherwise unsampled areas. The first test of abiotic surrogacy involves validating those predictions. The second test will be whether or not predictions based on abiotic surrogates are found to be useful in marine resource management.

## 6. References

- Abele, L. G., & Patton, W. K. (1976). The size of coral heads and the community biology of associated decapod crustaceans. *Journal of Biogeography*, 3, 35-47.
- Aberle, N., and Witte, U. 2003. Deep-sea macrofauna exposed to a simulated sedimentation event in the abyssal NE Atlantic: in situ pulse-chase experiments using <sup>13</sup>C-labelled phytodetritus. *Marine Ecology Progress Series* 251: 37-47.
- Allredge, A.L. and M.W. Silver (1988). Characteristics, dynamics and significance of marine snow. *Progress in Oceanography* 20: 41-82.
- Alongi, D. M. (1990). The ecology of tropical soft-bottom benthic ecosystems. *Oceanography and Marine Biology*, 28, 381-496.
- Anderson, M. J., Diebel, C. E., Blom, W. M., & Landers, T. J. (2005). Consistency and variation in kelp holdfast assemblages: Spatial patterns of biodiversity for the major phyla at different taxonomic resolutions. *Journal of Experimental Marine Biology and Ecology*, 320(1), 35-56.
- Anderson, T. J., Cochrane, G. R., Roberts, D. A., Chezar, H., & Hatcher, G. (2007). A rapid method to characterize seabed habitats and associated macro-organisms. In: B. J. Todd, & H. G. Greene, *Mapping the Seafloor for Habitat Characterization: Geological Association of Canada Special Paper 47* (pp. 71-79): Geological Association of Canada.
- Angel, M. V., (1984). Detrital organic fluxes through pelagic ecosystems: In: *Flows of energy and materials in marine ecosystems: theory and practice*, M. R., J. Fasham, editor, NATO Conference Series IV, Marine Science, Plenum Press, New York, pp. 475-516.
- Ardron, J., Gjerde, K., Pullen, S., & Tilot, V. (2008). Marine spatial planning in the high seas. *Marine Policy*, 32(5), 832-839.
- Attrill, M. J., Stafford, R., & Rowden, A. A. (2001). Latitudinal diversity patterns in estuarine tidal flats: indications of a global cline. *Ecography*, 24, 318-324.
- Ault, T., & Johnson, C. (1998). Spatially and temporally predictable fish communities on coral reefs. *Ecological Monographs*, 1, 5-50.
- Auster, P. J., & Langton, R. W. (1999). The effects of fishing on fish habitat. *American Fisheries Society Symposium*, 22, 150-187.
- Austin, M. P., Nicholls, A. O., & Margules, C. R. (1990). Measurement of the realized qualitative niche - environmental niches of 5 *Eucalyptus* species. *Ecological Monographs*, 60(2), 161-177.
- Austin, M. P., & Smith, T. M. (1989). A new model for the continuum concept. *Vegetatio*, 83(1-2), 35-47.
- Bagnold, R. A. (1963). Mechanics of Marine Sedimentation. In: M. N. Hill, *The Sea*, vol. 3 (pp. 507-527). New York: Wiley Interscience.
- Bailey (1998). *Ecoregions: the Ecosystem Geography of the Oceans and Continents*. New York: Springer-Verlag.

- Bamber, R. N., Evans, N. J., & Robbins, R. S. (2008). The marine soft-sediment benthic communities of Hong Kong: a comparison of submarine cave and open habitats. *Journal of Natural History*, 42(9-12), 953-965.
- Barnard, J. L. (1991). Amphipodological agreement with Platnick. 25, 1675-1676.
- Barnes, D. K. A., & Griffiths, H. J. (2008). Biodiversity and biogeography of southern temperate and polar bryozoans. *Global Ecology and Biogeography*, 17(1), 84-99.
- Barrett, N., Sanderson, J. C., Lawler, M., Halley, V., & Jordan, A. (2001). Mapping of inshore marine habitats in south eastern Tasmania for marine protected area planning and marine management. Technical Report. Tasmanian Aquaculture and Fisheries Institute: Hobart.
- Barros, F., Underwood, A. J., & Archambault, P. (2004). The influence of troughs and crests of ripple marks on the structure of subtidal benthic assemblages around rocky reefs. *Estuarine Coastal and Shelf Science*, 60(4), 781-790.
- Beaman, R. J., Daniell, J. J., & Harris, P. T. (2005). Geology-benthos relationships on a temperate rocky bank, eastern Bass Strait, Australia. *Marine and Freshwater Research*, 56(7), 943-958.
- Beaman, R. J., & Harris, P. T. (2007). Geophysical variables as predictors of megabenthos assemblages from the northern Great Barrier Reef, Australia. In: B. J. Todd, & H. G. Greene, *Mapping the Seafloor for Habitat Characterization*: Geological Association of Canada.
- Begon, M., Harper, J. L., & Townsend, C. R. (1996). *Ecology: Individuals, Populations and Communities*. Melbourne: Blackwell Scientific Publications.
- Bell, G. (2000). The distribution of abundance in neutral communities. *American Naturalist*, 155(5), 606-617.
- Bellwood, D. R., Hughes, T. P., Connolly, S. R., & Tanner, J. (2005). Environmental and geometric constraints on Indo-Pacific coral reef biodiversity. *Ecology Letters*, 8(6), 643-651.
- Belyayev, G.M., Vinogradova, N.G., Levenshteyn, R.Y., Pasternak, F.P., Sokolva, M.N., Filtatova, Z.A., (1973). Distribution patterns of deep-water bottom fauna related to the idea of the biological structure of the ocean. *Oceanography* 13: 114-121
- Benkendorff, K., & Davis, A. R. (2002). Identifying hotspots of molluscan species richness on rocky intertidal reefs. *Biodiversity and Conservation*, 11, 1959-1973.
- Bertrand, Y., Pteijel, F., & Rouse, G. W. (2006). Taxonomic surrogacy in biodiversity assessments, and the meaning of Linnaean ranks. *Systematics and Biodiversity*, 4(2), 149-159.
- Bindoff, N. L., Willebrand, J., Artale, V., Cazenave, A., Gregory, J., Gulev, S., Hanawa, K., Le Quere, C., Levitus, S., Nojiri, Y., Shum, C. K., Talley, L. D., & Unnikrishnan, A. (2007). Observations: oceanic climate change and sea level. In: S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, & H. L. Miller, *Climate Change 2007: the Physical Science Basis*. Cambridge, UK: Cambridge University Press.
- Blake, J.A., and Hilig, B. (1994). Dense infaunal assemblages on the continental slope off Cape Hatteras, North Carolina. *Deep-Sea Research II* 41: 875-900.

- Blott, S. J., & Pye, K. (2001). GRADISTAT: A grain size distribution and statistics package for the analysis of unconsolidated sediments. *Earth Surface Processes and Landforms*, 26(11), 1237-1248.
- Bolton, J. J. (1996). Patterns of species diversity and endemism in comparable temperate brown algal floras. *Hydrobiologia*, 327, 173-178.
- Bonsdorff, E., & Pearson, T. H. (1999). Variation in the sublittoral macrozoobenthos of the Baltic Sea along environmental gradients: A functional-group approach. *Australian Journal of Ecology*, 24(4), 312-326.
- Bouchet, P. (2006). The Magnitude of Marine Biodiversity. In: C. M. Duarte, *The Exploration of Marine Biodiversity: Scientific and Technological Challenges* (pp. 31-64). Paris: Fundacion BBVA.
- Brandt, A., Brix, S., Brokeland, W., Choudhury, M., Kaiser, S., & Malyutina, M. (2007). Deep-sea isopod biodiversity, abundance, and endemism in the Atlantic sector of the Southern Ocean-Results from the ANDEEP I-III expeditions. *Deep-Sea Research Part II*, 54(16-17), 1760-1775.
- Brandt, A., & Schnack, K. (1999). Macrofaunal abundance at 79N off East Greenland: opposing data from epibenthic-sledge and box-corer samples. *Polar Biology*, 22, 75-81.
- Bremner, J., Rogers, S. I., & Frid, C. L. J. (2006). Matching biological traits to environmental conditions in marine benthic ecosystems *Journal of Marine Systems*, 60, 302-316.
- Brey, T., Klages, M., Dahm, C., Gorny, M., Gutt, J., Hain, S., Stiller, M., Arntz, W. E., Wägele, J.-W., & Zimmermann, A. (1994). Antarctic benthic diversity. *Nature*, 368, 297.
- Briggs, J. C. (1974). *Marine Zoogeography*. New York: McGraw-Hill.
- Brown, C. J., Hewer, A. J., Meadows, W. J., Limpenny, D. S., Cooper, K. M., Rees, H. L., & Vivian, C. M. G. (2001). *Mapping of gravel biotopes and an examination of the factors controlling the distribution, type and diversity of their biological communities*. Lowestoft, CEFAS.
- Brown, J. H., & Gibson, A. C. (1983). *Biogeography*. St Louis: C.V. Mosby Co.
- Butman, C. A., & Grassle, J. P. (1992). Active habitat selection by *Capitella* sp. I larvae. 1. Two-choice experiments in still water and flume flows. *Journal of Marine Research*, 50(4), 669-715.
- Callaway, R. (2006). Tube worms promote community change. *Marine Ecology-Progress Series*, 308, 49-60.
- Carney, R. S. (2005). Zonation of deep biota on continental margins. *Oceanography and Marine Biology - an Annual Review*, Vol. 43, 43, 211.
- Cartes, J.E., Grémare, A., Maynou, F., Villora-Morenon, S., and Dinata, A. (2002). Bathymetric changes in the distributions of particulate organic matter and associated fauna along a deep-sea transect down the catalan sea slope (Northwestern Mediterranean). *Progress in Oceanography*, 53: 29-56.

- Chao, A., Chazdon, R. L., Colwell, R. K., & Shen, T. (2005). A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecology Letters*, 8, 148-159.
- Chivian, E., & Bernstein, A. (2008). *Sustaining Life: How Human Health Depends on Biodiversity*. Oxford University Press.
- Clark, M. R., Tittensor, D., Rogers, A. D., Brewin, P., Schlacher, T., Rowden, A., Stocks, K., & Consalvey, M. (2006). *Seamounts, Deep-Sea Coral and Fisheries: Vulnerability of Deep-Sea Corals to Fishing on Seamounts Beyond Areas of National Jurisdiction*. Cambridge, UK: UNEP-WCMC.
- Clarke, K. R., & Gorley, R. N. (2006). *PRIMER v6: User Manual / Tutorial*. Plymouth, U.K.: PRIMER-E Ltd.
- Clarke, K. R., & Warwick, R. M. (1998). A taxonomic distinctness index and its statistical properties. *Journal of Applied Ecology*, 35, 523-531.
- Clarke, K. R., & Warwick, R. M. (2001). *Change in Marine Communities: An Approach to Statistical Analysis and Interpretation*. Plymouth, U.K.: PRIMER-E Ltd.
- Cloney, R. A., Young, C. M., & Svane, I. (2006). Phylum Chordata: Urochordata. In: C. M. Young, *Atlas of Marine Invertebrate Larvae* (pp. 565-594). Barcelona: Elsevier.
- Coleman, N., Cuff, W., Moverley, J., Gason, A. S. H., & Heislars, S. (2007). Depth, sediment type, biogeography and high species richness in shallow-water benthos. *Marine and Freshwater Research*, 58(3), 293-305.
- Connell, J. H. (1978). Diversity in tropical rain forests and coral reefs. *Science*, 199, 1302-1310.
- Cosson, N., Sibuet, M., Galeron, J. (1997). Community structure and spatial heterogeneity of the deep-sea macrofauna at three contrasting stations in the tropical northeast Atlantic. *Deep-Sea Research I* 44: 247-269.
- Costello, M. J., Bouchet, P., Emblow, C. S., & Legakis, A. (2006). European marine biodiversity inventory and taxonomic resources: state of the art and gaps in knowledge. *Marine Ecology-Progress Series*, 316, 257-268.
- Cowie, G., and Hedges, J.I., 1992. Sources and reactivity of amino acids in a coastal marine environment. *Limnology and Oceanography*, 37(4) 703-724.
- Cox, C. B., & Moore, P. D. (2005). *Biogeography : an ecological and evolutionary approach* Blackwell Publishing.
- Crame, J. A. (2000). Evolution of taxonomic diversity gradients in the marine realm: evidence from the composition of recent bivalve faunas. *Paleobiology*, 26(2), 188-214.
- Creswell, I. D., & Thackway, R. (1998). Mapping a great big sea. *Global Biodiversity*, 8(1).
- Cribb, T. H., Bray, R. A., Barker, S. C., Adlard, R. D., & Anderson, G. R. (1994). Ecology and diversity of digenean trematodes of reef and inshore fishes of Queensland. *International Journal for Parasitology*, 24(6), 851-860.

- Culliney, J. L. (1974). Larval development of the giant scallop *Placopecten magellanicus* (Gmelin). *Biological Bulletin*, 147, 321-332.
- Currie, D. J., Mittelbach, G. G., Cornell, H. V., Field, R., Guegan, J. F., Hawkins, B. A., Kaufman, D. M., Kerr, J. T., Oberdorff, T., O'Brien, E., & Turner, J. R. G. (2004). Predictions and tests of climate-based hypotheses of broad-scale variation in taxonomic richness. *Ecology Letters*, 7(12), 1121-1134.
- Currie, D. R., & Parry, G. D. (1996). The effect of scallop dredging on a soft sediment community: a large-scale experimental study. *Marine Ecology Progress Series*, 134, 131-150.
- Danovaro, R., Fabiano, M., and Della Croce, N. 1993. Labile organic matter and microbial biomasses in deep-sea sediments (Eastern Mediterranean Sea). *Deep-sea Research I*, 40(5), 953-965.
- Danovaro, R., Della Croce, N., Eleftheriou, A., Fabiano, M., Papadopoulou, N., Smith, C., Tselepidis, A., 1995. Meiofauna of the deep Eastern Mediterranean Sea: distribution and abundance in relation to bacterial biomass, organic matter composition and other environmental factors. *Progress in Oceanography*, 36, 329-341.
- Danovaro, R., Dell'Anno, A., and Fabiano, M., 2001. Bioavailability of organic matter in the sediments of the Porcupine Abyssal Plain, northeast Atlantic. *Marine Ecology Progress Series*, 220, 25-32.
- Darwin, C. (1839). *The Voyage of the 'Beagle'*. London: Dutton.
- Dauer, D. M., Ranasinghe, J. A., & Weisberg, S. B. (2000). Relationships between benthic community condition, water quality, sediment quality, nutrient loads, and land use patterns in Chesapeake Bay. *Estuaries*, 23(1), 80-96.
- Dauwe, B.J., Middleburg, J.J., Herman, P.M.J., and Heip, C.H.R. 1999. Linking diagenetic alteration of amino acids and bulk organic matter reactivity. *Limnology and Oceanography*, 44(7), 1809-1814.
- Dayton, P. K. (1971). Competition, disturbance, and community organization: the provision and subsequent utilization of space in a rocky intertidal community. *Ecological Monographs*, 41, 351-389.
- Degraer, S., Verfaillie, E., Willems, W., Adriaens, E., Vincx, M., & Van Lancker, V. (2008). Habitat suitability modelling as a mapping tool for macrobenthic communities: An example from the Belgian part of the North Sea. *Continental Shelf Research*, 28(3), 369-379.
- Dell'Anno, A., Mei, M. L., Pusceddu, A., & Danovaro, R. (2002). Assessing the trophic state and eutrophication of coastal marine systems: a new approach based on the biochemical composition of sediment organic matter. *Marine Pollution Bulletin*, 44, 611-622.
- Denisenko, S. G., Denisenko, N. V., Lehtonen, K. K., Andersin, A. B., & Laine, A. O. (2003). Macrozoobenthos of the Pechora Sea (SE Barents Sea): community structure and spatial distribution in relation to environmental conditions. *Marine Ecology Progress Series*, 258, 109-123.
- Diaz, R. J., Solan, M., & Valente, R. M. (2004). A review of approaches for classifying benthic habitats and evaluating habitat quality. *Journal of Environmental Management*, 73, 165-181.

- Dittmann, S. (1998). Spatial and temporal patterns of platyhelminth assemblages in intertidal sediments of northeast Australia. *Hydrobiologia*, 383, 41-47.
- Dornelas, M., Connolly, S. R., & Hughes, T. P. (2006). Coral reef diversity refutes the neutral theory of biodiversity. *Nature*, 440, 80-82.
- Dundas, K., and Przeslawski, R. 2009. Deep-sea *Lebensspuren*: Biological features on the seafloor of the eastern and western Australian margin. *Geoscience Australia Record* 2009/26, pp. 76.
- Dye, A. H. (2005). Meiobenthos in intermittently open/closed coastal lakes in New South Wales: spatial and temporal patterns in densities of major taxa. *Marine and Freshwater Research*, 56(8), 1055-1067.
- Dye, A. H. (2006). Is geomorphic zonation a useful predictor of patterns of benthic infauna in intermittent estuaries in New South Wales, Australia? *Estuaries and Coasts*, 29(3), 455-464.
- Environmental Conservation Council (Victoria) (2000). *Marine, Coastal and Estuarine Investigation: Final Report*. Melbourne.
- Edgar, G. J. (1999). Experimental analysis of structural versus trophic importance of seagrass beds. I. Effects on macrofaunal and meiofaunal invertebrates. *Vie Et Milieu-Life and Environment*, 49(4), 239-248.
- Edgar, G. J. (2001). *Australian Marine Habitats*. Sydney: Reed New Holland Publishers Australia Pty Ltd.
- Ekman, S. (1953). *Zoogeography of the Sea*. London: Sigwick & Jackson.
- Ellingsen, K. E., Brandt, A., Ebbe, B., & Linse, K. (2007). Diversity and species distribution of polychaetes, isopods and bivalves in the Atlantic sector of the deep Southern Ocean. *Polar Biology*, 30(10), 1265-1273.
- Ellingsen, K. E., Clarke, K. R., Somerfield, P. J., & Warwick, R. M. (2005). Taxonomic distinctness as a measure of diversity applied over a large scale: the benthos of the Norwegian continental shelf. *Journal of Animal Ecology*, 74(6), 1069-1079.
- Ellingsen, K. E., & Gray, J. S. (2002). Spatial patterns of benthic diversity: is there a latitudinal gradient along the Norwegian continental shelf? *Journal of Animal Ecology*, 71(3), 373-389.
- Erlandsson, J., Kostylev, V., & Williams, G. A. (1999). A field technique for estimating the influence of surface complexity on movement tortuosity in the tropical limpet *Cellana grata gould*. *Ophelia*, 50, 215-224.
- Ernault, A., Freire-Diaz, S., Langlois, E., & Alard, D. (2006). Are similar landscapes the result of similar histories? *Landscape Ecology*, 21(5), 631-639.
- Etter, R. J., & Grassle, J. F. (1992). Patterns of species diversity in the deep sea as a function of sediment particle size diversity. *Nature*, 360, 1200-1216.
- Etter, R. J., & Mullineaux, L. S. (2001). Deep sea communities. In: M. D. Bertness, S. D. Gaines, & M. E. Hay, *Marine Community Ecology* (pp. 367-393). Sunderland MA: Sinauer.

Fauchald, K., & Jumars, P.A. (1979). The diet of worms: a study of polychaete feeding guilds. *Annual Review of Oceanography and Marine Biology* 17, 193-284.

Ferns, L., Bunting, B., Culquhoun, F., Coombes, B., De Kievit, J., De Luca, F., Edmunds, M., Griffiths, T., Holt, A., Leigh, C., Lloyd, S., Mackay, P., McGoldrick, M., Mumford, J., Pye, B., Riddell, K., Rodrigue, M., Strudwick, E., & Supple, R. (2003). Victoria's System of Marine National Parks and Marine Sanctuaries: Management Strategy 2003 - 2010. In: P. Victoria. Melbourne: Parks Victoria.

Ferns, L. W. (1999). *Environmental inventory of Victoria's marine ecosystems Stage 4 (Part 1)*. East Melbourne, Australia: Parks, Flora and Fauna Division, Department of Natural Resources and Environment.

Folk, R. L. (1974). *Petrology of Sedimentary Rocks*. Austin, TX.: Hemphills.

Folt, C. L., Chen, C. Y., Moore, M. V., & Burnaford, J. (1999). Synergism and antagonism among multiple stressors. *Limnology and Oceanography*, 44, 864-877.

Fortin, M. J., Keitt, T. H., Maurer, B. A., Taper, M. L., Kaufman, D. M., & Blackburn, T. M. (2005). Species' geographic ranges and distributional limits: pattern analysis and statistical issues. *Oikos*, 108(1), 7-17.

Foster, M. S. (2001). Rhodoliths: Between rocks and soft places. *Journal of Phycology*, 37(5), 659-667.

Franklin, J. (1995). Predictive vegetation mapping: Geographic modelling of biospatial patterns in relation to environmental gradients. *Progress in Physical Geography*, 19(4), 474-499.

Fraser, R. H., & Currie, D. J. (1996). The species richness-energy hypothesis in a system where historical factors are thought to prevail: Coral reefs. *American Naturalist*, 148(1), 138-159.

Freeman, S. M., & Rogers, S. I. (2003). A new analytical approach to the characterisation of macro-epibenthic habitats: linking species to the environment. *Estuarine Coastal and Shelf Science*, 56(3-4), 749-764.

Fry, B. (2006). *Stable Isotope Ecology*. New York: Springer.

Gage, J.D. and Tyler, P.A. (1991). *Deep-sea Biology: A Natural History of Organisms at the Deep-Sea Floor*. Cambridge University Press, Cambridge.

Gage, J. D. (1996). Why are there so many species in deep-sea sediments? *Journal of Experimental Marine Biology and Ecology*, 200(1-2), 257-286.

Gage, J. D. (2004). Diversity in deep-sea benthic macrofauna: the importance of local ecology, the larger scale, history and the Antarctic. *Deep-Sea Research Part II*, 51(14-16), 1689-1708.

Galeron, J., Sibuet, M., Vanreusel, A., Mackenzie, K., Gooday, A. J., Dinet, A., & Wolff, G. A. (2001). Temporal patterns among meiofauna and macrofauna taxa related to changes in sediment geochemistry at an abyssal NE Atlantic site. *Progress in Oceanography*, 50(1-4), 303-324.

Galil, B. S. (2000). A sea under siege -alien species in the Mediterranean. *Biological Invasions*, 2, 177-186.

- Galy, V., Bouchez, J., & France-Lanord, C. (2007). Determination of total organic carbon content and delta C-13 in carbonate-rich detrital sediments. *Geostandards and Geoanalytical Research*, 31(3), 199-207.
- Gambi, C., & Danovaro, R. (2006). A multiple-scale analysis of metazoan meiofaunal distribution in the deep Mediterranean Sea. *Deep-Sea Research Part I*, 53(7), 1117-1134.
- Gardner, R. H., & Engelhardt, K. A. M. (2008). Spatial processes that maintain biodiversity in plant communities. *Perspectives in Plant Ecology Evolution and Systematics*, 9(3-4), 211-228.
- Gladstone, W. (2007). Requirements for marine protected areas to conserve the biodiversity of rocky reef fishes. *Aquatic Conservation-Marine and Freshwater Ecosystems*, 17(1), 71-87.
- Gogina, M., Glockzin, M., Zettler, M.L. (2010). Distribution of benthic macrofaunal communities in the western Baltic Sea with regard to near-bottom environmental parameters. 1. Causal analysis. *Journal of Marine Systems*, 79, 112-123.
- Gooday, A. J. (2002). Biological Responses to Seasonally Varying Fluxes of Organic Matter to the Ocean Floor: A Review *Journal of Oceanography*, 58, 305-332.
- Grassle, J. F., & Maciolek, N. J. (1992). Deep sea species richness - regional and local diversity estimates from quantitative bottom samples. *American Naturalist*, 139(2), 313-341.
- Gray, J. S. (1974). Animal-sediment relationships. *Oceanography and Marine Biology Annual Review*, 12, 223-261.
- Gray, J. S. (1997). Marine biodiversity: patterns, threats and conservation needs. *Biodiversity and Conservation*, 6, 153-175.
- Gray, J. S. (2000). The measurement of marine species diversity, with an application to the benthic fauna of the Norwegian continental shelf. *Journal of Experimental Marine Biology and Ecology*, 250, 23-49.
- Gray, J. S. (2001). Marine diversity: the paradigms in patterns of species richness examined. *Scientia Marina*, 65, 41-56.
- Gray, J. S. (2002). Species richness of marine soft sediments. *Marine Ecology-Progress Series*, 244, 285-297.
- Green, M. O., Hewitt, J. E., & Thrush, S. F. (1998). Seabed drag coefficient over natural beds of horse mussels (*Atrina zelandica*). *Journal of Marine Research*, 56, 613-637.
- Greene, H. G., O'Connell, V. M., Wakefield, W. W., & Byrlesky, C. K. (2007a). The offshore Edgecumbe lava field, southeast Alaska: geologic and habitat characterization of a commercial fishing ground. In: B. J. Todd, & H. G. Greene, *Mapping the Seafloor for Habitat Characterization*, Vol. 47 (pp. 277-295). St Johns, Canada: Geological Association of Canada.
- Greene, H.G., Vallier, T.L., Bizzarro, J.J., Watt, S., Dieter, B.E. (2007b). Impacts of bay floor disturbances on benthic habitats in San Francisco Bay. In: Todd, B.J., Greene, H.G., *Mapping the Seafloor for Habitat Characterization*, Vol 47 (p 401-419). St Johns, Canada: Geological Association of Canada.

- Greene, H.G., Bizzarro, J.J., O'Connell, V.M., Brylinsky, C.K. (2007c). Construction of digital potential marine benthic habitat maps using a coded classification scheme and its application. In: Todd, B.J., Greene, H.G., *Mapping the Seafloor for Habitat Characterization*, Vol 47 (p 141-155). St Johns, Canada: Geological Association of Canada.
- Greene, H. G., Yoklavich, M. M., Starr, R. M., O'Connell, V. M., Wakefield, W. W., Sullivan, D. E., McRea, J. E., & Cailliet, G. M. (1999). A classification scheme for deep seafloor habitats. *Oceanologica Acta*, 22(6), 663-678.
- Greene, H. G., Yoklavich, M. M., Sullivan, D., & Cailliet, G. M. (1995). A Geophysical Approach to Classifying Marine Benthic Habitats: Monterey Bay as a Model. In: T. O'Connell, & W. Wakefield, *Applications of Side Scan Sonar and Laser-Line Systems in Fisheries Research*, vol. 9 (pp. 15-30): Alaska Fish and Game.
- Grove, S. L., Probert, P. K., Berkenbusch, K., & Nodder, S. D. (2006). Distribution of bathyal meiofauna in the region of the Subtropical Front, Chatham Rise, south-west Pacific. *Journal of Experimental Marine Biology and Ecology*, 330(1), 342-355.
- Gustafson, E. J. (1998). Quantifying landscape spatial pattern: What is the state of the art? *Ecosystems*, 1(2), 143-156.
- Gutt, J., & Piepenburg, D. (2003). Scale-dependent impact on diversity of Antarctic benthos caused by grounding of icebergs. *Marine Ecology-Progress Series*, 253, 77-83.
- Hagberg, J., Jonzen, N., Lundberg, P., & Ripa, J. (2003). Uncertain biotic and abiotic interactions in benthic communities. *Oikos*, 100(2), 353-361.
- Hall, S.J.(1994). Physical disturbance and marine benthic communities: life in unconsolidated sediments. *Oceanography and Marine Biology Annual Review*, 32, 179–239.
- Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Argosa, C., Bruno, J. F., Casey, K. S., Ebert, C., Fox, H. E., Fujita, R., Heinemann, D., Lenihan, H. S., Madin, E. M. P., Perry, M. T., Selig, E. R., Spalding, M., Steneck, R., & Watson, R. (2008). A global map of human impact on marine ecosystems. *Science*, 319, 948-952.
- Harris, J., Haward, M., Jabour, J., & Woehler, E. J. (2007a). A new approach to selecting Marine Protected Areas (MPAs) in the Southern Ocean. *Antarctic Science*, 19(2), 189-194.
- Harris, P., Heap, A., Whiteway, T., & Post, A. (2007b). Applications of Geoscience to Australia's Representative Marine Protected Area Program., *Alaska Habitat Mapping Workshop*. Anchorage: Geoscience Australia.
- Harris, P. T., Heap, A. D., Whiteway, T., & Post, A. (2008). Application of biophysical information to support Australia's representative marine protected area program. *Ocean & Coastal Management*, 51, 701-711.
- Harris, P. T., & Whiteway, T. (2009). High seas marine protected areas: Benthic environmental conservation priorities from a GIS analysis of global ocean biophysical data. *Ocean & Coastal Management*, 52(1), 22-38.
- Hart, D. E., & Kench, P. S. (2007). Carbonate production of an emergent reef platform, Warraber Island, Torres Strait, Australia. *Coral Reefs*, 26(1), 53-68.

- Hawkins, B. A. (2001). Ecology's oldest pattern? *Endeavour*, 25(3), 133-134.
- Hawkins, B. A., Porter, E. E., & Diniz, J. A. F. (2003). Productivity and history as predictors of the latitudinal diversity gradient of terrestrial birds. *Ecology*, 84(6), 1608-1623.
- Hayden, B. P., Ray, G. C., & Dolan, R. (1984). Classification of coastal and marine environments. *Environmental Conservation*, 11, 199-207.
- He, F., & Legendre, P. (2002). Species diversity patterns derived from species-area models. *Ecology*, 83(5), 1185-1198.
- Heap, A. D., Harris, P. T., Hinde, A., & Woods, M. (2005). *National Marine Bioregionalisation of Australia's Exclusive Economic Zone*. Canberra, Geoscience Australia.
- Hedgpeth, J. W. (1957). Marine Biogeography. *Geological Society of America Memoirs*, 67, 17-28.
- Hemer, M. A. (2006). The magnitude and frequency of combined flow bed shear stress as a measure of exposure on the Australian continental shelf. *Continental Shelf Research*, 26(11), 1258-1280.
- Hessler, R.R. (1974). The structure of deep benthic communities from central oceanic waters. In: *The Biology of the Ocean Pacific*, C.B. Miller (ed), pp. 79-94. Oregon state University Press, Corvallis.
- Hillebrand, H., Gruner, D. S., Borer, E. T., Bracken, M. E. S., Cleland, E. E., Elser, J. J., Harpole, W. S., Ngai, J. T., Seabloom, E. W., Shurin, J. B., & Smith, J. E. (2007). Consumer versus resource control of producer diversity depends on ecosystem type and producer community structure. *Proceedings of the National Academy of Sciences of the United States of America*, 104(26), 10904-10909.
- Hinojosa-Arango, G., & Riosmena-Rodriguez, R. (2004). Influence of rhodolith-forming species and growth-form on associated fauna of rhodolith beds in the central-west Gulf of California, Mexico. *Marine Ecology-Pubblicazioni Della Stazione Zoologica Di Napoli I*, 25(2), 109-127.
- Hogue, E. W., & Miller, C. B. (1981). Effects of sediment micro-topography on small-scale spatial distributions of meiobenthic nematodes. *Journal of Experimental Marine Biology and Ecology*, 53(2-3), 181-191.
- Holmes, K. W., Van Niel, K. P., Radford, B., Kendrick, G. A., & Grove, S. L. (2008). Modelling distribution of marine benthos from hydroacoustics and underwater video. *Continental Shelf Research*, 28, 1800-1810.
- Hooper, J. N. A., Kennedy, J. A., & Quinn, R. J. (2002). Biodiversity 'hotspots', patterns of richness and endemism, and taxonomic affinities of tropical Australian sponges (Porifera). *Biodiversity and Conservation*, 11(5), 851-885.
- Hourston, M., Warwick, R. M., Valesini, F. J., & Potter, I. C. (2005). To what extent are the characteristics of nematode assemblages in nearshore sediments on the west Australian coast related to habitat type, season and zone? *Estuarine Coastal and Shelf Science*, 64(4), 601-612.

- Howard-Williams, C., Peterson, D., Lyons, W. B., Cattaneo-Vietti, R., & Gordon, S. (2006). Measuring ecosystem response in a rapidly changing environment: the Latitudinal Gradient Project. *Antarctic Science*, 18(4), 465-471.
- Hunt, B., & Vincent, A. C. J. (2006). Scale and sustainability of marine bioprospecting for pharmaceuticals. *Ambio*, 35(2), 57-64.
- Hutchinson, G. E. (1957). Concluding Remarks. *Cold Spring harbor Symposia on Quantitative Biology*, 22, 415-427.
- IMCRA (2006). *Interim marine and coastal regionalisation for Australia: an ecosystem-based classification for marine and coastal environments. Version 4.0*. Canberra: Department of the Environment and Heritage, Commonwealth of Australia.
- Inoue, T., Suda, Y., & Sano, M. (2008). Surf zone fishes in an exposed sandy beach at Sanrimatsubara, Japan: does fish assemblage structure differ among microhabitats? *Estuarine and Coastal Shelf Science*, 77, 1-11.
- Jackson, J. B. C. (2008). Ecological extinction and evolution in the brave new ocean. *Proceedings of the National Academy of Sciences*, 105, 11458-11465.
- Jenkins, G., & Wheatley, M. (1998). The influence of habitat structure on nearshore fish assemblages in a southern Australian embayment: Comparison of shallow seagrass, reef-algal and unvegetated sand habitats, with emphasis on their importance to recruitment. *Journal of Experimental Marine Biology and Ecology*, 221, 147-172.
- Jensen, A., & Frederickson, R. (1992). The fauna associated with the bank-forming deepwater coral *Lophelia pertusa* (Scleractinaria) on the Faroe shelf. *Sarsia*, 77, 53-69.
- Jie, H., Zhinan, Z., Zishan, Y., & Widdows, J. (2001). Differences in the benthic-pelagic particle flux (biodeposition and sediment erosion) at intertidal sites with and without clam (*Ruditapes philippinarum*) cultivation in eastern China. *Journal of Experimental Marine Biology and Ecology*, 261(2), 245-261.
- Jones, G., & Syms, C. (1998). Disturbance, habitat structure and the ecology of fishes on coral reefs. *Australian Journal of Ecology*, 23, 287-297.
- Jones, N. S. (1950). Marine bottom communities. *Biological Reviews of the Cambridge Philosophical Society*, 25(3), 283-313.
- Kadmon, R., & Allouche, O. (2007). Integrating the effects of area, isolation, and habitat heterogeneity on species diversity: A unification of island biogeography and niche theory. *American Naturalist*, 170(3), 443-454.
- Kalmar, A., & Currie, D. J. (2006). A global model of island biogeography. *Global Ecology and Biogeography*, 15(1), 72-81.
- Karadzic, B., Marinkovic, S., & Katarinovski, D. (2003). Use of the beta-function to estimate the skewness of species responses. *Journal of Vegetation Science*, 14(6), 799-805.
- Ke, X., Collins, M. B., & Poulos, S. E. (1994). Velocity structure and sea bed roughness associated with intertidal (sand and mud) flats and saltmarshes of the wash. *UK Journal of Coastal Research*, 10, 702-715.

- Kelleher, G., Bleakley, C., & Wells, S. (1995). *A Global Representative System of Marine Protected Areas*. Washington D.C., The Great Barrier Reef Marine Park Authority.
- Kendall, M., Jensen, O., Alexander, C., Field, D., McFall, G., Bohne, R., & Monaco, M. (2005). Benthic mapping using sonar, video transects and an innovative approach to accuracy assessment: a characterization of bottom features in the Georgia Bights. *Journal of Coastal Research*, *21*, 1154-1165.
- Kennard, M. J., Harch, B. D., Pusey, B. J., & Arthington, A. H. (2006). Accurately defining the reference condition for summary biotic metrics: A comparison of four approaches. *Hydrobiologia*, *572*, 151-170.
- Kennet, J. P. (1982). *Marine Geology*. Englewood Cliffs NJ: Prentice Hall Inc.
- Kerswell, A. P. (2006). Global biodiversity patterns of benthic marine algae. *Ecology*, *87*(10), 2479-2488.
- Klitgaard, A. B. (1995). The fauna associated with outer shelf and upper slope sponges (Porifera, Demosptongiae) at the Faroe Islands, northeastern Atlantic. *Sarsia*, *80*, 1-22.
- Kloser, R. J., Williams, A., & Butler, A. J. (2007). Exploratory surveys of seabed habitats in Australia's deep ocean using remote sensing - needs and realities. In: B. J. Todd, & H. G. Greene, *Mapping the Seafloor for Habitat Characterization*, vol. 47 (pp. 93-110). St Johns, Canada: Geological Society of Canada.
- Knight, T. W., & Morris, D. W. (1996). How many habitats do landscapes contain? *Ecology*, *77*, 1756-1764.
- Koppelman, R., & Weikert, H. (2007). Spatial and temporal distribution patterns of deep-sea mesozooplankton in the eastern Mediterranean - indications of a climatically induced shift? *Marine Ecology-an Evolutionary Perspective*, *28*(2), 259-275.
- Kostylev, V. E., Courtney, R. C., Robert, G., & Todd, B. J. (2003). Stock evaluation of giant scallop (*Placopecten magellanicus*) using high-resolution acoustics for seabed mapping. *Fisheries Research*, *60*(2-3), 479-492.
- Kostylev, V. E., Todd, B. J., Fader, G. B. J., Courtney, R. C., Cameron, G. D. M., & Pickrill, R. A. (2001). Benthic habitat mapping on the Scotian Shelf based on multibeam bathymetry, surficial geology and sea floor photographs. *Marine Ecology-Progress Series*, *219*, 121-137.
- Kukowski, N., Harnpel, A., Hoth, S., & Bialas, J. (2008). Morphotectonic and morphometric analysis of the Nazca plate and the adjacent offshore Peruvian continental slope - Implications for submarine landscape evolution. *Marine Geology*, *254*(1-2), 107-120.
- Lampitt, R., Antia, R. (1997). Particle flux in the deep seas: regional characteristics and temporal variability. *Deep-Sea Research I*, *44*: 1377-1404
- Lanier, A., Romsos, C., & Goldfinger, C. (2007). Seafloor habitat mapping on the Oregon continental margin: A spatially nested GIS approach to mapping scale, mapping methods, and accuracy quantification. *Marine Geodesy*, *30*(1-2), 51-76.

- Larsen, F. W., & Rahbek, C. (2005). The influence of spatial grain size on the suitability of the higher-taxon approach in continental priority-setting. *Animal Conservation*, 8, 389-396.
- Legendre, P., & Legendre, L. (1998). *Numerical Ecology*. Amsterdam: Elsevier Science.
- Leigh Jr, E. G. (2007). Neutral theory: a historical perspective. *Journal of Evolutionary Biology*, 20(6), 2075-2091.
- Lenihan, H. S. (1999). Physical-biological coupling on oyster reefs: How habitat structure influences individual performance. *Ecological Monographs*, 69(6), 251-275.
- Lenihan, H. S., & Micheli (2001). Soft-Sediment Communities. In: M. D. Bertness, S. D. Gaines, & M. E. Hay, *Marine Community Ecology* (p. 550). Sunderland MA: Sinauer.
- Levin, L. A., Etter, R. J., Rex, M. A., Gooday, A. J., Smith, C. R., Pineda, J., Stuart, C. T., Hessler, R. R., & Pawson, D. (2001). Environmental influences on regional deep-sea species diversity. *Annual Review of Ecology and Systematics*, 32, 51-93.
- Levin, L. A., & Gage, J. D. (1998). Relationships between oxygen, organic matter and the diversity of bathyal macrofauna. *Deep Sea Research. II*, 45(1/3), 129.
- Levin, S. A. (1992). The Problem of Pattern and Scale in Ecology. *Ecology*, 73(6), 1943-1967.
- Lewis, A., Stein, J. L., Stein, J. A., Nix, H. A., Mackey, B. G., & Bowyer, J. K. (1991). *An assessment of the regional conservation adequacy of Tasmania*. Canberra, Resource Assessment Commission.
- Lewis, D. W., & McConchie, D. (1994). *Analytical Sedimentology*. New York: Chapman and Hall.
- Li, H., & Reynolds, J. F. (1993). A new contagion index to quantify spatial patterns in landscape. *Landscape Ecology*, 8, 155-162.
- Lillesand, T. M., & Kiefer, R. W. (1994). *Remote Sensing and Image Interpretation*. John Wiley and Sons. USA.
- Lindholm, J. B., Auster, P. J., Ruth, M., & Kaufman, L. (2001). Modelling the effects of fishing and implications for the design of marine protected areas: juvenile fish responses to variations in seafloor habitat. *Conservation Biology*, 15, 424-437.
- Lomolino, M. V., & Weiser, M. D. (2001). Towards a more general species-area relationship: diversity on all islands, great and small. *Journal of Biogeography*, 28(4), 431-445.
- Longhurst, A. R. (1998). *Ecological Geography of the Sea*. San Diego: Academic Press.
- Lyne, V., & Hayes, D. (2005). *Pelagic regionalisation: national marine bioregionalisation project*. Hobart, National Oceans Office, CSIRO Marine Research.
- MacArthur, R. H., & Wilson, E. O. (1963). An equilibrium theory of insular zoogeography. *Evolution*, 17(4), 373.

- Mackey, B. G., Nix, H. A., Hutchinson, M. F., McMahan, J. P., & Fleming, P. M. (1988). Assessing representativeness of places for conservation reservation and heritage listing. *Environmental Management*, *12*(4), 501-514.
- Mackey, B. G., Nix, H. A., Stein, J. A., Cork, S. E., & Bullten, F. T. (1989). Assessing the representativeness of the wet tropics of Queensland World Heritage property. *Biological Conservation*, *50*, 239-261.
- Magierowski, R. H., & Johnson, C. R. (2006). Robustness of surrogates of biodiversity in marine benthic communities. *Ecological Applications*, *16*(6), 2264-2275.
- Magurran, A. E. (2004). *Measuring Biological Diversity*. Blackwell Publishing.
- May, R. M. (1992). Biodiversity - bottoms up for the oceans. *Nature*, *357*(6376), 278-279.
- Mayr, E. (1996). What is a species and what is not? *Philosophy of Science*, *63*, 262-277.
- McGill, B. J. (2003). A test of the unified neutral theory of biodiversity. *Nature*, *422*, 881-885.
- McGuinness, K. A. (1984). Species-area relations of communities on intertidal boulders: testing the null hypothesis. *Journal of Biogeography*, *11*, 439-456.
- Menge, B. A., Daley, B. A., Lubchenco, J., Sanford, E., Dahlhoff, E., Halpin, P. M., Hudson, G., & Burnaford, J. L. (1999). Top-down and bottom-up regulation of New Zealand rocky intertidal communities. *Ecological Monographs*, *69*(3), 297-330.
- Meynard, C. N., & Quinn, J. F. (2007). Predicting species distributions: a critical comparison of the most common statistical models using artificial species. *Journal of Biogeography*, *34*(8), 1455-1469.
- Micheli, F., Peterson, C. H., Mullineaux, L. S., Fisher, C. R., Mills, S. W., Sancho, G., Johnson, G. A., & Lenihan, H. S. (2002). Predation structures communities at deep-sea hydrothermal vents. *Ecological Monographs*, *72*(3), 365-382.
- Mittelbach, G. G., Schemske, D. W., Cornell, H. V., Allen, A. P., Brown, J. M., Bush, M. B., Harrison, S. P., Hurlbert, A. H., Knowlton, N., Lessios, H. A., McCain, C. M., McCune, A. R., McDade, L. A., McPeck, M. A., Near, T. J., Price, T. D., Ricklefs, R. E., Roy, K., Sax, D. F., Schluter, D., Sobel, J. M., & Turelli, M. (2007). Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecology Letters*, *10*(4), 315-331.
- Moore, D. M., Lees, B. G., & Davey, S. M. (1991). A new method for predicting vegetation distributions using decision tree analysis in a geographic information system. *Environmental Management*, *15*(1), 59-71.
- Mossop, J. (2004). Legal challenges to the establishment of high-seas marine protected areas. In: *4th World Fisheries Congress* (pp. 1503-1515). Vancouver, CANADA: Amer Fisheries Soc.
- Motta, J. J. C., Underwood, A. J., Chapman, M. G., & Rossi, F. (2003). Benthic assemblages in sediments associated with intertidal boulder-fields. *Journal of Experimental Marine Biology and Ecology*, *285*, 383-401.

- Mumby, P. J., Green, E. P., Edwards, A. J., & Clark, C. D. (1997). Measurement of seagrass standing crop using satellite and digital airborne remote sensing. *Marine Ecology Progress Series*, 159(6), 51-60.
- Mumby, P. J., & Harborne, A. R. (1999). Development of a systematic classification scheme of marine habitats to facilitate regional management and mapping of Caribbean coral reefs. *Biological Conservation*, 88, 155-163.
- Murphy, D. D., & Duffus, D. A. (1996). Conservation biology and marine biodiversity. *Conservation Biology*, 10(2), 311-312.
- Murray, J. M. H., Meadows, A., & Meadows, P. S. (2002). Biogeomorphological implications of microscale interactions between sediment geotechnics and marine benthos: a review. *Geomorphology*, 47, 15-30.
- Nakamura, Y., & Sano, M. (2005). Comparison of invertebrate abundance in a seagrass bed and adjacent coral and sand areas at Amitori Bay, Iriomote Island, Japan. *Fisheries Science*, 71(3), 543-550.
- Neira, C., & Hoepner, T. (1994). The role of *Heteromastus filiformis* (Capitellidae, Polychaeta) in organic carbon cycling. *Ophelia*, 39(1), 55-73.
- Newell, R. C., Seiderer, L. J., & Hitchcock, D. R. (1998). The impact of dredging works in coastal waters: a review of the sensitivity to disturbance and subsequent recovery of biological resources on the sea bed. *Oceanography and Marine Biology: an Annual Review*, 36, 127-178.
- Nicolaidou, A., & Papadopoulou, K. N. (1989). Factors affecting the distribution and diversity of polychaetes in Amvrakikos Bay, Greece. *Marine Ecology (Pubblicazioni Della Stazione Zoologica Di Napoli I)*, 10(3), 193-204.
- Norling, K., Rosenberg, R., Hulth, S., Gremare, A., & Bonsdorff, E. (2007). Importance of functional biodiversity and species-specific traits of benthic fauna for ecosystem functions in marine sediment. *Marine Ecology-Progress Series*, 332, 11-23.
- O'Hara, T. (2000). Habitats as surrogates for faunal and floral assemblages associated with rocky reefs along the Victorian coast. In: L. W. Ferns, & D. Hough, *Environmental Inventory of Victoria's Marine Ecosystems, Stage 3 (2nd edition) - Understanding Biodiversity Representativeness of Victoria's Rocky Reefs*. East Melbourne: Department of Natural Resources and Environment
- O'Hara, T. D., & Poore, G. C. B. (2000). Patterns of distribution for southern Australian marine echinoderms and decapods. *Journal of Biogeography*, 27, 1321-1335.
- O'Neill, R. V., Krummel, R. H., Gardner, R. H., Sugihara, G., Jackson, B., deAngelis, D. I., Milne, B. T., Turner, M. G., Zygmunt, B., Christensen, S. W., Dale, V. H., & Graham, R. L. (1998). Indices of landscape pattern. *Landscape Ecology*, 16, 153-163.
- Olabarria, C. (2006). Faunal change and bathymetric diversity gradient in deep-sea prosobranchs from Northeastern Atlantic. *Biodiversity and Conservation*, 15(11), 3685-3702.

- Paavola, R., Muotka, T., Virtanen, R., Heino, J., Jackson, D., & Maki-Petays, A. (2006). Spatial scale affects community concordance among fishes, benthic macroinvertebrates, and bryophytes in streams. *Ecological Applications*, *16*(1), 368-379.
- Palumbi, S. R. (2001). The Ecology of Marine Protected Areas. In: M. D. Bertness, S. D. Gaines, & M. E. Hay, *Marine Community Ecology* (pp. 509-532). Sunderland MA: Sinauer Associates, Inc.
- Parmar, R. M., Arora, R. K., Rao, M. V., & Thyagarajan, K. (2006). OCEANSAT-2 mission and its applications - art. no. 64070C. In: *Conference on GEOSS and Next-Generation Sensors and Missions* (pp. C4070-C4070). Goa, INDIA: Spie-Int Soc Optical Engineering.
- Parry, D. M., Kendall, M. A., Pilgrim, D. A., & Jones, M. B. (2003). Identification of patch structure within marine benthic landscapes using a remotely operated vehicle. *Journal of Experimental Marine Biology and Ecology*, *285*, 497-511.
- Passlow, V., O'Hara, T. D., Daniell, T., Beaman, R. J., & Twyford, L. M. (2006). *Sediments and Benthic Biota of Bass Strait: an Approach to Benthic Habitat Mapping*. Canberra, Geoscience Australia. 2004/23.
- Paterson, G. L. J., Lambshead, P. J. D., & Gage, J. D. (1992). Down the slippery slope: A study of polychaete assemblages from bathymetric transects in the Rockall Trough. *Polychaete Research Newsletter*, No., *14*, 2.
- Patzner, R. A. (1999). Habitat utilization and depth distribution of small cryptobenthic fishes (Blenniidae, Gobiesocidae, Gobiidae, Tripterygiidae) in Ibiza (Western Mediterranean Sea). *Environmental Biology of Fishes*, *55*, 207-214.
- Pauly, D., Watson, R., & Alder, J. (2005). Global trends in world fisheries: impacts on marine ecosystems and food security. *Philosophical Transactions of the Royal Society B-Biological Sciences*, *B 360*, 5-12.
- Pearson, T. H. (2001). Functional group ecology in soft sediment marine benthos: The role of bioturbation. *Oceanography and Marine Biology, An Annual Review*, *39*, 233-267.
- Petch, D. A. (1986). Selective deposit-feeding by *Lumbrineris* cf. *latreilli* (Polychaeta: Lumbrineridae), with a new method for assessing selectivity by deposit-feeding organisms. *Marine Biology* *93*, 443-448.
- Peterson, G., Allen, C. R., & Holling, C. S. (1998). Ecological resilience, biodiversity and scale. *Ecosystems*, *1*, 6-18.
- Piepenburg, D., Schmid, M. K., & Gerdes, D. (2002). The benthos off King George Island (South Shetland Islands, Antarctica): further evidence for a lack of a latitudinal biomass cline in the Southern Ocean. *Polar Biology*, *25*(2), 146-158.
- Pitcher, C. R., Doherty, P., Arnold, P., Hooper, J., Gribble, N., Bartlett, C., Browne, M., Campbell, N., Cannard, T., Cappel, M., Carini, G., Chalmers, S., Cheers, S., Chetwynd, D., Colefax, A., Coles, R., Cook, S., Davie, P., De'ath, G., Devereux, D., Done, B., Donovan, T., Ehrke, B., Ellis, N., Ericson, G., Jacobsen, I., Johnson, J., Jones, M., Kinninmoth, S., Kistle, S., Last, P., Leite, A., Marks, S., McLeod, I., Oczkowicz, S., Robinson, M., Rose, C., Seabright, D., Sheils, J., Sherlock, M., Skelton, P., Smith, D., Smith, G., Speare, P., Stowar, M., Strickland, C., Van der Geest, C., Venables, W., Walsh, C., Wassenberg, T., Welna, A., & Yearsley, G. (2007).

- Seabed Biodiversity on the Continental Shelf of the Great Barrier Reef World Heritage Area.*, AIMS/CSIRO/QM/QDPI CRC Reef Research Task Final Report.
- Ponder, W., Hutchings, P., & Champman, R. (2002). *Overview of the Conservation of Australian Marine Invertebrates: A Report for Environment Australia*. Sydney, Australian Museum.
- Poore, G. C. B., Just, J., & Cohen, B. (1994). Composition and diversity of Crustacea Isopoda of the southeastern Australian continental slope. *Deep-Sea Research*, 41(4), 677-693.
- Poore, G. C. B., McCallum, A. W., & Taylor, J. (2008). *Decapod Crustacea of the continental margin of southwestern Australia and central Western Australia: preliminary identifications of 524 species from FRV Southern Surveyor voyage SS10-2005*. Melbourne, Museum Victoria. 11.
- Poore, G. C. B., & O'Hara, T. D. (2007). Marine biogeography and biodiversity of Australia. In: S. D. Connell, & B. M. Gillanders, *Marine Ecology* (pp. 177-198). Melbourne: Oxford University Press.
- Post, A. L., Wassenberg, T. J., & Passlow, V. (2006). Physical surrogates for macrofaunal distributions and abundance in a tropical gulf. *Marine and Freshwater Research*, 57(5), 469-483.
- Prada, M. C., Appeldoorn, R. S., & Rivera, J. A. (2008). Improving coral reef habitat mapping of the Puerto Rico insular shelf using side scan sonar. *Marine Geodesy*, 31, 49-73.
- Przeslawski, R., Davis, A.R. (2007). Does spawning behaviour minimize exposure to environmental stressors for encapsulated gastropod embryos on rocky shores? *Marine Biology*, 152: 991-1002.
- Rees, S. A., Opdyke, B. N., Wilson, P. A., & Henstock, T. J. (2007). Significance of Halimeda bioherms to the global carbonate budget based on a geological sediment budget for the Northern Great Barrier Reef, Australia. *Coral Reefs*, 26(1), 177-188.
- Reineck, H. E., & Singh, I. B. (1980). *Depositional Sedimentary Environments: with Reference to Terrigenous Clastics*. Berlin: Springer - Verlag.
- Reise, K. (1981). *Gnathostomulida* abundant alongside polychaete burrows. *Marine Ecology Progress Series*, 6(3), 329-333.
- Rex, M. A., Crame, J. A., Stuart, C. T., & Clarke, A. (2005). Large-scale biogeographic patterns in marine mollusks: A confluence of history and productivity? *Ecology*, 86(9), 2288-2297.
- Rex, M. A., Etter, R. J., Morris, J. S., Crouse, J., McClain, C. R., Johnson, N. A., Stuart, C. T., Deming, J. W., Thies, R., & Avery, R. (2006). Global bathymetric patterns of standing stock and body size in the deep-sea benthos. *Marine Ecology-Progress Series*, 317, 1-8.
- Rhoads, D. C. (1974). Organism-sediment relationships on the muddy sea floor. *Oceanography and Marine Biology Annual Review*, 12, 263-300.
- Riitters, K. H. , O'Neil, R. V., Hunsaker, C. T., Wickham, J. D., Yankee, D. H., Timmins, S. P., Jones, K. B., & Jackson, J. B. (1995). A factor analysis of landscape pattern and structure metric. *Landscape Ecology*, 10, 23-39.
- Romsos, C. G., Goldfinger, C., Robison, R., Milstein, R. L., & Chaytor, J. D. (2007). Development of a regional seafloor surficial geologic habitat map for the continental margins of

- Oregon and Washington, USA. In: B. J. Todd, & H. G. Greene, *Mapping the Seafloor for Habitat Characterization.*, vol. Special Paper 47 (pp. 219-243): Geological Association of Canada.
- Ross, P. M., Thrush, S. F., Montgomery, J. C., Walker, J. W., & Parsons, D. M. (2007). Habitat complexity and predation risk determine juvenile snapper (*Pagrus auratus*) and goatfish (*Upeneichthys lineatus*) behaviour and distribution. *Marine and Freshwater Research*, 58(12), 1144-1151.
- Rowden, A. A., Clark, M. R., & Wright, I. C. (2005). Physical characterisation and a biologically focused classification of "seamounts" in the New Zealand region. *New Zealand Journal of Marine and Freshwater Research*, 39(5), 1039-1059.
- Rowe, G.T., Sibuet, M.S., Deming, J., Kripounoff, A., Tietjen, J., Macko, S., Theroux, R. (1991). "Total sediment biomass and preliminary estimates of organic carbon in deep-sea sediments. *Journal of Marine Research*, 43: 925-950.
- Ryan, D. A., Brooke, B. P., Bostock, H. C., Radke, L. C., Siwabessy, P. J. W., Margvelashvili, N., & Skene, D. (2007a). Bedload sediment transport dynamics in a macrotidal embayment, and implications for export to the southern Great Barrier Reef shelf. *Marine Geology*, 240(1-4), 197-215.
- Ryan, D. A., Brooke, B. P., Collins, L. B., Kendrick, G. A., Baxter, K. J., Bickers, A. N., Siwabessy, P. J. W., & Pattiaratchi, C. B. (2007b). The influence of geomorphology and sedimentary processes on shallow water benthic habitat distribution: Esperance Bay, Western Australia. *Estuarine and Coastal Shelf Science*, 72(379-386).
- Sanders, H. L. (1958). Benthic Studies in Buzzards Bay. I. Animal-Sediment Relationships. *Limnology and Oceanography*, 3(3), 245.
- Sanders, J. L., Kendall, M. A., Hawkins, A. J. S., & Spicer, J. I. (2007). Can functional groups be used to indicate estuarine ecological status? *Hydrobiologia*, 588, 45-58.
- Sarkar, S., Justus, J., Fuller, T., Kelley, C., Garson, J., & Mayfield, M. (2005). Effectiveness of environmental surrogates for the selection of conservation area networks. *Conservation Biology*, 19(3), 815-825.
- Scanes, P. R., Mann, R. A., Manning, T. M., & Chapman, J. C. (1993). Observations on the biota of an estuarine beach at Hardys Bay (NSW, Australia) following a spillage of aldrin. *Marine Pollution Bulletin*, 26(12), 687-691.
- Schemske, D. (2002). Tropical diversity: patterns and processes. In: R. Chazdon, & T. Whitmore, *Ecological and Evolutionary Perspectives on the Origins of Tropical Diversity: Key Papers and Commentaries*. Chicago IL: University of Chicago Press.
- Schubert, C.J., Niggerman, J., Klockgether, G., and Ferdelman, G., 2005. Chlorin Index: A new parameter for organic matter freshness in sediment. *Geochemistry Geophysics and Geosystems*, 6(3), 1-12.
- Sebesvari, Z., F. Esser, & Harder, T. (2006). Sediment-associated cues for larval settlement of the infaunal spionid polychaetes *Polydora cornuta* and *Streblospio benedicti*. *Journal of Experimental Marine Biology and Ecology* 337, 109-120.

- Seiter, K., Hensen, C., Schroter, J., Zabel, M., 2004. Organic carbon content in surface sediments – defining regional provinces. *Deep-Sea Research I*, 51, 2001-2026
- Seiter, K., Hensen, C., Zabel, M., 2005. Benthic carbon mineralisation on a global scale. *Global Biogeochemical Cycles*, 19, 1-26.
- Sherman, K., & Alexander, L. M. (1989). *Biomass Yields and Geography of Large Marine Ecosystems*. Boulder: Westview Press.
- Shulman, M. J., & Bermingham, E. (1995). Early-life histories, ocean currents, and the population-genetics of Caribbean reef fishes. *Evolution*, 49(5), 897-910.
- Smith, C.R., Berelson, W., DeMaster, D.J., Dobbs, F.C., Hammond, D., Hoover, D.J., Popoe, R.H. and Stephen (1997). Latitudinal variation in benthic processes in the abyssal equatorial Pacific: control by biogenic particle flux. *Deep-Sea Research II*, 44(9-10): 2295-2317.
- Smith, F. A. U., & Witman, J. D. (1999). Species diversity in subtidal landscapes: maintenance by physical processes and larval recruitment. *Ecology*, 80(1), 51-69.
- Smith, K. A., & Otway, N. M. (1997). Spatial and temporal patterns of abundance and the effects of disturbance on under-boulder chitons. *Molluscan Research.*, 18, 43-57.
- Snelgrove, P. V. R. (2001). Diversity of Marine Species. *Encyclopedia of Ocean Sciences* (pp. 748-757). Oxford: Academic Press.
- Snelgrove, P. V. R., & Butman, C. A. (1994). Animal-sediment relationships revisited: cause versus effect. *Oceanography and Marine Biology, An Annual Review*, 32, 111-177.
- Sousa, W. P. (2001). Natural disturbance and the dynamics of marine benthic communities. In: M. D. Bertness, S. D. Gaines, & M. E. Hay, *Marine Community Ecology* (pp. 85-132). Sunderland MA: Sinauer Associates, Inc.
- Spalding, M. D. (2007). Marine ecoregions of the World: a bioregionalisation of coastal and shelf areas. *Bioscience*, 57, 573-583.
- St-Onge, P., & Miron, G. (2007). Effects of current speed, shell length and type of sediment on the erosion and transport of juvenile softshell clams (*Mya arenaria*). *Journal of Experimental Marine Biology and Ecology*, 349(1), 12-26.
- Stehli, F. G., McAlester, A. L., & Helsley, C. E. (1967). Taxonomic diversity of recent bivalves and some implications for geology. *Geological Society of America Bulletin*, 78, 455-466.
- Stevens, T., & Connolly, R. M. (2004). Testing the utility of abiotic surrogates for marine habitat mapping at scales relevant to management. *Biological Conservation*, 119(3), 351-362.
- Strathmann, M. F., & Strathmann, R. R. (2007). An extraordinarily long larval duration of 4.5 years from hatching to metamorphosis for teleplanic veligers of *Fusitriton oregonensis*. *Biological Bulletin*, 213(2), 152-159.
- Suetsugu, K., & Ohta, S. (2005). Day and nighttime changes in species composition of deep-sea demersal fishes. *Journal of Oceanography*, 61(2), 187-196.

- Svensson, J. R., Lindegarth, M., Siccha, M., Lenz, M., Molis, M., Wahl, M., & Pavia, H. (2007). Maximum species richness at intermediate frequencies of disturbance: Consistency among levels of productivity. *Ecology*, 88(4), 830-838.
- Taghon, G. L. (1982). Optimal foraging by deposit-feeding invertebrates: roles of particle size and organic coating. *Oecologia Berlin*, 52(3), 295-304.
- Taghon, G. L. (1988). The benefits and costs of deposit feeding in the polychaete *Abarenicola pacifica*. *Limnology and Oceanography*, 33(5), 1166-1175.
- Tett, P., Gowen, R., Mills, D., Fernandes, T., Gilpin, L., Huxham, M., Kennington, K., Read, P., Service, M., Wilkinson, M., & Malcolm, S. (2007). Defining and detecting undesirable disturbance in the context of marine eutrophication. *Marine Pollution Bulletin*, 55(1-6), 282-297.
- Thackway, R., & Cresswell, I. D. (1992). *Environmental Regionalisations of Australia: a User Oriented Approach*. Canberra, Environmental Resources Information Network, Australian National Parks and Wildlife Service.
- Thackway, R., & Cresswell, I. D. (1995). *An Interim Biogeographic Regionalisation for Australia: a Framework for Establishing the National System of Reserves, Version 4*. Canberra, Australian Nature Conservation Agency.
- Thiyagarajan, V., L. Soo, & Qian, P.-Y. (2005). The role of sediment organic matter composition in larval habitat selection by the polychaete *Capitella* sp I. *Journal of Experimental Marine Biology and Ecology* 323, 70-83.
- Thorson, G. (1956). Marine level-bottom animal communities of recent seas. *Journal of Paleontology*, 30(4), 1001-1002.
- Thorson, G. (1957). Bottom Communities. In: Hedgpeth, Joel W. (editor). *Memoirs of the Geological Society of America*, 67(1), 461-534.
- Thouzeau, G., Robert, G., & Ugarte, R. (1991). Faunal assemblages of benthic megainvertebrates inhabiting sea scallop grounds from eastern Georges Bank, in relation to environmental factors. *Marine Ecology-Progress Series*, 74(1), 61-82.
- Thrush, S. F., Hewitt, J. E., Herman, P. M. J., & Ysebaert, T. J. (2005). Multi-scale analysis of species-environment relationships. *Marine Ecology Progress Series*, 302, 13-26.
- Tissot, B. N., Yoklavich, M. M., Love, M. S., York, K., & Amend, M. (2006). Benthic invertebrates that form habitat on deep banks off southern California, with special reference to deep sea coral. *Fishery Bulletin*, 104(2), 167-181.
- Tolimieri, N. (1998). The relationship among microhabitat characteristics, recruitment and adult abundance in the stoplight parrotfish, *Sparisoma viride*, at three spatial scales. *Bulletin of Marine Science*, 62, 253-268.
- Tsuchiya, M., & Nishihira, M. (1986). Islands of *Mytilus edulis* as a habitat for small intertidal animals - effect of *Mytilus* age structure on the species composition of the associated fauna and community organization. *Marine Ecology-Progress Series*, 31(2), 171-178.
- Turner, J. T. (2002). Zooplankton fecal pellets, marine snow and sinking phytoplankton blooms. *Aquatic Microbial Ecology*, 27(1), 57-102.

- Ugland, K. I., Gray, J. S., & Ellingsen, K. E. (2003). The species-accumulation curve and estimation of species richness. *Journal of Animal Ecology*, 72(5), 888-897.
- Urbanski, J. A., & Szymelfenig, M. (2003). GIS based mapping of benthic habitats. *Estuarine, Coastal and Shelf Science*, 56, 99-109.
- Van Dover, C. (2000). *The ecology of deep-sea hydrothermal vents*. Princeton, NJ: Princeton University Press.
- Vasslides, J. M., & Able, K. W. (2008). Importance of shoreface sand ridges as habitat for fishes off the northeast coast of the United States. *Fishery Bulletin*, 106(1), 93-107.
- Veron, J. E. N. (1995). *Corals in Space and Time. The Biogeography and Evolution of the Scleractinia*. Sydney: University of New South Wales Press.
- Vetter, E. W. (1995). Detritus-based patches of high secondary production in the nearshore benthos. *Marine Ecology Progress Series* 120, 251-262.
- Vierros, M., Cresswell, I., Briones, E. E., Rice, J., & Ardrón, J. e. (2008). *Draft Report on Global Open Oceans and Deep Sea-Habitats (GOODS) Bioregional Classification*. Rome, United Nations Environmental Programme. UNEP/CBD/SBSTTA/13/INF/19.
- Volkov, I., Banavar, J. R., Hubbell, S. P., & Maritan, A. (2003). Neutral theory and relative species abundance in ecology. *Nature*, 424, 1035-1037.
- Wakeham, S.G., Lee, C., Hedges, J.I., Hernes, P.J., Peterson, M.L. 1997. Molecular indicators of diagenetic status in marine organic matter. *Geochimica et Cosmochimica Acta*, 61(24), 5363-5369.
- Walker, B. K., Riegl, B., & Dodge, R. E. (2008). Mapping coral reef habitats in southeast Florida using a combined technique approach. *Journal of Coastal Research*, 24(4), 1138-1150.
- Wentworth, C. K. (1922). A scale of grade and class terms for clastic sediments. *Journal of Geology*, 30, 377-392.
- Wheatcroft, R. A., Smith, C. R., & Jumars, P. A. (1989). Dynamics of surficial trace assemblages in the deep sea. *Deep Sea Research*, 36, 71-91.
- Whitlatch, R., B. (1981). Animal-sediment relationships in inter-tidal marine benthic habitats: some determinants of deposit feeding species diversity. *Journal of Experimental Marine Biology & Ecology*, 53, 31-45.
- Whitmire, C. E., Embley, R. W., Wakefield, W. W., Merle, S. G., & Tissot, B. N. (2007). A quantitative approach for using multibeam sonar data to map benthic habitats. In: B. J. Todd, & H. G. Greene, *Mapping the Seafloor Habitat Characterization*. (pp. 111-126). St Johns, Canada: Geological Association of Canada.
- Widdicombe, S., Kendall, M. A., & Parry, D. M. (2003). Using the surface-features created by bioturbating organisms as surrogates for macrofaunal diversity and community structure. *Vie Et Milieu-Life and Environment*, 53(4), 179-186.

- Widdows, J., Brinsley, M. D., Salkeld, P. N., & Elliott, M. (1998). Use of annular flumes to determine the influence of current velocity and bivalves on material flux at the sediment-water interface. *Estuaries*, *21*, 552-559.
- Wildish, D. J., & Kristmanson, D. (1997). *Benthic Suspension Feeders and Flow*. New York: Cambridge University Press.
- Williams, A., & Bax, N. J. (2001). Delineating fish-habitat associations for spatially based management: an example from the south-eastern Australian continental shelf. *Marine and Freshwater Research*, *52*(4), 513-536.
- Williams, A., Gowlett-Holmes, K., & Althaus, F. (2006). Biodiversity Survey of Seamounts & Slopes of the Norfolk Ridge and Lord Howe Rise: Final Report to the Department of the Environment and Heritage (National Oceans Office). Hobart: CSIRO, 591 pp.
- Willig, M. R., Kaufman, D. M., & Stevens, R. D. (2003). Latitudinal gradients of biodiversity: pattern, process, scale and synthesis. *Annual Review of Ecology and Systematics*, *34*, 273-309.
- Wilson, E. O. (1994). *The Diversity of Life*. London: Penguin.
- With, K. A. (1997). The application of neutral landscape models in conservation biology. *Conservation Biology*, *11*(6), 1169-1080.
- Witman, J. D. (1985). Refuges, biological disturbance, and rocky subtidal community structure in New England. *Ecological Monographs*, *55*(4), 421-445.
- Wlodarska-Kowalczyk, M., & Kedra, M. (2007). Surrogacy in natural patterns of benthic distribution and diversity: selected taxa versus lower taxonomic resolution. *Marine Ecology-Progress Series*, *351*, 53-63.
- Wolanski, E. (2001). *Oceanographic Processes of Coral Reefs: Physical and Biological Links in the Great Barrier Reef*. Boca Raton: CRC Press.
- Yeom, D.-H., & Adams, S. M. (2007). Assessing effects of stress across levels of biological organization using an aquatic ecosystem health index. *Ecotoxicology and Environmental Safety*, *67*(2), 286-295.